

# GAC–MAC 2013: FIELD GUIDE SUMMARY

## Ordovician–Silurian Boundary Interval in the Williston Basin Outcrop Belt of Manitoba: A Record of Global and Regional Environmental and Biotic Change

GAC–MAC Winnipeg 2013,  
post-meeting field trip

Robert Elias<sup>1</sup> and Graham Young<sup>2</sup>

<sup>1</sup>Department of Geological Sciences  
University of Manitoba  
Winnipeg, MB, Canada, R3T 2N2  
E-mail: eliasrj@cc.umanitoba.ca

<sup>2</sup>The Manitoba Museum  
190 Rupert Avenue  
Winnipeg, MB, Canada, R3B 0N2

### FIELD TRIP OBJECTIVES

During latest Ordovician to earliest Silurian time, deposition and biotas were influenced globally by environmental changes that led to glaciation and subsequent deglaciation in south-polar Gondwana. At the centre of equatorial Laurentia, the Williston Basin provides a record of global and regional change in an epicontinental sea. Classic localities in southern Manitoba will be compared with sites north of Grand Rapids and northeast of The Pas, including the only known exposures of the Ordovician – Silurian boundary in this vast basin. New research, integrating lithostratigraphy, biostratigraphy, and C-isotope stratigraphy, suggests that deposits of latest Ordovician (Hirnantian) age are present and that the Ordovician – Silurian boundary is at a higher position than previously thought.

Strata exposed in southern and

west-central Manitoba provide a superb opportunity to examine this extraordinary time in Earth's history. This field trip will focus on the Stony Mountain, Stonewall (Figs. 1, 2), and Fisher Branch formations. These predominantly carbonate strata record environments and events leading up to and following the major Hirnantian glacial episode. The succession of fossils represents the culmination of the 'Great Ordovician Biodiversification Event', the Late Ordovician mass extinction, and the Early Silurian recovery. The Williston Basin outcrop belt in Manitoba provides a dip section, from relatively basinal deposits in the south to more marginal in the north. On Day 1, we will visit the type sections of the three formations in the southern part of the outcrop belt. Exposures of correlative strata in two areas within the northern part of the outcrop belt will be examined on Days 2 and 3.

This field trip complements several GAC–MAC 2013 Special Sessions, including *Williston Basin and Other Intracratonic Basins – Sedimentology, Stratigraphy, Paleontology and Resources*; *Advances in Phanerozoic Geoscience in the Hudson Bay Basin, Foxe Basin and Hudson Strait*; and *Evolving Carbonate Environments from Precambrian to Modern: Sedimentological*



**Figure 1.** Type section of the Stonewall Formation at Stonewall Quarry Park, Stonewall, MB.



**Figure 2.** Upper portion of the Stonewall Formation, including the reddish *t*-marker bed, north of Grand Rapids, MB.

*and Geochemical Perspectives*; and the symposium on *Life and Times of Phanerozoic Seas (in Honour of Rolf Ludvigsen)*.

### OTHER INFORMATION

The round trip by minivans in southern and west-central Manitoba will be about 1700 km, with stops at easily accessible roadcuts, outcrops, and abandoned quarries. The longest round-trip hike will be about 1 or 2 easy km. We will be visiting these sites over three days beginning on May 25, so there will be substantial driving time and each day will be rather long. The trip will end late on May 27, so do not plan subsequent travel on that day.