


SERIES



Classic Rock Tours 2. Exploring a Famous Ophiolite: A Guide to the Bay of Islands Igneous Complex in Gros Morne National Park, Western Newfoundland, Canada

Andrew Kerr

*Memorial University
Department of Earth Sciences
St. John's, Newfoundland and Labrador, A1B 3X5, Canada
E-mail: akerr@mun.ca*

SUMMARY

Ophiolites are complex assemblages of ultramafic and mafic igneous rocks that are now widely considered to be pieces of ancient oceanic crust that were emplaced on to the continents courtesy of global plate tectonics. However, most examples were originally considered parts of enormous layered mafic intrusions and so were interpreted in that light. The new understanding of ophiolites in the late 1960s and early 1970s was a crucial part of the global Earth Science revolution, and they are now central to all plate tectonic models developed for ancient orogenic belts. Although their equivalence to oceanic crust is now well established, many ophiolites may not be 'typical' examples of such, and not all examples are identical. Most ophiolites likely formed in subduction-influenced environments rather than at mid-ocean ridges. Ophiolites remain

important foci for research in the 21st century, and many questions remain about their environments of formation and especially their mechanisms of emplacement onto the continents.

Although it was not the first to be seen as a relic of a vanished ocean, the Bay of Islands Igneous Complex in western Newfoundland is one of the best preserved and most easily accessible ophiolites in the world. In the late 20th century, research work in this area proved highly influential in understanding the oceanic crust, and in unravelling the diachronous events involved in the progressive destruction of an ancient stable continental margin as arcs and microcontinental blocks were accreted along it. Parts of the Tablelands Ophiolite lie within Gros Morne National Park, which is a UNESCO world heritage site because of its importance to our understanding of global tectonics. The wider region around the park also includes the Cabox Aspiring Geopark Project, now also in the process of seeking recognition through UNESCO.

This article provides background information on ophiolites and the development of our ideas about them, and links this material to four self-guided field excursions that allow examination of many classic features. These excursions range from a collection of roadside outcrops, to some relatively easy hiking excursions on official National Park trails, and eventually to a more challenging off-trail hike that ascends to the summit plateau of the Tablelands to visit rare exposures of the Moho (the Mohorovičić Discontinuity, i.e. the lower boundary of the Earth's crust) and the underlying upper mantle rocks. Collectively, the field stops should allow geologically-minded visitors to experience some amazing geology in a spectacular and sometimes surreal landscape.

RÉSUMÉ

Les ophiolites sont des assemblages complexes de roches ignées ultramafiques et mafiques qui sont maintenant généralement considérées comme des fragments de croûte océanique ancienne qui ont été charriés sur les continents grâce à la tectonique globale des plaques. Cependant, la plupart des exemples étaient à l'origine considérés comme faisant partie de vastes intrusions mafiques stratifiées et ont donc été interprétés dans ce contexte. La nouvelle compréhension des ophiolites à la fin des années 60 et au début des années 70 a été un élément crucial de la révolution des sciences de la Terre. Les ophiolites sont désormais au cœur de tous les modèles tectoniques des plaques développés pour les anciennes ceintures orogéniques. Bien que leur équivalence avec la croûte océanique soit maintenant bien établie, de nombreuses ophiolites

lites peuvent ne pas en être des exemples « typiques », et tous les exemples ne sont pas identiques. La plupart des ophiolites se sont probablement formées dans des environnements influencés par la subduction plutôt qu'au niveau des dorsales océaniques. Les ophiolites restent un thème de recherche important au XXI^e siècle et de nombreuses questions subsistent quant à leurs environnements de formation et notamment à leurs mécanismes de mise en place sur les continents.

Bien qu'il n'ait pas été le premier à être identifié comme un vestige d'un océan disparu, le complexe igné de la baie des Îles, dans l'ouest de Terre-Neuve, fait partie des ophiolites les mieux conservées et les plus facilement accessibles au monde. À la fin du XX^e siècle, les travaux de recherche dans ce domaine ont joué un rôle déterminant dans la compréhension de la croûte océanique et dans la compréhension des événements diachrones impliqués dans la destruction progressive d'une ancienne marge continentale stable au fur et à mesure de l'accrétion d'arcs et de blocs microcontinentaux. Une partie des Tablelands Ophiolite se trouve dans le parc national du Gros-Morne, site classé au patrimoine mondial de l'UNESCO en raison de son importance pour notre compréhension de la tectonique globale. La région plus large autour du parc comprend également le projet Cabox Aspiring Geopark, qui est également à la recherche d'une reconnaissance dans le cadre de l'UNESCO.

Cet article fournit des informations de base sur les ophiolites et le développement de nos idées à leur sujet, et relie ce matériel à quatre excursions autoguidées qui permettent d'examiner de nombreuses caractéristiques classiques. Ces excursions vont d'une collection de visites d'affleurements au bord de la route, à des randonnées relativement faciles sur les sentiers officiels du parc national, et finalement à une randonnée plus difficile hors-piste menant au plateau sommital des Tablelands pour visiter de rares affleurements du Moho (la discontinuité de Mohorovičić, c'est-à-dire la limite inférieure de la croûte terrestre) et des roches du manteau supérieur sous-jacentes. Collectivement, les visites sur le terrain devraient permettre aux visiteurs amateurs de géologie de faire l'expérience d'une géologie remarquable dans un paysage spectaculaire et parfois surréaliste.

Traduit par la Traductrice

PREAMBLE

In the late 1960s, ideas long held by geologists about stratigraphy, mountain belts, the origins of igneous rocks and many other concepts were revised and in some cases entirely reversed. Unusual mafic and ultramafic 'intrusions', which had long puzzled both mapmakers and petrologists, assumed greater importance when it was first proposed that they might be remnants of the vanished oceans that global tectonics predicted. The term *ophiolite* is now almost as familiar as 'basalt' to undergraduates, but few of them realize that its use long predates ideas of continental drift, or that these enigmatic rocks took so long to understand. Most of the ophiolites around the world were once studied largely by petrologists and mineralogists. These vast igneous massifs were logically interpreted as large layered intrusions that were younger than the surround-




Figure 1. The location of Gros Morne National Park and the adjoining Cabox Aspiring Geopark areas in western Newfoundland. The map also shows the fundamental three-fold division of Newfoundland into the Humber Zone (the ancient continental margin of Laurentia), the Avalon Zone (Late Precambrian rocks that correlate with similar rocks in Europe and North Africa) and the Central Mobile Belt (Dunnage and Gander Zones, which represents island arcs and other regions formed within the proto-Atlantic or Iapetus Ocean). Terminology for the Appalachian Orogen in Newfoundland after Williams (1979, 1995).

ing rocks, even if the contact relationships were enigmatic. The evidence for the true nature of ophiolites was, of course, always there for us to record but it was discounted because the implications extended beyond prevailing concepts. Such is often the case in the history of geology.

There are many well-established (but variably understood) ophiolites around the world, and the list continues to expand (e.g. Dilek and Furnes 2011, 2014). Ophiolites are key elements in our interpretations of ancient orogenic belts and the development of tectonic models for them (e.g. Dalziel et al. 2000). The ultramafic rocks within ophiolites commonly form unusual and striking landscapes, so it is not surprising that several important examples are within National Parks or other reserves. Gros Morne National Park in western Newfoundland (Fig. 1) includes parts of the Cambrian–Ordovician Bay of Islands Igneous Complex (BOIC), which was first recognized as an ophiolite in the late 1960s. Western Newfoundland is a key location where ideas linking ancient rocks and the modern oceanic crust were first integrated (e.g. Stevens 1965; Church and Stevens 1971; Williams and Stevens 1974). The BOIC was not the first ophiolite to be interpreted in this way, but it has proved to be one of the most influential, and it is certainly one of the most spectacular. The ophiolites are one of the primary reasons why Gros Morne National Park is now designated as a UNESCO World Heritage Site. Beyond the National Park boundaries, the BOIC also forms the core of the Cabox 'Aspiring Geopark' proposal, also under the broad umbrella of UNESCO (see www.caboxgeopark.org for more information). Ophiolite enthusiasts undoubtedly have their own favourites (e.g. Bédard 2014), but most would agree that western New-

foundland is one of the best places on Earth to explore an ophiolite and understand the story that it tells. This article is intended to help make that process easier for geologically-minded visitors to this now-famous area.

This article does not pretend to be a specialized research contribution, but instead sets out to review background information and safely guide those who wish to independently explore and appreciate this influential 'type area'. Much of the material herein is derived from previous field trip guides and articles of a more specialized nature (e.g. Malpas 1987; Calon et al. 1988; Williams and Cawood 1989; Berger et al. 1992; Cawood and Suhr 1992; Stevens et al. 2003). Additional information, and most photographs, are from many personal visits and a long-dormant project with the late Robert K. Stevens to write a more comprehensive geological guidebook to the National Park. The numerous publications referenced in the text should remain the primary citations for details of the geology, rather than this more generalized overview. The article is intended to be understood by readers who lack specialized knowledge of tectonic processes or igneous rocks, but it assumes a general understanding of geological terminology and concepts appropriate to introductory Earth Science courses. A short glossary is provided in the appendix to explain specific terms that may be unfamiliar, and one of the many published dictionaries of geology might also prove useful. For those with minimal knowledge of geology, the excellent book by Hild (2012) covers parts of Gros Morne, including the Tablelands and provides information intended for a wider non-scientific audience.

OPHIOLITE ESSENTIALS

The Development of the Concept

The term *ophiolite* long predates the plate tectonics revolution. It is derived by combining *ophio-* (Greek for 'snake') and *-lite* (a Greek suffix for 'stone'), so its original meaning is *snakestone*. This curious name reflects the abundance of serpentinites - as you might guess, serpentine minerals were so named because their weathering textures resemble snakeskins. Ophiolite was an early descriptive term used for mafic and ultramafic rocks, commonly altered and variably deformed, that were first described within the Italian Alps (Brongniart 1821). The later term *Steinmann Trinity* captured the common association of such serpentinites with altered basalts ('spilites') and cherts - a threesome usually found together somewhere within ancient orogenic belts. Steinmann (1927) noted the deep-water character of the laminated cherts, and implied a link between these three rock types and the deep ocean floor. Interestingly, the area discussed by Steinmann is a rather atypical ophiolite, as it is missing some characteristic components (e.g. Bernoulli 2001; Desmurs et al. 2001). For over a century, ophiolites were interpreted as large intrusions, but they invariably had ambiguous relationships with surrounding sedimentary and volcanic rocks. The lack of large contact metamorphic aureoles around them seemed inconsistent with high-temperature mafic or ultramafic magmas, and it was hard to confirm that they intruded their country rocks. Also, prevailing ideas about

igneous processes (e.g. Wager and Deer 1939) suggested that such ultramafic rocks had to be derived by processes of crystallization and mineral accumulation within much larger mafic magma chambers, which never seemed to be preserved in the vicinity. They were truly a puzzle.

The idea that ophiolites might be *tectonically* emplaced in a solid state was suggested first for the California Coast Ranges, and then for similar bodies in the South Island of New Zealand (e.g. Coleman 1967, 1971). However, the idea that they might originate elsewhere did not solve the puzzle, because it was not clear where 'elsewhere' might be. Prior to the oceanic drilling and geophysical programs of the 1960s, geologists had limited knowledge about oceanic crust and little inkling of the many ways in which continents and oceans could interact. A new interpretation of ophiolites emerged in the light of clearer evidence for the mobility of continents and the transience of oceans - they were seen as slices of oceanic crust and underlying mantle. Although the first such musings came from California, interpretation of the Troodos Mountains (Cyprus) as uplifted oceanic crust (Gass 1968) first brought the idea into the scientific mainstream. Coleman (1971) first coined the term *obduction* for the process by which ophiolites are tectonically emplaced, but the exact details of this process are still open to debate, as discussed in a later section.

With new interpretations of Mesozoic ophiolites in hand, it was logical to think that Paleozoic examples might also be relics of long-vanished oceans. If so, ophiolites could provide key evidence for the operation of global plate tectonics through Earth's history. The Bay of Islands Igneous Complex in western Newfoundland (Fig. 1) was one of the first to be reinterpreted (Stevens 1965; Church and Stevens 1971; Dewey and Bird 1971), and it remains a classic area. Western Newfoundland (and Gros Morne National Park in particular) has wider importance in plate tectonics, because the rocks reveal how various components of an ancient continental margin were transported and juxtaposed as island arcs, and microcontinental blocks were progressively 'accreted' to North America from the adjoining Iapetus Ocean (see later discussion for the origins of this name). The spectacular ultramafic rocks that symbolize Gros Morne National Park are the best-known part of this story, but they are only one chapter in a much longer tale.

To say that ophiolites are critical to understanding Earth history is a huge understatement. Almost all Paleozoic orogenic belts contain ophiolites, although their preservation varies widely, but in all cases they constrain plate-tectonic models. Ophiolites of Precambrian age remain elusive, but several are now widely accepted, including the well-preserved Paleoproterozoic Purtiniq ophiolite of northern Quebec (e.g. Scott et al. 1992). Ophiolites remain important research topics today, and there is even a scientific journal completely devoted to their study (*Ophioliti*, published in Italy; www.ofioliti.it). My general article and guide simply cannot do justice to decades of study and many remaining questions. Several papers in the journal *Elements* (2014, volume 10, issue 2) provide reviews with extensive bibliographies. An earlier GSA Special Paper






Figure 2. The major components of the ophiolite suite, their interpretation and their relationship to layer structure defined in the modern oceanic crust by seismic and other geophysical methods. Modified after Williams and Malpas (1972) with specific reference to the Bay of Islands Igneous Complex, but also applicable elsewhere. More detailed views of various types of ophiolites are provided by Dilek et al. (2000) and Dilek and Furnes (2014).

(Dilek et al. 2000) provides a wealth of more detailed information on specific examples. The following sections summarize some key aspects, with specific application to the western Newfoundland examples highlighted in this article.

Components of Ophiolites and Links to Modern Oceanic Crust

Although terminology varies, seven principal components are recognized within ophiolites and equated with specific parts of the oceanic crust (Fig. 2 after Williams and Malpas 1972; Dilek and Furnes 2011, 2014), even though our *direct* knowledge of the latter is limited to the uppermost 2 km or so. From top to bottom, these are:

1. Sedimentary rocks (typically cherts or shales) that were deposited on the lavas forming the deep-seafloor.
2. Pillow lavas of basaltic composition that were extruded to form the seafloor.
3. 'Sheeted dykes' of basaltic (diabase) composition, which are interpreted as the feeder systems to the overlying pillow lavas.
4. Gabbroic rocks representing high-level fractionated magma chambers, commonly gradational with the overlying sheeted diabase dyke systems.
5. Layered gabbros, which record cumulate processes in the deeper sections of mid-crustal magma chambers.
6. Ultramafic cumulate rocks, notably feldspathic dunite and olivine-rich gabbro, which define the transition between crust and mantle, i.e. the Moho, which is shorthand for 'Mohorovičić Discontinuity'.
7. Massive (but commonly deformed) ultramafic rocks (harzburgite, dunite and pyroxenite) that represent the

residual mantle material from partial melting that produced the overlying oceanic crust section.

Geophysical studies of the modern oceanic crust indicate that it is layered, and four layers are generally defined on the basis of its seismic velocity structure. The seven components listed above for ophiolites are broadly equated with these seismically-defined layers (Layers 1 to 4; Fig. 2), but they do not necessarily have individual seismic expression. The original definitions suggested from a landmark Penrose conference (Penrose Field Conference Participants 1972) excluded the uppermost sedimentary rocks, and it is now recognized that modern oceanic crust may lack parts of layer 2 (pillow basalts and sheeted dykes) and layer 3 (gabbro) in areas of strong extension. Such is apparently the case for the ophiolitic rocks that first defined Steinmann's trinity (Desmurs et al. 2001). Ancient ophiolites are rarely complete, as they may be structurally dismembered during emplacement and/or truncated by later erosion. The dyke- and basalt-dominated upper parts of the ophiolite assemblage are the most vulnerable to removal by both mechanisms, and may be absent. Ultramafic rocks representing the upper mantle are more robust, although they may be variably serpentinized. Subsea hydrothermal systems cause extensive alteration, and emplacement involves low-grade metamorphism, so the original features of many ancient ophiolites are cryptic.

Although not specifically described above, many ophiolites also contain minor quantities of granitoid rocks, which are typically plagioclase-rich (albite) tonalites, also known as *trondhjemites*. These are interpreted as residual magmas from deeper fractionating magma chambers, or as products of localized partial melting within the crust (e.g. Malpas 1979; Jenner et al. 1991; Dilek and Furnes 2014). Intermediate and felsic magmas of trondhjemite type are also known in modern arc-type settings (Dilek and Furnes 2014), and their presence in ancient ophiolites is important in understanding their tectonic settings.

Some ophiolite complexes, including the BOIC, also contain discontinuous belts of metamorphic rocks that are located structurally *below* the ultramafic rocks. These are commonly of mafic composition, and are interpreted as thermal and tectonic aureoles developed during obduction. These metamorphic aureoles provide very important evidence for the tectonic emplacement of ophiolites and their high ambient temperatures compared to structurally underlying rocks. The metamorphic aureole of the BOIC was one of the first to be recognized and interpreted in this fashion (Williams and Smyth 1973).

Ophiolites host specific styles of economic mineralization although these are not in themselves diagnostic features (e.g. Jébrak and Marcoux 2015). Examples include sea-floor sulphide deposits (volcanogenic massive sulphides, or VMS) in volcanic rocks, and podiform to massive chromite accumulations in peridotites. Chromite occurrences are typically small, but many ophiolites were also explored for stratiform chromitites and platinum-group element deposits because the mafic and ultramafic rocks were considered to represent parts of large layered mafic intrusions, which host important deposits of both types. Important asbestos deposits occur in many

ophiolites, formed by syn- and post-emplacement alteration of the ultramafic rocks; notable Canadian examples include the Baie Verte area in Newfoundland and the Thetford Mines area in Quebec (Jébrak and Marcoux 2015). Diamonds are reported from some ophiolite complexes (e.g. Yang et al. 2014), although such concentrations are not thought to be economically significant.

Many studies over many years have outlined the strong similarities between ancient ophiolites and the modern oceanic crust in terms of rock types, composition and structure. The comparison is not perfect in all details for all examples, but the discrepancies lie mostly in the relative abundance of rock types, or petrological details, rather than gross anatomy. Our *direct* knowledge is probably more complete for ancient ophiolites than for the modern oceanic crust, and we cannot be certain that original relative thicknesses are retained following their obduction. More importantly, many ophiolite sequences probably represent substrates to island arcs rather than true mid-ocean-ridge settings (see later discussion) and such arc environments are intrinsically more varied in character. Notwithstanding continuing discussion as to exactly *where* individual ancient ophiolites were formed and *how* they were emplaced, the link to the crust of the modern oceans remains convincing, and such a conclusion is also supported by the similarities of seismic profiles obtained from the BOIC (and other ophiolites) to those from the modern ocean basins (e.g. Salisbury and Christensen 1978).

The Varieties and Flavours of Ophiolites

Ophiolites are diverse in character, with contrasts in their regional geological settings, and more subtle geochemical and isotopic differences. Recent reviews (e.g., Dilek and Furnes 2014; Pearce 2014) define several major categories, and some of these can be subdivided according to their finer details.

The clearest distinction is into so-called 'Tethyan' and 'Cordilleran' types, which have different geological settings. The Mesozoic *Tethyan* ophiolites, including the Troodos complex in Cyprus, and the vast Semail ophiolite in Oman (considered to be the largest known ophiolite; e.g. Searle and Cox 1999; Goodenough et al. 2014) are regionally associated with shallow-water platformal sedimentary sequences. The magmatic stratigraphy of these ophiolites is relatively intact, and they seem to have been transported across passive continental margins. The Paleozoic ophiolites of the Appalachians, although unrelated to the much younger Tethys Ocean, are also of this general type. In contrast, *Cordilleran* ophiolites, exemplified by those of western North America and the wider circum-Pacific region, lack such association with platformal sequences, and are associated with deep-water clastic rocks (turbidites) and low-temperature, high-pressure metamorphism, including blueschists. Their regional tectonic association is with long-lived convergent margins and active subduction zones rather than previously stable continental shelves. Cordilleran ophiolites tend also to be more dismembered and/or deformed, and are typically more altered than their Tethyan counterparts. Ophiolites defined in ancient orogenic belts include examples of both types, but the best-known

examples generally correspond to the Tethyan type. Paleoproterozoic ophiolites in northern Quebec (e.g. Scott et al. 1992) likely also belong in this category.

Geochemical data indicate that although there are a few ancient and modern ophiolites that have mid-ocean-ridge basalt (MORB) affinities, most were not formed in such settings. Miyashiro (1972) pointed out that basalts from the Troodos ophiolite in Cyprus had arc-like geochemical traits, suggesting formation in an island arc or back-arc basin, rather than their proposed ridge-type setting (e.g. Gass 1968). This seems to be a general trait, although some larger ophiolites (e.g. Semail in Oman) may record both early ridge-type and later island-arc-type components (e.g. Pearce 2014; Goodenough et al. 2014). The geochemical patterns of ophiolites also define several subtypes (e.g. Dilek and Furnes 2014; Pearce 2014) but this geochemical variation appears to be independent of the more fundamental division into Tethyan and Cordilleran types. Most ophiolites are now considered to have formed in fore-arc or back-arc settings. From the perspective of a field geologist the rocks from these various ophiolite flavours look essentially the same, but the inference of arc-related settings is very important for ideas about the mechanisms of emplacement, as discussed below.

The Emplacement of Ophiolites

Density contrasts between broadly granitic continental crust and mafic oceanic crust indicate that *obduction* of the latter, rather than its subduction, demands special circumstances. Under normal circumstances, dense ocean crust should descend below less dense continental crust, as in subduction zones. However, the attempted subduction of continental crust during terrane accretion or continental collision provides a mechanism by which buoyant continental crust could 'rebound', carrying with it parts of the upper oceanic plate. In simple terms, trying to subduct continental crust beneath oceanic crust is akin to trying to push a buoyant inflatable cushion beneath the water in a swimming pool. This general premise is central to most models for ophiolite emplacement (e.g. Searle and Stevens 1984; Dilek and Furnes 2014). Tethyan-type ophiolites would be emplaced across continental margins when the latter enter a subduction zone that dips *towards* the adjoining ocean, rather than beneath the continent (Fig. 3). Models of this type also explain stratigraphic evolution from a stable shallow-water sequence to deeper-water clastic sedimentation as the volcanic arc approaches and sheds its detritus over the foundering continental shelf (e.g. Church and Stevens 1971; Williams and Stevens 1974; Cawood and Suhr 1992). Intervening sedimentary environments from the outer continental shelf and the continental slope are then assembled tectonically ahead of and below the oceanic crustal slab as it approaches and sheds detritus across them, and are progressively transported across the former continental shelf. The end result is an ordered sequence of tectonic slices in which the distance of tectonic transport increases from bottom to top, culminating in the ophiolite section. The actual assembly and transport of all these components is diachronous, and ophiolite emplacement is the final step. In a general






Figure 3. Possible tectonic setting for generation and emplacement of Tethyan-type ophiolites, such as the Bay of Islands Igneous Complex, in which an irregular continental margin approaches a curvilinear subduction zone that dips away from the continent. (A) Ophiolites form in places where oceanic crust remains trapped in embayments (also termed ‘reentrants’) of the continental margin after adjacent promontories (also termed ‘salients’) enter the subduction zone. (B) The tectonic architecture along lines X–X’ and Y–Y’ is shown for Time points 1 and 2. At Time 2, continued subduction of the trapped oceanic crust and fore-arc extension generates new oceanic crust, which is hot and buoyant, and is emplaced onto the continental margin at Time 3 (C), in conjunction with adjoining sedimentary and igneous rock assemblages. For further details and discussion of emplacement mechanisms, see Cawood and Suhr (1992) and Dilek and Furnes (2014). The very schematic diagram is based on models of Cawood (1991) and Cawood and Suhr (1992).

sense, this is the prevailing model for the emplacement of the BOIC and also for the wider geological history of Gros Morne National Park.

Cawood and Suhr (1992) suggested that the generation of the ophiolites and their eventual emplacement both resulted from the irregular geography of the ancient continental margin compared to that of the offshore island arc (Fig. 3). In this model, promontories along the margin (also known as ‘salients’) will be the first to encounter the subduction zone, but oceanic crust will remain in other areas where there is an embayment in the margin (also known as a ‘reentrant’). In such areas, the oceanic crust continues to subduct, leading to its stretching (extension) and perhaps its eventual detachment from the associated continent, promoting the formation of ‘new’ and hot oceanic crust. It is this newly-formed and relatively buoyant oceanic crust that is eventually obducted onto the approaching continent when remnant basins are finally closed (Fig. 3). In detail, the model is far more complex than these few sentences would suggest, but it is applicable to other ophiolites, such as those of Oman (e.g. Cawood 1991). For further details of other emplacement models and discussions related to other ophiolites, readers should consult Dilek and Furnes (2014) and its extensive bibliography. Although many ideas about ophiolites are now well accepted, there remains

much debate about the exact mechanisms of their emplacement.

GEOLOGICAL FRAMEWORK

Gros Morne in the Context of the Appalachian Orogenic Belt

Gros Morne National Park is located on the west coast of Newfoundland, about 70 km north of Corner Brook (Fig. 1). The island of Newfoundland is part of the Appalachian–Caledonian Orogenic Belt, which contains rocks ranging in age from Precambrian to Carboniferous, recording multiple orogenic events of broadly early to middle Paleozoic age, i.e. Ordovician, Silurian, Devonian and Carboniferous. This large orogenic belt was dispersed when the modern Atlantic Ocean opened during the Mesozoic, and other parts of it now occur in northwestern Europe and Scandinavia, the east coast of Greenland, and in various Arctic islands (Fig. 4).

The Appalachian–Caledonian Orogenic Belt was formed when the continent of North America (Laurentia) was amalgamated with present-day Europe and parts of present-day Africa and South America (part of a larger continental assembly called ‘Gondwanaland’). This took place in several discrete stages, and the first stage involved the closure of a major ocean basin that existed from the late Precambrian to the late Ordovician. This ocean, which all but disappeared in Silurian times, is now called Iapetus, honouring the father of Atlantis in Greek mythology (Harland and Gayer 1972), but the concept of a precursor ocean to the modern Atlantic was made famous by J. Tuzo Wilson, in his classic paper entitled “*Did the Atlantic close and then reopen?*” (Wilson 1966). He noted that because the line along which the modern Atlantic opened was not the same as that along which its predecessor had closed, pieces of Europe and Africa were left attached to North America, and pieces of North America were left attached to Europe (Fig. 4). His analysis was, not surprisingly, strongly influenced by the geology of Newfoundland. Harold (Hank) Williams had earlier published a paper that outlined the concept of a “two-sided symmetrical system” for the Newfoundland Appalachians (Williams 1964). Newfoundland has huge importance in the Appalachian Orogen because it preserves both sides of the belt, and also the largest surviving pieces of the Iapetus Ocean. It also provides a vital link between the Appalachians and the Caledonian orogenic belt of Britain and Scandinavia (Fig. 4).

During the closure of the Iapetus Ocean, several island arcs were developed over subduction zones along both the Laurentian and Gondwanaland margins of Iapetus. In the middle Ordovician (~ 470 m.y. ago), the Laurentian continent collided with an island arc or arcs along a subduction zone that dipped away from the continent. Laurentia, and its stable continental shelf, was pushed underneath rocks of oceanic affinity, including deep-water sedimentary rocks, igneous rocks formed in the island arcs, and the deeper oceanic crust with its underlying mantle. These oceanic rocks were then transported tectonically for hundreds of kilometres towards the interior of Laurentia to form what are generally called the *Taconic*




Figure 4. Reconstruction of the Appalachian-Caledonian Orogen and surrounding areas obtained by closing the modern Atlantic and fitting the surrounding continental shelves. Modified from Williams (1995). Note how some parts of modern Europe (e.g. Scotland, Norway) are of North American affinity, and some parts of modern North America (e.g. parts of Newfoundland and Nova Scotia) are of European or African affinity.

Allochthons, named for an area in upstate New York (Fig. 4). The term 'allochthon' (adjective, allochthonous) refers to a collection of rock units that have moved over large lateral distances during and after tectonic collisions. One of the largest of the Taconic Allochthons underlies much of western New-

foundland, including part of Gros Morne National Park. It is known as the *Humber Arm Allochthon* (e.g. Williams and Cawood 1989; Figs. 4 and 5).

Gros Morne National Park is important because it preserves many of these various tectonic elements in one small area. It contains the Precambrian basement rocks of North America and a relatively complete section through the Cambrian-Ordovician continental shelf sedimentary sequence. Sitting structurally above these are the allochthonous rocks transported from the deeper parts of the continental shelf and the Iapetus Ocean, represented by the Humber Arm Allochthon. The Bay of Islands Igneous Complex is the most famous of these transported components. Other allochthonous components include mafic volcanic rocks and plutonic rocks formed in an island arc, and deep-water sedimentary rocks of both clastic and carbonate affinity. The latter represent the ancient continental slope and adjacent basin, and contain remarkable fossil assemblages that help to constrain geologic time (e.g. James et al. 1989). These rocks are in their own way every bit as unique and spectacular as the ophiolites, although they form less prominent landscapes. They include the Global Stratotype Section for the Cambrian-Ordovician boundary, located at Green Point, also within Gros Morne National Park (Cooper et al. 2001).

The evolution of the Appalachian-Caledonian Orogenic Belt did not end with the events recorded in Gros Morne National Park. Volcanism, plutonism and deformation continued through the Ordovician, the Silurian, and well into the Devonian. Gros Morne National Park largely lay outside the influence of these later events, which is another reason for its importance in understanding early events associated with ophiolites. The final chapter in the biography of the Park was written in the Pleistocene, when the spectacular landscape of today was sculpted by glaciation. The most obvious results of glaciation are the much-photographed fjords such as Bonne Bay and Western Brook Pond, but there are lesser features that attest to the power of the ice sheets. Berger et al. (1992) provide some useful summary information on Quaternary history and landscapes, and references to more detailed sources.

An Overview of the Geology of Gros Morne National Park

Gros Morne National Park may be conveniently divided into eight main geological 'packages' (Figs. 6 and 7; after Williams and Cawood 1989; Colman-Sadd et al. 1990; Berger et al. 1992). Two of these packages are in their original locations (i.e. they are *autochthonous*) and the remainder are allochthonous, but some have travelled further than others have. The order of description below corresponds broadly to their structural position (Fig. 7) rather than to their actual stratigraphic age. Subsequent sections provide more background information on the Little Port Complex and the Bay of Islands Igneous Complex (Fig. 5), which are the main foci of the suggested excursions.

Precambrian Basement Rocks (Long Range Inlier)

The oldest autochthonous rocks in the park are Precambrian gneisses and granites of the Long Range Inlier (Fig. 5), which underlie most of the high, barren land in the west (Fig. 6).




Figure 5. Generalized geology of part of Newfoundland and a more detailed inset view of the ophiolite massifs and related rock units in the Humber Arm Allochthon. Geology after Colman-Sadd et al. (1990), Williams and Cawood (1989), Cawood and Suhr (1992). Note that several other ophiolite sequences exist in central Newfoundland, where they represent parts of the Iapetus Ocean and related basins associated with island arcs.

These rocks were formed well over a billion years ago; they form part of the region of the Canadian Shield known as the Grenville Province. These basement rocks are unconformably overlain by Lower Paleozoic sedimentary rocks in the south of the Park, but are separated from them by west-directed reverse faults in the north (Fig. 7). This west-directed transport and uplift is late (Silurian or Devonian in age) compared to the initial Middle Ordovician emplacement of the allochthons.

Autochthonous Cambrian and Ordovician Sedimentary Rocks

Autochthonous Cambrian and Ordovician strata unconformably overlie the Precambrian basement (Figs. 6 and 7), and record the initiation and development of a stable continental shelf, dominated by carbonate deposition. Gros Morne is not the best place to understand Cambrian and Ordovician stratigraphy, as it has more than a few structural complications, but the succession is relatively complete. The youngest rocks of the autochthonous succession are sandstones that were derived from the east, rather than from continental regions to the west. These youngest sedimentary rocks formed from the erosion of an offshore landmass during early stages of the obduction process (e.g. Church and Stevens 1971; James et al. 1989). This eastward-derived detritus thus records the uplift and erosion of island arcs and ophiolitic rocks during the

attempted subduction of the continental margin, as discussed above.

Allochthonous Cambrian and Ordovician Sedimentary Rocks of the Cow Head Group

The Cow Head Group, in the north of the park (Figs. 6 and 7), comprises carbonate strata formed on the outer continental shelf and the continental slope. This remarkable assemblage of deep-water carbonate rocks is correlated with the autochthonous succession using fossils, and it contains the Global Stratotype Section for the Cambrian–Ordovician boundary located at Green Point (Fig. 6; Cooper et al. 2001). The Cow Head Group represents a variety of coexisting depositional environments, but the carbonate rocks in each structural slice are eventually overlain by deep-water clastic rocks derived from the east. The strata of the Cow Head Group thus record the obduction process in a similar manner to the autochthonous sequence (see above), but the change in the sedimentary environment and provenance occurred at an *earlier* time than in the autochthonous rocks, because the allochthonous sedimentary rocks were originally closer to the subduction zone. The inundation of the continental margin by detritus from the east is thus *diachronous*, i.e. it occurred at different times in different locations.




Figure 6. Simplified geology of Gros Morne National Park and surrounding areas, showing major rock assemblages only. Adapted from Berger et al. (1992), Williams and Cawood (1989).




Figure 7. Schematic west to east cross-sections for the northern and southern areas of Gros Morne National Park, showing the inferred structural relationships between autochthonous basement and cover, and various packages of allochthonous rocks, including the Tablelands Ophiolite. Based on Berger et al. (1992), Williams and Cawood (1989).

Allochthonous Cambrian and Ordovician Sedimentary Rocks of the Bonne Bay Group

In the southern area of Gros Morne National Park, allochthonous Cambrian and Ordovician sedimentary rocks (Fig. 6) are mostly of clastic type rather than carbonate-dominated, and are interbedded with lesser mafic volcanic rocks and some deep-water carbonate rocks. These rocks, generally assigned to the Bonne Bay Group, are interpreted to be derived from more distal, deeper parts of the Iapetus Ocean, perhaps underlain by the transition between continental and oceanic crust. Importantly, some sandstones within this package contain detrital chromite, which is a diagnostic indicator for derivation from ultramafic source regions such as the ophiolite suite (Stevens 1965, 1970).

Allochthonous Late Precambrian basalts of the Skinner Cove Formation

Basaltic rocks of the Skinner Cove Formation occur only in a small area of the park, between Bonne Bay and Trout River (Fig. 6) and were originally thought to represent a seamount or oceanic island formed in the Iapetus Ocean, because of their 'alkaline' geochemical traits (Baker 1979). More recent U–Pb geochronology and paleomagnetic data (Cawood et al. 2001; Hodych et al. 2004) indicate a Late Precambrian age (~ 550 Ma). This suggests that they represent older volcanism that might have been associated with the initial rifting of Laurentia and the establishment of a continental margin, as discussed by Cawood et al. (2001).

Allochthonous Igneous Rocks of the Little Port Complex

In the south of the park, allochthonous sedimentary rocks and alkali basalts are in turn structurally overlain by a complex assemblage of igneous rocks, including peridotite, gabbro, diabase, pillow basalt and trondhjemite (Figs. 6 and 7). These rocks are generally termed the Little Port Complex although the more general term 'Coastal Complex' is applied by some (Casey et al. 1985; see also Fig. 5). These are believed to mostly represent higher parts of the oceanic crust developed in an island-arc type setting within the Iapetus Ocean. Minor sedimentary rocks characteristic of deep-water oceanic environments are also present.

Allochthonous Igneous Rocks of the Bay of Islands Complex (Tablelands Ophiolite)

The structurally highest geological package in Gros Morne National Park comprises the spectacular ophiolites of the Tablelands region, representing mostly the upper mantle and the lower sections of the Paleozoic oceanic crust (Figs. 5 to 7). South of Trout River Pond, the ophiolites of the adjacent North Arm Mountain Massif are downfaulted and tilted, and better reveal higher levels of the oceanic crust (gabbro, sheeted dykes and basalts), and also metamorphic rocks at the base of the complex. The metamorphic aureole to the ophiolite is also exposed along the eastern side of the Tablelands Ophiolite, which represents its base. The metamorphic aureoles were probably developed from mélanges consisting largely of mafic volcanic rocks, as initially discussed by Williams and Smyth

(1973). The various components of the Bay of Islands Igneous Complex are shown in Figure 5.

Mélange Zones

The remaining geological package in Gros Morne consists of rocks formed *during* the tectonic transport of allochthonous rocks over the autochthonous sedimentary sequences. These generally chaotic and confusing rocks are termed *mélanges* (Figs. 6 and 7). The most extensive belt is in the Rocky Harbour area, and consists of a wide variety of coherent blocks, ranging from a few metres to several hundred metres in size, encased in a shaly matrix. The belt continues on the south side of Bonne Bay. Similar (but thinner) mélange zones separate higher structural slices in the southern part of the park. The mélanges developed under generally cool conditions, but high temperatures and pressures prevailed at the base of the ophiolite sequences, where hot mantle peridotites were exhumed (see previous section).

Geometric Arrangement of Autochthonous and Allochthonous Rocks

The general arrangement of the various rock packages described above is illustrated by schematic cross-sections (Fig. 7). Allochthonous rocks always sit structurally above autochthonous rocks, except in the north of the park, where later reverse faults locally place Precambrian basement above the Cow Head Group. In the region south of Bonne Bay, the stacking order is seen most clearly. The allochthonous sedimentary rocks of the Bonne Bay Group form the base of the stack, with the Little Port Complex and the Tablelands ophiolites sitting structurally above them. The Cow Head Group is absent here, but would sit below the Bonne Bay Group. The exact position of the Skinner Cove Formation relative to the Bonne Bay Group is more difficult to establish, although it clearly sits below the Little Port Complex. The sequence from bottom to top within the allochthonous rocks also represents a geographic sequence, in which higher slices travelled progressively greater distances from the east, across the ancient continental margin represented by the autochthonous sequence.

Ophiolites in Western Newfoundland - A Brief Research History

The history of research in the ophiolites of western Newfoundland is an interesting topic in its own right, and cannot be fully explored here. The ultramafic and mafic rocks were noted by early mappers, but little attention was paid to them until the 1930s, when chromite concentrations attracted more attention (e.g. Snelgrove 1934). Other studies during this time were completed by Ingerson (1935), Cooper (1936) and also by a young geologist from Princeton named Harry Hess. Hess would later be known as one of the founding figures in the plate tectonics revolution, and an interesting paper from that period (Hess 1938) revealed that he thought at length about the wider puzzle of the ultramafic rocks, in addition to the details of local geology (e.g. Buddington and Hess 1937). All early workers considered the rocks to be developed as part of

a large layered intrusion or intrusions. Smith (1958) completed the first regional mapping of the entire igneous complex and his synthesis followed the same premise.

Stevens (1965, 1970) provided the first recognition that the Bay of Islands Complex was a well-defined ophiolite suite, and was tectonically emplaced above the sedimentary rocks formerly considered to be older country rocks. There followed a long period of detailed investigation, involving workers from Memorial University and several other institutions, aimed both at understanding magmatic processes in the oceanic crust, and also the later structural and tectonic evolution of the complex. Calon et al. (1988) provide more complete details of all this research, including Ph.D. research projects from that period. Important contributions were also made by Dewey and Casey (2013) and Casey et al. (1985). The work of Suhr (1991) was particularly important in developing models for ophiolite emplacement, as discussed above (e.g. Cawood and Suhr 1992). Compared to the 1980s and 1990s, relatively little research has been completed in the area during the 21st century. Current research activity from Memorial University is centred on alkaline springs, where serpentinization reactions are producing methane (Szponar et al. 2013; Morrill et al. 2014); such environments are considered possible analogues for the surface on Mars.

Allochthonous Igneous Rocks of the Little Port Complex

Most of the localities discussed for the excursions are assigned to the Bay of Islands Igneous Complex (Figs. 5 and 6), but some stops in pillow lavas, sheeted dykes and trondhjemites are actually within the structurally underlying Little Port Complex. The following outline is drawn from Williams and Malpas (1972), Malpas (1979) and Jenner et al. (1991).

The Little Port Complex contains a bewildering assortment of plutonic and volcanic rocks, largely of mafic composition. Amongst the plutonic rocks, the oldest are foliated gabbros, peridotites and their altered equivalents. These were intruded by quartz diorites and trondhjemites. Plutonic rocks of all composition were cut by mafic dykes, of which there are probably several generations. The volcanic rocks of the Little Port Complex formed in submarine environments, and are dominated by pillow lavas and pillow breccias, associated with red cherts and black shales. Minor silicic volcanic rocks are reported in the area south of the park, and hydrothermal alteration of the type associated with sulphide mineralization occurs locally.

The confusing geology of the Little Port Complex is further complicated by alteration and low-grade metamorphism of many of the rocks, such that their original character is hard to establish. Nevertheless, many of its components likely represent the upper layers of the oceanic crust, and they are similar to rocks found locally within the Bay of Islands Complex. U–Pb dating of the trondhjemites near Trout River gave a Cambrian age of ca. 505 Ma, which contrasts with the younger Ordovician age (ca. 484 Ma) of the adjacent Bay of Islands Igneous Complex (Jenner et al. 1991). Based on this, and on aspects of its geochemistry, the Little Port Complex was rein-

terpreted as part of an older island arc assemblage (Jenner et al. 1991).

Allochthonous Igneous Rocks of the Bay of Islands Igneous Complex (Tablelands Ophiolite)

The Tablelands Ophiolite is the northernmost of four massifs of ultramafic and mafic rocks that form the Bay of Islands Igneous Complex (Fig. 5). A detailed review of all research is beyond the scope of this article, and the following general summary is derived largely from Calon et al. (1988), Cawood and Suhr (1992) and Williams (1995). Information specific to the Tablelands area is contained in Calon et al. (1988) and summarized in Stevens et al. (2003). The general geology of the Tablelands is indicated in Figure 8, redrawn and simplified from Suhr (1991), and also including work by T.J. Calon.

The Bay of Islands Igneous Complex contains all the components of the ophiolite suite. Ultramafic and gabbroic rocks are most extensive, and these are the only areas that are easily accessible. The lower structural contact of the ophiolites, where unmodified by later faulting, is marked by thin discontinuous zones of polydeformed amphibolites and schists. Individual ophiolite massifs within the Bay of Islands Complex, including the Tablelands Ophiolite, are disposed in synclinal structures that expose mantle rocks around their edges and preserve the higher levels of the oceanic crust in their centres. In addition to mafic and ultramafic rocks, there are minor rocks of trondhjemitic composition, which are generally similar to those of the Little Port Complex. Dunning and Krogh (1986) and Jenner et al. (1991) found that the Bay of Islands Complex is significantly younger than the Little Port Complex, at ca. 490–485 Ma. Casey et al. (1985) had previously proposed that these rocks were synchronous, representing mid-ocean ridge and transform fault environments, respectively. Jenner et al. (1991) also presented geochemical and isotopic data suggesting that the ophiolites formed in a supra-subduction zone environment. As discussed above, Cawood and Suhr (1992) proposed a model in which the relatively young oceanic crust now represented by the Bay of Islands Complex was generated within embayments in the Laurentian margin following initial collision of adjacent promontories with an offshore island arc.

The geology of the Tablelands area is described in detail by Calon et al. (1988) and Suhr (1992); the following summary is derived from these sources, and also from reviews by Malpas (1987) and Stevens et al. (2003). The various rock types are best seen in the east of the area, where there are essentially six units. The basal unit is the metamorphic aureole, which is separated from the ultramafic rocks proper by a mylonitic zone developed in the ultramafic rocks (Suhr and Cawood 1993). This basal mylonitic zone is overlain by a thick section of ‘depleted mantle tectonites’, which are mainly harzburgite with lesser amounts of dunite, and dykes and veins of pyroxenite. The mantle tectonites are so named because they contain strong fabrics developed in a high-temperature environment. The adjective *depleted* refers to the fact that these rocks are believed to have lost partial melt during the formation of the overlying oceanic crust.






Figure 8. Generalized geology of the western part of the Tablelands Ophiolite in Gros Morne National Park, adapted from a larger map by Suhr (1991), also incorporating work by T.J. Calon included in Stevens et al. (2003). The approximate route of the hike to the Moho area (Excursion 4) is shown, but for navigation, the detailed maps of Figure 9 are required.

The mantle tectonites are overlain by a zone believed to represent originally similar peridotites that were extensively injected by these partial melts. These include harzburgites and abundant dunites, locally with podiform chromite concentrations. This complex unit is sometimes referred to as the ‘transition zone’.

The transition zone is overlain by the interesting unit generally termed the ‘critical zone’, which was so named to reflect comparisons with classic layered mafic intrusions such as the Bushveld Complex and Skaergaard Intrusion (e.g. Smith 1958). This zone consists of strongly deformed ultramafic to mafic cumulates, forming a diffuse boundary between variably melt-depleted mantle rocks and the oceanic crustal rocks that formed from those melts. It represents the Moho, or Mohorovičić Discontinuity. The uppermost unit in the central part of the Tablelands Ophiolite consists of variably deformed to massive or locally layered gabbro, which locally contains diabase dykes.

The Moho is a complex zone, within which the *geophysical Moho* (i.e. the transition from mafic to ultramafic rocks, yielding a seismic velocity contrast) sits a few hundred metres above the *petrological Moho*, which is a more subtle boundary between cumulate rocks and melt-depleted rocks of the mantle. These separate definitions of the Moho are confusing to some (including the author) because both are petrological transitions, but the more obvious lithological contrast is not labelled as such.

The uppermost dyke- and volcanic-dominated parts of the oceanic crust were eroded from the Tablelands, but such rocks do occur within the adjacent North Arm Mountain massif, south of Trout River Pond (Fig. 5). Access to this area without a boat is difficult, as it is rugged and there are no defined trails, so it is not included in suggested excursions. However, the sheeted dykes are clearly visible from viewpoints along the Trout River hiking trail. The Moho is also exposed on the southern shore of Trout River Pond, as is the metamorphic




Figure 9. Extracts from National Topographic System (NTS) maps showing the stop localities and hiking routes for Excursions 2 and 4, which in part include off-trail hiking. (A). Part of NTS map area 12H/05 (Lomond) showing Excursion 2. (B) Part of NTS map area 12G/08 (Trout River). The original scale for both maps is 1:50,000, but may differ in this reproduction; the pale blue grid squares represent 1 km on each side. Copyright for the base maps rests with the Government of Canada.

aureole, but these areas are not visible from any of the excursions.

SUMMARY OF FIELD EXCURSIONS

Overview

This article outlines four excursions within Gros Morne National Park, each of which offers a different view of the Tablelands Ophiolite and associated rocks. The most ambitious is Excursion 4, which visits the Moho and underlying upper mantle rocks on the high plateau at around 600 m elevation. This is a strenuous venture and requires special considerations and preparation. Two other hiking excursions explore the lower country around Winterhouse Brook (Excursion 2) and Trout River Pond (Excursion 3), mostly using marked National Park trails. These are part-day ventures that can be completed in less-than-ideal weather, although they are far more pleasant if the sun shines. Excursion 1 is a collection of roadside stops on Route 431 and some associated short optional hikes that allow visitors to see typical rock types, including some that are absent or inaccessible in the Tablelands. Excursion 1, or parts of it, could easily be combined with either Excursion 2 or Excursion 3 to make a full day. However, Excursion 4 requires a full day with a good weather forecast.

Descriptive information for excursions comes in part from previous publications, notably Malpas (1987), Calon et al. (1988) and Stevens et al. (2003). The map of Berger et al. (1992) is an excellent resource with annotated panoramas drawn by the late Ian Brookes, and is highly recommended. However, its large size makes it difficult to use outside on a windy Newfoundland day! Figure 8 provides a geological map for part of the Tablelands Ophiolite and Figure 9 provides topographic maps showing the routes for Excursions 2 and 4. The photographs included in Figures 10 to 32 illustrate most of the outcrops described below.

Safety Considerations and Park Protocols

Visitors to all the excursion areas should be attentive to safety. Roadside outcrops require careful attention to traffic, as Route 431 is narrow and lacks a hard shoulder in some areas. Wherever possible, use the parking areas designated by Parks Canada. Newfoundland’s unpredictable weather can be hazardous as well as inconvenient. In high-elevation areas such as the Tablelands plateau, weather can deteriorate rapidly, and fog can obscure routes, leading to disorientation and unexpected encounters with cliffs. Unprepared hikers risk hypothermia if

they are caught by changing weather. Even on the marked National Park trails it is wise to have sturdy footwear, with strong ankle support, and this is essential for any off-trail hiking, as the terrain is rugged. In warm sunny summer weather, biting insects will almost inevitably accompany hikers. This article provides descriptive information and maps for off-trail hiking (Fig. 9), but hikers should carry a compass and a general topographic map. The use of a global positioning system (GPS) receiver is strongly recommended, because coordinates are provided for all locations, to allow them to be found precisely. The area is located mostly within National Topographic System (NTS) 1: 50,000-scale map areas 12G/08 (Trout River) and 12H/05 (Lomond), aside from the Woody Point area, which is within NTS area 12H/12 (Gros Morne).

Gros Morne National Park is a protected area, and the collection of samples (including loose material) is prohibited; similar restrictions apply to the removal of plants and flowers, and there are strict regulations about interaction with any wildlife. Formal research activities in the park, even if they do not involve sampling, require approval through Parks Canada. There are facilities for disposal of refuse on most official park trails, but off-trail hikers are responsible for removing any waste to protect the natural environment. Further information on Park regulations and other aspects of visiting can be found on the Parks Canada website (www.pc.gc.ca). Other information connected to visiting the Park is available at www.newfoundlandlabrador.com and its numerous links. Readers are asked to carefully respect all restrictions and procedures that are imposed to protect this World Heritage Site.

Getting to Gros Morne National Park and the Tablelands

Gros Morne National Park is located on the west coast of Newfoundland, north of Corner Brook (Fig. 1). Access is via Route 430, also known as ‘the Viking Trail’, which departs from Route 1 (the Trans-Canada Highway) at Deer Lake, where the regional airport is located. There is no public transportation to the park so most visitors will need a vehicle; rentals are available at Deer Lake airport and in the City of Corner Brook. Most accommodation and services are on the north side of Bonne Bay, around Rocky Harbour and Norris Point. The drive from these areas to the Tablelands will take slightly more than one hour, depending on traffic. There is also some accommodation on the south side of Bonne Bay, mostly around Woody Point, where the Park interpretation centre is located. Unfortunately, there is no longer a car ferry linking Woody Point and Norris Point, but a summer foot passenger ferry links the two communities, and transportation is also available from Woody Point to the Tablelands trailhead, allowing access to Excursion 3 and possibly Excursion 4 without driving. However, there are only a few return ferry trips each day, and the last water taxi run from Woody Point in the summer of 2019 was at 5.30 pm. Details of the schedule are best obtained onsite in the communities as it may vary from day to day and with the season.

All excursions are on the south side of Bonne Bay, accessed by Route 431, which branches from Route 430 at Wiltondale, and terminates at Trout River. This scenic road ini-

tially passes through autochthonous Cambrian–Ordovician carbonate rocks, and then passes into allochthonous transported sedimentary rocks before entering the Bay of Islands Igneous Complex and Little Port Complex between Woody Point and Trout River (see Fig. 6).

Finding the Localities

Roadside outcrops are located using kilometerage from known locations such as junctions or bridges, or by reference to Park facilities, but note that odometer readings in different vehicles may vary by as much as 10%. Stop descriptions include Universal Transverse Mercator (UTM) coordinates derived from GPS readings, which are generally accurate to ± 15 m or so. All UTM coordinates are in Zone 21, and use the NAD 1983 datum. Coordinates can be converted to the earlier NAD 1927 datum by subtracting 63 m from the Easting coordinate and subtracting 219 m from the Northing coordinate. For greater certainty and convenient reference, complete coordinates for NAD 1983, NAD 1927 and latitude/longitude systems are provided in Table 1.

EXCURSION 1: ROADSIDE OUTCROPS AND SHORT HIKES

General Information

This excursion proceeds from east to west, mostly along Route 431, although it does not have to be completed in that direction. Short optional hikes start at the Discovery Centre, at the trailhead for Green Gardens, and near the end of Route 431 in Trout River. Some roadside stops are adjacent to parking areas provided by Parks Canada, but others require careful parking on the gravel shoulder.

Stop 1.1: Panoramic Views of Bonne Bay and the Tablelands Ophiolite (43230E, 5482896N)

From Wiltondale, follow Route 431 to the South Arm of Bonne Bay, passing through the communities of Glenburnie, Birchy Head and Shoal Brook. Stop 1.1 is located just before the junction where Route 431 branches left up a steep hill, signposted for Trout River and the Discovery Centre. It is included because of its spectacular views (Fig. 10). Most geological elements of Gros Morne National Park can be seen from Stop 1.1.

The view of the Tablelands looming over Bonne Bay is justly famous, as is the view of the pyramid-like ‘Peak of Tenerife’, which represents the metamorphic aureole of the ophiolite, sitting beneath and adjacent to the igneous rocks. The mountain was named for its similarity in shape to the high volcano (Mount Teide) that dominates the island of Tenerife in the Canary Islands, off the northwest coast of Africa. ‘Gibraltar Peak’ is the flat-topped mountain across Bonne Bay, and was so named because it resembles the famous Mediterranean fortress. It is made up, in part, of transported volcanic rocks of rift-related origin from the earliest (late Precambrian) structural slices of the Humber Arm Allochthon. These features, and ‘Table Mountain’ itself, were named by Captain James Cook, who was one of the first Europeans to explore the west coast of Newfoundland from 1765 to 1767. The South Arm of Bonne Bay is incised into a regional

Table 1. A compilation of geographic coordinates for all locations discussed in this article, with reference to Universal Transverse Mercator (UTM) systems (NAD1927 and NAD1983 Canada data) and Latitude/Longitude, expressed as decimal degrees and minutes. The coordinates quoted in the article text are all with respect to NAD 1983 datum, but can be converted to NAD 1927 by subtracting 63 m from Easting and 219 m from Northing values. Note that this approximate conversion is valid only for this restricted geographic area.

Stop Identification (see text of paper)	UTM Easting (NAD 1927)	UTM Northing (NAD 1927)	UTM Easting (NAD 1983)	UTM Northing (NAD 1983)	Longitude (degree; min; sec)	Latitude (degree; min; sec)
Excursion 1: Roadside stops and assorted optional hikes						
1.1	433167	5482677	433230	5482896	57; 55; 19.39663 West	49; 28; 13.20201 North
1.2	433837	5483818	433900	5484037	57; 54; 46.78002 West	49; 30; 19.27930 North
1.3	431560	5482815	431623	5483034	57; 56; 39.36309 West	49; 29; 45.90000 North
1.4	432605	5482118	432668	5482337	57; 55; 46.98959 West	49; 29; 23.75213 North
1.5	430810	5481257	430873	5481476	57; 57; 15.66331 West	49; 28; 55.15211 North
1.6	430356	5481157	430419	5481376	57; 57; 38.16110 West	49; 28; 51.72842 North
1.7	422090	5483295	422153	5483514	58; 04; 30.45272 West	49; 29; 57.37263 North
1.8	417798	5481967	417861	5482186	58; 08; 02.79644 West	49; 29; 12.33030 North
1.9	417872	5481578	417935	5481797	58; 07; 58.82839 West	49; 28; 59.77221 North
1.10	418311	5482356	418374	5482575	58; 07; 37.59336 West	49; 29; 25.17524 North
Excursion 2: The upper mantle and aureole around Winterhouse Brook						
2.1	430572	5479490	430635	5479709	57; 57; 26.37465 West	49; 27; 57.84461 North
2.2	430370	5479525	430433	5479744	57; 57; 36.43175 West	49; 27; 58.89493 North
2.3	430955	5479640	431018	5479859	57; 57; 07.44214 West	49; 28; 2.85763 North
2.4	431690	5480100	431753	5480319	57; 56; 31.21336 West	49; 28; 18.04892 North
2.5	432020	5480110	432083	5480329	57; 56; 14.82384 West	49; 28; 18.50538 North
2.6	431385	5480425	431448	5480644	57; 56; 46.56974 West	49; 28; 28.44825 North
2.7	430665	5480265	430728	5480484	57; 57; 22.24324 West	49; 28; 22.97488 North
Excursion 3: Gabbro and peridotite units around Trout River Pond						
3.1	418458	5478951	418521	5479170	58; 07; 27.76096 West	49; 27; 35.00419 North
3.2	420560	5477260	420623	5477479	58; 05; 42.13569 West	49; 26; 41.25596 North
3.3	420860	5477110	420923	5477329	58; 05; 27.13141 West	49; 26; 36.53990 North
3.4	422010	5476365	422073	5476584	58; 04; 29.50011 West	49; 26; 12.95051 North
3.5	422880	5475845	422943	5476064	58; 03; 45.94084 West	49; 25' 56.51053 North
Excursion 4: The lower oceanic crust, moho region, and upper mantle on the high plateau						
4.1	424385	5482570	424448	5482789	58; 02; 35.85827 West	49; 29; 34.92936 North
4.2	424070	5482165	424133	5482384	58; 02; 51.23560 West	49; 29; 21.67655 North
4.3	424255	5481670	424318	5481889	58; 02; 41.69836 West	49; 29; 05.73158 North
4.4	424400	5481595	424463	5481814	58; 02; 34.44025 West	49; 29; 03.36750 North
4.5	424895	5480285	424958	5480504	58; 02; 08.94262 West	49; 28; 21.17102 North
4.6	424830	5480110	424893	5480329	58; 02; 12.05238 West	49; 28; 15.47626 North
4.7	424795	5480160	424858	5480379	58; 02; 13.82561 West	49; 28; 17.07972 North
4.8	424485	5479460	424548	5479679	58; 02; 28.74639 West	49; 27; 54.27814 North
4.9	424425	5479360	424488	5479579	58; 02; 31.65825 West	49; 27; 51.01374 North
4.10	424554	5478860	424617	5479079	58; 02; 24.90628 West	49; 27; 34.88189 North
4.11	425080	5478925	425143	5479144	58; 01; 58.82263 West	49; 27; 37.21857 North
4.12	425900	5479735	425963	5479954	58; 01; 18.63647 West	49; 28; 03.80293 North
4.13	425395	5480925	425458	5481144	58; 01; 44.53440 West	49; 28; 42.11214 North

Note: All UTM coordinates are in Zone 21. Conversion of coordinates between NAD 1927 and NAD 1983 datums, and the calculation of latitude and longitude were completed using the NTV2 online utility program, provided by Natural Resources Canada, at (webapp.geod.nrcan.gc.ca/geod/tools-outils/ntv2.php). UTM coordinates from GPS measurements are considered to be accurate to +/- 15 m for most localities.




Figure 10. Composite image of the view across the south arm of Bonne Bay, from near Woody Point (Stop 1.1). The view includes part of the Tablelands Ophiolite, the metamorphic aureole beneath it (Peak of Tenerife area) and allochthonous sedimentary rocks of the Humber Arm Allochthon (Gibraltar Peak). The peaks and Table Mountain were named by Captain James Cook after locations he had visited on other voyages.

mélange unit, remnants of which can be seen along its west shore. Gros Morne mountain (elevation 807 m) is clearly visible from the stop, and is the round peak resembling a bald head. Gros Morne is often cited as the highest point on the island of Newfoundland, but is actually the runner-up for that honour; a peak in the Lewis Hills (also part of the Bay of Islands Igneous Complex) is slightly higher at 814 m.

Stop 1.2: Gabbro and Mélange at the Woody Point Lighthouse (433900E, 5484037N)

From Stop 1.1, continue on Route 431 into the village of Woody Point, and take the road nearest to the shore through the small business district. The lighthouse is at the north end of the village, and is easily accessed by a short path. This stop, and several that follow, are located within the Little Port Complex, which sits structurally beneath the ultramafic rocks of the Bay of Islands Igneous Complex.

The Woody Point lighthouse is a much-photographed locality that affords excellent panoramic views of Bonne Bay, including Gros Morne mountain (Fig. 11). It is built upon a relatively homogeneous, fine- to medium-grained gabbro, which is one of the simpler and more homogeneous outcrops in the Little Port Complex. However, the gabbro is actually a large block within the wide mélange zone that sits at the base of the Little Port Complex. The cliffs below the lighthouse expose part of this mélange, which consists of a sheared, broken shale matrix containing lozenges of sandstone. Cracks in some of the sandstone blocks have been injected by the shale matrix material. The cliff exposures are visible from close to the lighthouse, but there is no safe route to descend to the shoreline.

Stop 1.3: Lookout Hills Trail, Tablelands Ophiolite Views, and Pillow Lavas (431623E, 5483034N)

From Stop 1.2, return through Woody Point, and take Route 431 (now signposted for Trout River), which climbs up the hill behind the village, and leads to the Discovery Centre. The Discovery Centre is the main interpretation location in this region of the park, with much of interest to visitors and abundant parking. The Lookout Hills Trail is a short but steep route that leads from the Discovery Centre to a nearby hilltop. The trail provides magnificent views, and some outcrops of variably preserved pillow lavas and pillow breccias. The viewing platform is also an excellent place to eat lunch on a nice day. The 5 km roundtrip trail involves an elevation gain of about 300 m and should only be attempted by visitors in good physical condition; about two hours should be allowed for the roundtrip hike.

The seemingly endless ascent leads to the junction for the summit loop, where the right hand branch leads quickly to the summit viewing platform. The trail junction has exceptional views of the Tablelands, and is a good place to see and photograph the view in the evening light, when the ultramafic mountains may appear blood-red (Fig. 12). The view includes the canyon of Winterhouse Brook, which is part of the hiking route for Excursion 2. At the summit (corresponding to the listed coordinates) the outcrop behind the wooden viewing platform consists of deformed pillow lava, forming part of the Little Port Complex. A short distance further along the trail is a rocky area where several large blocks have been arranged in circles for making fires. Some of these blocks are actually entire pillows, with chilled exteriors and radial cooling cracks.




Figure 11. The lighthouse at Woody Point (Stop 1.2) built on a mélangé unit including large blocks of altered gabbro (Little Port Complex). Gros Morne mountain (807 m) is seen clearly in the background above the community of Norris Point, across Bonne Bay.



Figure 12. The Tablelands Ophiolite and the deep valley of Winterhouse Brook, as seen from the highest viewpoint on the Lookout Hills Trail (Stop 1.3), above the Gros Morne Park Discovery Centre, near Woody Point. The area around Winterhouse Brook is the location of the Excursion 2 hiking route.



Figure 13. Mafic and granitoid rocks of the Little Port Complex at Stop 1.4, near the Gros Morne Discovery Centre. (A) Vein of trondhjemite (plagioclase-rich granite, at left) cutting altered and weakly metamorphosed gabbro and diabase. (B) Thin trondhjemite veins typically found all through this outcrop.

They have presumably been derived from outcrops by frost action. From the summit area, the trail eventually returns to the trail junction, and then back to the Discovery Centre via a series of boardwalks.

Stop 1.4: Varitextured gabbro, diabase dykes and trondhjemite (432668E, 5482337N)

From the Discovery Centre, walk or drive on Route 431 towards Trout River. This large and very obvious quarry is a very short distance uphill from the Discovery Centre, and was the source of most of the large blocks used in landscaping around the site. A few blocks remain, but the area is now easily accessible. The quarry face should be approached with caution, and many features are better seen in the blocks closer to the road.

The outcrop consists mostly of mafic intrusive rocks of the Little Port Complex (Fig. 13), and these exhibit wide vari-

ations in grain size, composition, texture and degree of alteration. Parts of it appear amphibolitic and are weakly foliated. It also contains white-weathering trondhjemite veins, but most of the examples in the outcrop face are thin and exposed on fracture surfaces. In addition to coarse-grained plutonic rocks, the outcrops contain a few areas of mafic dykes, representing higher layers of the oceanic crust, but these are not easy to find. These dyke-like units are mostly fine-grained gabbroic rocks rather than true diabase. This complexity is typical of many outcrops within the Little Port Complex. Although the overall appearance is very messy, this innate heterogeneity adds to the visual interest of the material when it is used for landscaping. Wider grey trondhjemite veins and dykes are most easily observed in large blocks at the uphill end of the quarry (Fig. 13), where there are also blocks of possible volcanic origin. The large blocks are the best places to see intrusive contacts between trondhjemite and the older gabbro, as the out-




Figure 14. Sheeted diabase dykes found at Stop 1.5, along with possible metavolcanic rocks, all part of the Little Port Complex. (A) General view of outcrops showing colour contrasts. (B) Detail of internal contact zone between two dykes.

crop is very fractured and loose material from the quarry face could be hazardous.

Stop 1.5: Sheeted(?) Diabase Dykes (430873E, 5481476N)

From Stop 1.4, continue on Route 431 for about 1.9 km, to a small outcrop located on the north side of route 431. Park as far off the road as possible, and watch carefully for traffic.

This outcrop is dominated by diabase and provides an interesting contrast with the next stop (1.6), which includes pillow lavas and deep-water sedimentary rocks. The outcrop is massive and well-jointed (blocky) which is more typical of diabase, and it contains some internal contacts, revealed by slight colour and grain size differences (Fig. 14). This suggests that it may consist of sheeted diabase dykes, typical of the upper part of the oceanic crust, located beneath pillow lavas formed on the seafloor. The outcrop is far from spectacular, but as far as I am aware it is the only easily accessible location that represents this important unit.

Stop 1.6: Pillow Lava and Seafloor Sedimentary Rocks, Hydrothermal Alteration and Sulphide Mineralization (430419E, 5481376N)

This outcrop is located opposite the Tablelands viewpoint on Route 431, the road towards Trout River, about 0.8 km west of Stop 1.5. The viewpoint provides safe parking for vehicles, but be attentive to traffic when crossing the road. The reference coordinate is the east end of the outcrop, but the entire outcrop is more than 300 m in length.

This is an extensive outcrop dominated by altered pillow lava, pillow breccia, red chert and red to black shale (Fig. 15). The outcrop is interpreted to represent the uppermost part of the ophiolitic basement to the island arc represented by the Little Port Complex, and part of its overlying sedimentary cover. The prominent rusty zone in the centre of the outcrop provides evidence of hydrothermal alteration and sulphide deposition, and probably represents part of a small submarine hydrothermal system, which vented mineral-rich hot brines

onto the ancient seafloor. The red and black sedimentary rocks immediately east of the sulphide-rich zone are cherts interpreted as seafloor sediment, perhaps formed from these exhaled brines.

The eastern part of this outcrop (corresponding to the coordinates listed above) contains well-preserved pillow lavas, which are west-younging based on their geometry and draping relationships. Compared to the previous stop, there are very few dykes, which is consistent with a higher level in the oceanic crust. There are also sections of pillow breccia in this outcrop, with sporadic entire pillows scattered through them (Fig. 15). These are located in the section between the pillow lavas and the sulphide zone, and also dominate the western end of the outcrop. The pillow breccias formed through explosive quenching of pillows, and through physical disaggregation when partly-solidified pillows rolled down submarine slopes.

Stop 1.7: Trondhjemite of the Little Port Complex (422153E, 5483514N)

From Stop 1.6, continue westward on Route 431, passing the trailhead for the Tablelands Hiking Trail (Excursion 2), and a second parking area which is the starting point for hiking up to the summit plateau of the Tablelands (Excursion 4). This parking area also connects to the Green Gardens Trail, but access to the trail is presently closed due to flooding damage. Continue on Route 431 towards its highest point, and park at the main trailhead for the Green Gardens Trail. The Green Gardens Trail is one of the most spectacular routes in the Park, and allows access to well-preserved volcanic rocks of the Skinner Cove Formation, which sit structurally below the Little Port Complex (see earlier description). A short hike of around 1 km along this trail to the first ridge leads to a trondhjemite outcrop, and also provides excellent views across Trout River Gulch to the Tablelands plateau. The vistas will probably make more of a lasting impression than the outcrop.

After leaving the trailhead, there is very little outcrop. The sides of the trail are lined with many orange and brown peri-




Figure 15. Features of Stop 1.6, an extensive outcrop of pillow lavas, pillow breccias, diabase and sedimentary rocks, all part of the Little Port Complex. (A) General view of pillow lava section. (B) brown-weathering pillow breccias, western part of outcrop. (C) Contact between reddish chert and black chert, both of which are interpreted as seafloor sedimentary facies. (D) A lonely basaltic lava pillow within a section of the outcrop dominated by pillow breccias.

dotite boulders and lesser amounts of grey gabbro and white trondhjemite. The landscape is forbidding and barren as the trail climbs gently, and is very typical of the entire Trout River Gulch area (Fig. 16). A few hundred metres from the trailhead, there is a gradual change in vegetation as low bushes and trees appear and cover the hillside. The appearance of this (relatively) lush vegetation probably marks the transition into the largely mafic rocks of the Little Port Complex.

Stop 1.7 consists of obvious white-weathering outcrops on the left hand side of trail, at the first ridge crest. These are coarse-grained, quartz-rich, leucocratic trondhjemite (Fig. 16). The outcrop has the slightly brecciated and fractured appearance typical of much of the Little Port Complex. The mafic minerals have been chloritized, so their original nature is unknown. If you are not continuing on the Green Gardens Trail towards the coast, return to the trailhead by the same route.

Stop 1.8: The Old Man of Trout River (417861E, 5482186N)

From Stop 1.7, continue westward on Route 431 to Trout River, and turn right at the junction on the edge of the village. Continue along the road through Trout River, and turn left on Riverside Drive, across a wooden bridge, and park by the signboards and wooden stairs. A short hiking trail leads from here to a raised seastack known as the "Old Man" (Fig. 17), and also provides access to the Lighthouse Trail (Stop 1.9). The seastack itself is a lineated to foliated amphibolitic gabbro, which includes rusty bands suggesting the presence of sulphides. A smaller collapsed sea stack next to the Old Man is (predictably) known as the Old Woman. The location provides excellent views of glacial features related to sea-level changes (Berger et al. 1992). The following summary is adapted from this source, and also from Brookes and Deardon (1981) and Stevens et al. (2003).





Figure 16. The route through Trout River Gulch. (A) The striking contrast between the barren rocks of the Tablelands Ophiolite (left) and the more ‘fertile’ wooded hills of the adjacent and structurally lower Little Port Complex (at extreme right). (B) Trondhjemite from the outcrops at Stop 1.7 (an optional short hike) from where there are good views of this area.



Figure 17. The Old Man of Trout River (Stop 1.8), a raised sea stack indicative of a prominent ancient shoreline, seen also is a terrace in the background, upon which most of the village is built.

The village of Trout River is underlain by a large terrace of gravel and sand that now separates the glacially-carved valley of Trout River Pond from the sea. This terrace is a classic ice-marginal marine delta, formed by meltwaters derived from a glacier that existed in Trout River Gulch. The top of this terrace is about 35 m above present day sea level, at the same level as the base of the raised seastack. This former sea level stand at 35 m is also represented by an obvious rock platform that extends on either side of the delta. These prominent features suggest that this was an important and long-lived sea-level stand. Radiocarbon dates obtained from shells indicate that the delta was deposited 12,700 years ago. Some lower terraces in the village mark later stages in the decline of sea level to its present position. Traces of a higher terrace and raised beaches around 70 m above present day sea level are dated at 13,400 years ago.

Stop 1.9: Trail to Trout River Lighthouse (417935E, 5481797N)

Just below the Old Man of Trout River (Stop 1.8) there is a junction on the access trail, signposted for the Lighthouse Trail. This is an easy walk of just over 1 km through some coastal forest and meadows that are full of wildflowers in summer, and leads to an automated lighthouse. There is no outcrop along the trail, but some low cliff-like outcrops visible to the south are part of same raised shoreline at 35 m that is defined by gravel terraces and the raised seastacks. At the lighthouse, the coastal cliffs are visible. These are extensively fractured rocks that are probably of mafic volcanic origin, and are assigned to the Little Port Complex. There are also good views across to the village of Trout River and outcrops on the north side of the harbour, which include some prominent pale grey areas formed from trondhjemites (Fig. 18). These can be visited by hiking along the Eastern Point Trail (Stop 1.10).

Stop 1.10: Trail to Eastern Point (418374E, 5482575N)

There is another short trail from the end of the road at the north end of the beach in Trout River to Eastern Point, which provides access to the trondhjemite outcrops visible from the Trout River lighthouse. To get to the trailhead, return to the main road through the village and turn towards the beach, and then drive or walk to the end of the road. The long wooden stairway ascending the gravel terrace is very obvious at the trailhead, but there is only limited parking. The hike is mostly of interest for its scenery and views, but there are outcrops of lineated coarse-grained trondhjemite at its furthest point (Fig. 18). Other outcrops in the area are black chloritic fractured rocks of uncertain origin, which are superficially similar to those seen in the cliffs at the lighthouse (Stop 1.9).




Figure 18. The view from the hiking trail that leads to Stop 1.8 and the Trout River lighthouse (Stop 1.9). (A) The light-coloured bands and areas on the opposite side of the Bay at Eastern Point are trondhjemite outcrops within the Little Port Complex. Note the prominent raised terrace and shoreline. (B) Strongly lineated trondhjemite from the location on Eastern Point (Stop 1.10) indicated by the white arrow in A.

EXCURSION 2: THE UPPER MANTLE AND METAMORPHIC AUREOLE AROUND WINTERHOUSE BROOK

General Information

A very popular Parks Canada trail commences near Route 431 and leads into the valley of Winterhouse Brook. This provides access to ultramafic rocks that represent the upper mantle section of the ophiolite, and also to much spectacular scenery, locally reminiscent of photos from the Mars Exploration Rovers. Excursion 2 includes this trail and then leaves the trail for some relatively easy off-trail hiking in the area east of Winterhouse Brook, ending at a beautiful river pool. The excursion reveals upper mantle peridotites containing strong tectonic fabrics, variably altered and serpentinized ultramafic rocks, and a small part of the metamorphic aureole that underlies the ophiolite. The Park trail is an easy walk including long boardwalks, but the off-trail hike requires sturdy footwear and adequate clothing, plus a map and (if possible) a GPS receiver. Some of the stops near the Park trail were described in the earlier article by Malpas (1987), but additional information is provided here.

Tablelands Trailhead to Winterhouse Brook

The parking area for the Tablelands interpretation trail is reached via a short branch road from Route 431, about 3 km west of the Discovery Centre. The hiking trail follows an old road for about 1.6 km, and is easy to walk. The route lies very close to the fault that separates ultramafic rocks to the south from mélanges and sedimentary rocks to the north. There are no outcrops along the trail, but there are many large blocks of brown-weathering peridotite derived from the huge cliffs that loom above the valley. Some blocks show spectacular fracture coatings of serpentine minerals. There are excellent views to the west, where the barren landscape of the Tablelands contrasts with the wooded hills of the Little Port Complex on the other side of Trout River Gulch. The flat landscape also shows

frost polygons (in which a central region of fine-grained gravelly material is surrounded by a ring of larger angular blocks on a scale of tens of metres). These are not always easy to see from ground level, but are widely developed. They result from frost-heaving and suggest the former presence of permafrost in this high-elevation area; similar frost polygons are widely present in the Canadian Arctic. At Winterhouse Brook, the Park trail diverges south from the old road, and after a few hundred metres it becomes a long boardwalk that leads to a small viewing platform. This is often a busy location in the summer season.

Stop 2.1: Winterhouse Brook Canyon and Alkaline Springs (430635E, 5479709N)

Stop 2.1 is located at the viewing platform at the end of the boardwalk, where there is abundant outcrop of massive brown-weathering peridotite in the stream and around the platform (Fig. 19). These are the most easily accessible peridotite outcrops in Gros Morne National Park, but they are generally homogeneous and featureless compared to the outcrops at Stop 2.2. Thin cross-cutting veins of serpentine minerals are prominent throughout the outcrops and locally yield photogenic fracture coatings (Fig. 19). The site has excellent views up the canyon of Winterhouse Brook, which is the largest of the river valleys that drain the Tablelands Ophiolite. The river course consists of numerous blocks, largely of ultramafic composition, and the bottom and sides of the valley are almost devoid of vegetation. The valley sides consist of unstable scree slopes that can be dangerous and difficult to negotiate. They are best avoided. The brook can be followed up into the canyon for around 1 km, although there is no defined trail, and the numerous peridotite blocks do not make for easy walking. Walking into the canyon is for the most part a scenic excursion, as the nearby outcrops at Stop 2.2 are probably more interesting for geologists.




Figure 19. Interesting features of the area around Winterhouse Brook Canyon (Stop 2.1 and vicinity). (A) A view into the canyon from the access trail. (B) View of the desolate landscape around the watercourse, underlain by ultramafic rocks of the mantle, viewing platform at the right hand side, as seen from Stop 2.2 (see Fig. 20). (C) Serpentine minerals on a fracture surface of a peridotite block. (D) Alkaline spring adjacent to Winterhouse Brook, showing white and cream travertine deposits from high-pH waters formed through subsurface serpentinization reactions.

About 75 m north from the viewing platform, the most accessible of several alkaline springs in the Tablelands Ophiolite is located adjacent to the boardwalk on the west bank of Winterhouse Brook. The presence of such alkaline springs was first noted by Stevens (1988). The alkaline groundwaters have deposited white and cream travertine-like layers over peridotite boulders and bedrock at this locality (Fig. 19). These unusual high-pH ground waters are produced by several chemical reactions involved in the formation of serpentine minerals by waters percolating through the ultramafic rocks. Sites such as this are suggested to be possible analogue sites for the surface of Mars, and a possible abiogenic source of methane (e.g. Szponar et al. 2013; Morrill et al. 2014). The springs at this locality are an active research project, and visitors should avoid walking on the travertine surfaces, which are fragile. Note that, although this is a spring, it is NOT suitable as drinking water, as its pH is in the same range as many domestic cleaning agents.

Stop 2.2: Depleted, Strongly Deformed Harzburgites and Pyroxenites, or “Mantle Tectonites” (430433E, 5479744N)

From the viewing platform at Stop 2.1, a faint trail leads west for about 200 m towards the base of the cliffs, providing abundant outcrops, which correspond to Stop 1b in the earlier article by Malpas (1987). The northern edge of these outcrops at the base of the cliff (right hand side as you approach them) is the primary area of interest. The layering in the outcrops becomes increasingly obvious as you hike up from the viewing platform. Layered rocks of similar type continue up the steep hill above the site, but this area can be hazardous and contains much loose rock.

The outcrops (Fig. 20) are the most accessible examples of harzburgites from the mantle tectonite sequence, and they have been strongly deformed under the extreme pressure and temperature conditions of the mantle. The layering is not of magmatic origin, but instead results largely from deformation




Figure 20. The ‘mantle tectonite’ outcrops above Winterhouse Brook (Stop 2.2). (A) General view towards Bonne Bay from above the site showing the landscape and the banding in the rocks. (B). Strongly banded harzburgite and dunite with concordant pyroxenite bands (best seen at upper left); the vertical vein consists of serpentine minerals and cuts the older banding. (C) One of several tight to isoclinal folds defined by pyroxenitic layers in these outcrops. (D) Thicker pyroxenite veins showing evidence of some later gentle folding.

processes. The harzburgites contain about 60% olivine and 30% Mg-rich orthopyroxene, with smaller amounts of clinopyroxene and oxide minerals. These are believed to be residual ultramafic rocks that have been depleted in their low-melting point constituents by partial melting related to the formation of the rocks of the overlying oceanic crust. The banded appearance of the outcrops largely reflects variations in the olivine to orthopyroxene ratio, which leads to differences in colour and weathering response. The contacts of individual bands appear abrupt, but they are generally gradational in detail; individual bands can be traced for up to 500 m from this stop. Some of the layers are up to 100 m thick, but most are just a few metres thick. The layering results from intense ductile deformation of original compositional heterogeneities in the rock, and the rotation of veins and dykes of different composition to their hosts. The process that produced the intense layering is called transposition, and the formation of these

rocks is closely analogous to the development of banded gneisses in high-grade metamorphic environments. The features evident in these spectacular outcrops are similar to those described from other ophiolite complexes around the world.

The banding in the ultramafic rocks is cut locally by less-deformed veins and dykes of variable composition; some are essentially pyroxenitic, whereas others are extremely olivine-rich and are almost dunitic. The pyroxenite veins and pods are best observed in the higher parts of the outcrop. Pyroxenites are readily identified because they lack the yellow-brown weathering crust developed on the olivine-rich rocks and they exhibit many crystal faces from the pyroxenes within them, which weather to a lustrous pale grey-green. On a sunny day, these pyroxenite veins are easy to see because they actually sparkle. The pyroxenite zones locally retain discordant contacts, but elsewhere are parallel with the regional layering because they have been reoriented by deformation. Some of






Figure 21. Typical appearance of ultramafic rocks representing the mantle around Winterhouse Brook. (A) Networks of serpentine veinlets dissecting peridotite. (B) More intense development of fractures and serpentine veins. Photos are from Stops 2.3 and 2.4, respectively.

these pyroxenitic bands are folded to form tight to isoclinal folds (Fig. 20), some of which are entirely contained within the regional layering, i.e. they are rootless intrafolial folds.

Both types of veins likely formed during partial melting of the mantle to produce mafic liquids. The dunitic veins have retained only olivine, which was the first mineral to crystallize from these liquids; pyroxenites result from coprecipitation of olivine, orthopyroxene and clinopyroxene. The gabbros in the western section of the Tablelands (see Excursions 3 and 4) represent the final crystallization products of the mafic liquids generated at this deeper level within the mantle.

Stop 2.3: Harzburgites With Serpentine Veining (431018E, 5479860N)

From Stop 2.2, walk downhill again towards the boardwalk, and continue across Winterhouse Brook. The brook is easily crossed at most times of the year. From the brook, walk eastward towards a partially tree-covered hill on the horizon that has a grey colour unlike the yellow-brown that dominates most of the landscape immediately ahead. The terrain is relatively flat and varies between gravelly material that is relatively pleasant to walk upon and areas of angular peridotite blocks that are very hard on the feet and ankles.

The outcrops of interest are located on a low broad ridge, about 0.5 km from Winterhouse Brook. They are harzburgites akin to those of Stop 2.2, but they lack the well-developed compositional layering. These outcrops instead show visually striking net-like patterns of serpentinized veinlets that are very typical of many ultramafic rocks in the Tablelands, and elsewhere (Fig. 21). The pattern is in many respects reminiscent of the veined and dissected appearance of olivine in thin sections of ultramafic rocks, and is an interesting example of how outcrop-scale and microscopic textures can mirror one another in these olivine-dominated rocks.

Stop 2.4: Strongly Serpentinized Ultramafic Rocks (431753E, 5480319N)

From Stop 2.3, continue walking for about 700 m to the prominent low grey-coloured hill noted above. The colour contrast might suggest that it consists of a different rock type to Stop 2.3, but this is not the case. The outcrop (Fig. 21) is more intensely veined and pervasively serpentinized compared to Stop 2.3, which is in turn more serpentinized than the well-preserved rocks at Stop 2.2. This is probably because the excursion route is getting progressively closer to the faulted contact between the Tablelands Ophiolite and adjacent the mélange units.

Stop 2.5: Amphibolites of the Metamorphic Aureole (432083E, 5480329N)

From Stop 2.4, walk southeast, crossing some more outcrops of strongly serpentinized ultramafic rocks, and then turn northwards to ascend a partially barren ridge with some small scrubby trees on its upper part. Head towards two distinctive pyramid-shaped peridotite boulders, and then climb as far as you can without actually getting into the trees, which is not advised, as they consist mostly of impenetrable wind-stunted spruce. The upper part of this small hill exposes amphibolites of the metamorphic aureole beneath the ophiolite. Stop 2.5 also provides some wonderful views of the east flank of Table Mountain, and reveals the steep forested mountains above Birchy Head, including the distinctive Peak of Tenerife (Fig. 22). The peak is also part of the metamorphic aureole, although it is not physically connected to the rocks exposed at Stop 2.5. The prominent mountain above Birchy Head consists of altered basalts and dykes representing a large tectonic inclusion of Little Port Complex rocks within the mélange zone. There is also a very good view northward to the South Arm of Bonne Bay, Woody Point and Norris Point.




Figure 22. Stop 2.5 located on the metamorphic aureole of the Tablelands Ophiolite. (A) The contrast between the eastern edge of the ophiolite (at right) and the forested hills of the metamorphic aureole, including the Peak of Tenerife. (B) Foliated and lineated fine-grained amphibolitic rock of the aureole, likely derived from mélangé including mafic volcanic material.

The rocks are not easily examined, but this is the only portion of the metamorphic aureole that is easily accessible on foot. The rock type is a fine-grained, strongly foliated to lineated amphibolite of mafic composition that locally shows some poorly developed compositional layering (Fig. 22). The original nature of the aureole rocks is not always clear, but they are generally of basaltic composition and are likely derived from rocks of the upper oceanic crust (i.e. sheeted diabase dykes and basalts) in mélanges that were "welded" to the ultramafic rocks during the earliest stages of obduction.

Stop 2.6: Severely Altered and Sheared Ultramafic Rocks (431448E, 5480643N)

From Stop 2.5, retrace the route across Stop 2.4, and then head northwest towards Stop 2.6, which is marked by a low rubbly hill with a prominent green meadow on its west side in the summer months. This spot is mostly included for its superb views of Bonne Bay, including Woody Point, Norris Point and Nedly Harbour, with Gros Morne mountain in the background. The outcrop, which is less spectacular, consists of severely altered and sheared ultramafic rocks, which superficially resemble shale. The outcrop is located very close to the faulted northern contact of the ophiolite, which probably explains why it is strongly sheared and retrogressed.

Stop 2.7: 'Xonotlite Layer' at the Northern Contact of the Ophiolite (430728E, 5480484N)

From Stop 2.6, walk a short distance northwest to the old road, which is marked by an abandoned telephone line. Continue northwestward across the rubbly peridotite and gravel. Just before a band of scrubby vegetation, there is a small stream, flowing westward. Follow this stream to the west. The valley gradually becomes more deeply incised, and eventually leads into the deeper lower valley of Winterhouse Brook, which is canyon-like. The descent along the brook is not difficult, but be wary of large blocks of ultramafic rock that might

be unstable. The valley runs just south of the fault that marks the northern edge of the Tablelands Ophiolite, and the outcrops are all strongly serpentinized ultramafic rocks. Some fracture surfaces exhibit spectacular fibrous green and blue-green serpentine minerals (Fig. 23). The stream eventually joins Winterhouse Brook. Stop 2.7 is visible downstream from the junction of the tributary brook with Winterhouse Brook, and it is marked by a prominent wall-like feature extending across the brook (Fig. 23). It corresponds to Stop 1a of Malpas (1987) and marks the contact of the ophiolite with the sedimentary rocks to the north.

The contact of igneous and sedimentary rocks at this location is a fault, and there is no metamorphic aureole preserved. However, boulders of aureole rocks that resemble those of Stop 2.5 are located here and there in the stream bed. The ultramafic rocks in this area are pervasively altered, and few original textures or minerals are preserved. The contact itself is marked by a resistant, pale grey unit that resembles a bed (Fig. 23). This is commonly referred to as the "xonotlite layer", after a rare mineral that it contains. It is actually a metasomatized ultramafic rock that also contains Ca-rich prehnite, calcite and wollastonite. Because the ultramafic rocks contain very little calcium, it is generally assumed that the source of the extra calcium in this layer was *outside* the ophiolite, i.e. within the adjoining sedimentary rocks. The metasomatism of the ultramafic rocks at the contact may have occurred during emplacement of the ophiolite, but could equally well reflect fluid migrations associated with later faulting.

Xonotlite is an unusual mineral with the formula $\text{Ca}_6\text{Si}_6\text{O}_{17}(\text{OH})_2$. It is relatively hard (6.5 in the Moh's hardness scale) which accounts for the resistant nature of this layer. It is named for a place called Tetela de Xonotla, in the state of Puebla, Mexico. The Winterhouse Brook locality was the first xonotlite to be described in Canada. The xonotlite was initially reported here by Smith (1958), and he provides the most detailed description that I have found. The mineral occurs at




Figure 23. The edge of the Tablelands Ophiolite in Winterhouse Brook, at the 'xonotlite layer' (Stop 2.7). (A) Contact between reddish shale and mélangé (left) and serpentinized ultramafic rocks (right), including the resistant xonotlite layer. (B) A view of the xonotlite layer crossing the brook, where it creates a waterfall and a tempting pool; note that access to the pool for swimming is now difficult and potentially hazardous. (C) Contact between the xonotlite layer (white, metasomatized ultramafic rock) and soft green serpentinite at right. (D) One of many spectacular fracture surfaces displaying serpentine minerals seen in this area.

another locality in the Bay of Islands Igneous Complex (Cox's Cove; Smith 1958) and also in calc-silicate metamorphic rocks near Rose Blanche, in southwestern Newfoundland (Brown 1978). Visitors should not expect to collect any samples, because Park regulations prohibit any removal of material. Also, the mineral is essentially invisible to the naked eye, as it is intergrown with the other minerals noted above.

The xonotlite layer forms a prominent waterfall in Winterhouse Brook, below which is a beautiful deep circular pool. Unfortunately, access to this pool for swimming is increasingly hazardous because the banks have become deeply eroded and very steep. Although it may be tempting, it is not recommended.

From Stop 2.7, cross Winterhouse Brook below or above the swimming hole, and carefully climb the western slope of the valley, sticking to the more stable vegetated slopes. At the top of the bank, turn upstream and walk south or southwest to rejoin the old road between Woody Point and Trout River.

This section of the hike crosses some very well-developed frost polygons before joining the old road. The walk back to the parking area should take 15 to 20 minutes.

EXCURSION 3: PERIDOTITE AND GABBRO IN THE TROUT RIVER POND AREA

General Information

The Trout River area has several features of geological interest, including the raised sea-stack of 'The "Old Man"' and other indications of changing sea level (see Excursion 1). The long lake of Trout River Pond occupies the valley between the Tablelands and North Arm Mountain, both of which are part of the Bay of Islands Igneous Complex (Fig. 5). Trout River Pond has excellent shoreline exposures that represent the upper crustal section of the latter, and also the Moho region, but these are only accessible by boat. A tourist excursion on the lake during the summer provides a chance to see these




Figure 24. The views from the location near Trout River Campground (Stop 3.1). (A) Trout River Pond separating the ultramafic rocks of the Tablelands (left) from the higher levels of the ophiolite sequence exposed in the North Arm Mountain massif (right). (B) A closer view of the mountain known as the “Elephant’s Head” (officially termed Narrows Head) showing its resemblance to the folded ears and long trunk of an elephant. Most of this prominent feature consists of sheeted diabase dykes.

from a distance, but the boat tour availability has varied in recent years. Excursion 3 is a hike of about 11 km roundtrip, which starts at the boat launching area, and follows an official Park trail; it is a relatively easy walk, without any significant elevation gain, and is well marked. The hiking trail encounters little outcrop except at its easternmost end, but it provides access to peridotites and gabbros of the Tablelands Ophiolite without actually climbing to the high plateau. The trail also provides many scenic views of the Tablelands and North Arm Mountain, including distant views of the sheeted dykes in the latter. At least 4–5 hours should be allowed for the return hike. A visit to the scenic viewpoint by the campground (Stop 3.1; signposted) is also highly recommended.

Stop 3.1: Viewpoint Near the Trout River Campground (418521E, 5479170N)

If entering Trout River from Route 431 westbound, turn left at the junction, and continue west beside the river, following the signs for the campground. The road towards the campground is the extension of Main Street in the village. Turn right at the junction for the day-use area, cross the river bridge, and ascend to the viewpoint, signposted on the right.

This viewpoint gives one of the more spectacular views in the Park (Fig. 24), and it contains a wealth of geological information. The south side of the Tablelands Ophiolite is clearly visible, and the contrast between red-weathering ultramafic rocks and grey-weathering gabbro is obvious. Miner’s Point, at the head of the lake, also consists of gabbro. A large detached piece of gabbro has moved downslope towards the lake in postglacial times; this is visited at a later stop on the hike. A vertical fault occupies the lake valley, and the hills to the south (right) are part of the North Arm Mountain massif. This area is dropped down relative to the Tablelands Ophiolite and also tilted towards the viewpoint, exposing the metamorphic aureole of the ophiolite at the far end of the lake. The hills on the south side of the lake thus expose higher sections of the

oceanic crust, such as sheeted dykes and gabbro, which are less abundant in the Tablelands. A prominent rounded peak has a long ridge that connects its summit to the shore of the lake, and rounded cliffs on either side (Fig. 24). This is known locally as the “Elephant’s Head”, because the long ridge resembles a trunk and the rounded cliffs resemble ears. It consists mostly of sheeted diabase dykes. The high peak in the south is Mount St. Gregory (674 m), which is one of the highest points in Newfoundland. The Mount St. Gregory area was well-known for numerous but unfortunately small copper-bearing veins, which is how Miner’s Point got its name.

Stop 3.2: Mélange and Shale Outcrops (420623E, 5477479N)

From the viewpoint (Stop 3.1) return towards Route 431, turn right after crossing the bridge, and then take the side road signposted for the day-use area. The trail starts just behind the toilets. There are no outcrops at all for the first 2.5 km, but the trail is very pleasant, running through thick woodland just above the shore of the lake. The lack of outcrop in the first part of the trail reflects the fact that it is largely within the mélange unit that sits beneath the Tablelands Ophiolite. Some reddish shales of this unit are exposed within a small stream at this location, but may not always be visible above the water level.

Stop 3.3: Stream Crossing and Views of Slumped Gabbro Mass (420923E, 5477329N)

A few hundred metres beyond Stop 3.2, the trail emerges into countryside that is more open and crosses a larger brook. The boulders in the brook include numerous peridotites and lineated to foliated gabbros. The latter resemble the gabbros that outcrop in the region of the Moho (see Excursion 4) and may be derived from similar sources.

This location has a view of huge grey outcrop that has a flat top, sloping to the south (Fig. 25). The grey outcrop con-



Figure 25. The large mass of gabbro (grey) sitting physically above peridotite (orange-brown) in the valley of Trout River Pond (Stop 3.3). This feature was interpreted by Smith (1958) to be a product of postglacial mass movements in the valley and its glacially oversteepened cliffs.

sists of gabbro, whereas the outcrops beneath it are clearly peridotite, and they have the distinctive red-brown colour of the Tablelands. However, the contact between these two rock types is not the Moho in this particular location. Smith (1958) proposed that this enormous crag of gabbro "slumped" into the valley of Trout River Pond in postglacial times, when the support of the ice sheets was lost following their melting. Large areas of slumping are identified by airphoto analysis of the areas, and smaller areas of gabbro scattered through the valley sides were interpreted in a similar way (Smith 1958). Further discussion of this idea is provided by Brookes and Deardon (1981) and Brookes (1993).

Stop 3.4: Gabbro Outcrops (422073E, 5476584N)

Stop 3.4 is located about 4.5 km from the trailhead. There are 'outcrops' immediately above the trail that belong to the slumped gabbro mass, and large blocks adjacent to the trail that represent the same material. The rock type is an altered and retrogressed foliated gabbro, which contains some discontinuous lenses of ultramafic composition, possibly feldspathic dunites. The gabbros are not as well-preserved as those found on the Tablelands high country excursion (see Excursion 4), but they are likely derived from the region of the Moho.

This location also affords excellent views of the narrows in Trout River Pond, and the imposing cliffs on the opposite side of the lake (Fig. 26). The highest cliff visible across the lake (part of the "Elephant's Head" noted at Stop 3.1) is mostly formed by sheeted diabase dykes, representing the feeder systems to seafloor mafic volcanic rocks. The lowermost part of the cliff consists of gabbro. The vertical dykes in the cliff face are just visible to the naked eye if you have keen eyes, but are more easily seen through binoculars.

From Stop 3.4 onwards, the trail passes through ultramafic rocks as evidenced by the barren country and numerous peridotite boulders. Note the view down to the beaches on either side of the narrows in Trout River Pond; these display the contrasts in geology very nicely. The beach on the north side is

orange, as it is largely of ultramafic composition, whereas the beach on the south side is grey, being dominantly mafic (Fig. 26).

Stop 3.5: Peridotites (422943E, 5476064N)

The outcrops at Stop 3.5 lie very close to the end of the trail, and are the most extensive areas of bedrock anywhere on the trail. They consist of brown-weathering peridotite and harzburgite, cut by both pyroxenite and dunite dykes (Fig. 27). The rocks contain a weakly developed fabric. Looking up towards the cliffs of the Tablelands Ophiolite, you may see several greyish patches indicating the locations of alkaline springs, similar to the one noted in Winterhouse Brook on Excursion 2. There are also very good views of the Trout River Pond valley and Gregory Plateau from this location.

EXCURSION 4: THE LOWER OCEANIC CRUST, THE MOHO AND THE MANTLE ON THE TABLELANDS SUMMIT PLATEAU

General Information

Excursion 4 is a lengthy off-trail hike, but well worth the effort. It visits the Moho, which marks the gradational transition between ultramafic rocks of the mantle and gabbroic rocks of the lower section of the oceanic crust. Early studies of the Bay of Islands Igneous Complex (BOIC) called this the "Critical Zone" (Smith 1958), and this term is retained in some accounts. This terminology is derived from large layered intrusions such as the Bushveld Complex of southern Africa, where it refers to the region where cumulus plagioclase first appears in the crystallization sequence. The hike initially crosses through gabbroic rocks of the lowermost oceanic crust and, if completed fully, eventually traverses some of the spectacular scenery formed by ultramafic rocks on the high plateau.

Specific safety concerns apply to this excursion. It should not be attempted unless you are in good physical condition and well prepared. It should never be attempted if the summit plateau is covered in cloud, and hikers should quickly descend if the weather deteriorates. The time required depends on how many of the stops below are included. A straight hike to the Moho and back on the same route can be done in about 5 hours, but if the return loop through the underlying ultramafic rocks is included, at least 7 or 8 hours should be allocated. Although it is quicker to retrace your steps from the Moho, the walk across the ultramafic plateau is a rather surreal experience. The area around the Moho is also described by Malpas (1987), and information is also provided by Stevens et al. (2003), but without locational and directional information.

Climbing from Route 431 to Stop 4.1

The starting point for the hike is the parking lot for the Green Gardens Trail on Route 431, between the Discovery Centre in Woody Point and Trout River. The UTM coordinate for this spot is approximately 424650E, 5283050N. Note that there are two separate trailheads for the Green Gardens Trail system, and that this is the first parking area on the right hand side when heading west on Route 431 towards Trout River. If trav-




Figure 26. Views from the Trout River Pond Hiking Trail, at Stop 3.5. (A) Another view of the “Elephant’s Head”, where sheeted diabase dykes can be seen by those with keen eyes or good binoculars. (B) The contrasting colours of cobble beaches on either side of Trout River Pond Narrows that reveal the contrast between peridotite (foreground) and gabbro or diabase (background).

elling east, it is the second parking lot on the left. In 2019, it was not visibly identified for the Green Gardens Trail, due to flood-related damage on parts of the trail. From the trailhead parking lot, walk a short distance west, and then leave the road to the left just past the large signboard. Ascend the steep slope on the west bank of a small stream.

Stop 4.1: Route Orientation (424448E, 5482789N)

There is no outcrop at this location, but it is a good place to take a rest. The site consists of a large pile of ultramafic boulders, which are probably frost-heaved blocks from underlying bedrock. From here, you can see your destination on the high ridge ahead. A prominent outcrop is visible just to the right of a notch where a small stream emerges. During the remainder of the steep ascent, keep this outcrop in sight and head directly towards it. You will probably have many chances to view the route ahead as you stop to regain your breath.

Stop 4.2: Coarse-Grained Gabbro of the Lower Oceanic Crust (424133E, 5482384N)

As you ascend the hill from stop 4.1 keep to the west (right hand side) of the stream valley. Note that the brown ultramafic talus becomes mixed with grey blocks of gabbroic composition as the hill is climbed. There are few definite outcrops enroute, because the surface is extensively frost-shattered, but the transition between the ultramafic rocks and the gabbro is marked, as elsewhere in the park, by increasing amounts of scrubby vegetation.

Stop 4.2 is the first large outcrop on the route, and it consists of coarse-grained gabbro. The contact between the ultramafic rocks of the lower slope and this gabbro is a later fault zone, and does not actually represent the Moho at this location. It is hard to see fresh surfaces in this outcrop. The gabbro is medium- to coarse-grained and consists largely of plagioclase and subophitic clinopyroxene, with lesser orthopyroxene. There is some local alteration of the mafic minerals to amphibole and chlorite. The absence of primary hydrous minerals



Figure 27. The visually striking peridotite-dominated landscape around the end of the Trout River Pond Hiking Trail (Stop 3.5). The upper, more remote, part of Trout River Pond is visible in the distance.

and the low modal abundance of orthopyroxene suggest that these rocks represent an intermediate level in layer 3 of the oceanic crust. There are also some local concentrations of diabase dykes in this general area (but not seen at this outcrop) that probably represent the roots of the overlying layer 2 (sheeted dykes) which has largely been eroded.

Stop 4.3: Coarse-Grained Gabbro (424318E, 5481889N)

From Stop 4.2, make a slight change of direction to southeast as you continue up the slope. This will eventually lead to the large outcrops of Stop 4.3, which is the prominent rocky hill first observed from Stop 4.1, and almost on the summit plateau. At this point, take a well-deserved rest and admire the wonderful views of Gros Morne mountain to the northeast, and the broad sweep of the coast directly to the north, beyond Rocky Harbour.





Figure 28. Examples of gabbroic rocks seen on the ascent to the Tablelands plateau (Stops 4.3 and 4.5). (A) Foliated gabbro, with prominent leucocratic bands. (B) Mafic cumulate rocks with brownish troctolitic intervals and grey gabbroic material; it is not clear if the foliation is magmatic or superimposed.

The outcrop at Stop 4.3 consists of coarse-grained gabbro that resembles the outcrops at Stop 4.2, but it is much better preserved. It contains a well-developed plagioclase alignment and foliation that at first sight appears to be of primary magmatic origin (Fig. 28). Some compositional layering is developed, but this is generally better seen in loose blocks than in the outcrops. Petrographic studies indicate that the layering and foliation are partly due to deformation and recrystallization under high-temperature ductile conditions rather than magmatic accumulation (Calon et al. 1988; Stevens et al. 2003). However, it is likely that this recrystallization and flow closely followed the crystallization of the host rocks, as these fabrics are locally cut by undeformed gabbro and anorthositic segregations.

Stop 4.4: Views of the Plateau and Moho Region (424463E, 5481814N)

From Stop 4.3, continue for about 150 metres to the east, to the true summit of the hill. The outcrops here are essentially the same as those described from Stop 4.3, but more extensive. From here, the next section of the hike is clearly visible. To the east, the Tablelands summit plateau is now visible, consisting largely of brown-weathering ultramafic rocks; these are incised by a shallow stream valley. To the right of the stream valley is a large barren hill with brown ultramafic rocks showing through a ragged cover of green grass, which is in turn flanked to the right by a more vegetated hill on which the rocks are grey in color. This hill is located on a bearing of approximately 120°, and consists of gabbro; the Moho separates these mafic rocks from the ultramafic rocks to the east. This grey hill represents Stop 4.5, and the route heads directly towards it.

Stop 4.5: Strongly Foliated and Lineated Gabbro (424958E, 5480504N)

From Stop 4.4, walk south-southeast for approximately 1.3 km to reach the north end of the long ridge that exposes the Moho. The terrain is generally flat, and the walking is for the

most part relatively easy, albeit swampy in places. Some sections consist of frost-heaved and broken rock that is hard on the ankles and requires some caution. However, this is nothing compared to the broken terrain that you will encounter later when you get into the ultramafic rocks in the final section of the hike.

Stop 4.5 is a prominent grey outcrop with many large frost-heaved boulders. The hike from Route 431 to this spot will generally take about 2 hours, and this is one of two places recommended for a possible lunch break. The second location is Stop 4.6, only 200 m south from which the views are superior, but likely to be windier. Stop 4.5 provides better shelter from the wind in the lee of some of the large blocks.

Outcrops at Stop 4.5 consist of coarse-grained, foliated to lineated gabbro, showing variably-developed compositional layering (Fig. 29). However, there are also many blocks of feldspathic dunite, which are typically chocolate-brown weathering, and contrast with the grey-weathering gabbro. South of Stop 4.5 towards the crest of the ridge, there are several outcrops of this material, many of which contain fine-grained plagioclase-rich (nearly anorthositic) layers. Minor chromite is visible on many weathered surfaces, and chromite-rich pods are locally visible, but there are no significant chromite accumulations. This is a gradational contact region, with feldspathic dunites to the east, gabbro to the west, and interlayered dunites and gabbros in the middle. It represents the geophysical Moho, despite the fact that there is an obvious contrast in petrology; the compositional shift from gabbro to peridotite would result in an increase in seismic velocity. In essence, the Moho at this site is a strongly deformed package of ultramafic and mafic rocks including dunite, feldspathic dunite (described as troctolite by Malpas 1987), pyroxene-rich gabbro and olivine gabbro, and a distinctive assemblage of anorthositic dykes and veins. The strong fabrics and numerous folds indicate intense plastic (ductile) deformation, and several features (notably rotated clinopyroxene megacrysts) indicate that the Moho was essentially a large-scale shear zone. The fabrics and the evidence of




Figure 29. Examples of rock types seen in the region of the Moho at Stop 4.6 and adjacent areas. (A) Typical outcrops showing the mixed character of the Moho region, with alternating zones of variably layered gabbro (grey) and olivine-rich ultramafic rocks (brown); in detail, such compositional domains are more like elongate lenses than layers. (B) Leucocratic layers, preserving contact relationships with associated mafic cumulate rocks. (C) Strongly foliated feldspathic dunite showing colour variations defined by the plagioclase to olivine ratio. (D) Sharply-defined plagioclase-rich bands (originally cumulates?) in a rock type similar to (C), but note also the cross-cutting plagioclase-rich vein in the centre of the photo. Photos (A) and (B) by John Waldron, University of Alberta.

deformation are in many respects similar to the features observed near Winterhouse Brook (Excursion 2) and in later parts of this excursion. Deformation of this intensity and type is almost never seen in the layered sequences of large mafic intrusions such as the Bushveld Complex, and its presence puzzled earlier geologists who interpreted the BOIC from this perspective.

Stop 4.6: Moho Region (424893E, 5480329N)

Stop 4.6 is located on the ridge crest about 200 m south of Stop 4.5, and is perhaps the best spot to view the Moho and to photograph the location with spectacular views in the background (Fig. 30). The contrast between grey gabbros to the west and brown dunites to the east is very clear here. There are numerous white anorthositic layers in the dunite; some of

these are discordant to the foliation (Fig. 29), implying that they represent originally cross-cutting veins that were later rotated into parallelism.

Stop 4.7: Deformed Augite Megacrysts (424858E, 5480379N)

Stop 4.7 is the grey layered outcrop that is visible from Stop 4.6, and located about 70 m to the northwest. It is a gabbroic outcrop that is notable for the presence of strong layering, foliation and many olivine-rich layers. It also contains some large clinopyroxene (augite) megacrysts, which have become augen due to pervasive shearing (Fig. 31). The outcrop also contains some intrafolial isoclinal folds, which are much harder to see than the megacrysts.



Figure 30. Panoramic view of the Moho location on the Tablelands (near Stop 4.6), with pale grey gabbro in the background and interlayered mafic and ultramafic rocks in the foreground. The view in the far distance extends to the Lookout Hills and the north side of Bonne Bay.

Stop 4.8: Deformed Gabbro and Views to the South (424548E, 5479679N)

From Stop 4.7, walk east to the ridge crest again, and then walk south along the ridge. There is not very much outcrop enroute to Stop 4.8, and most of the ridge consists of frost-heaved rubble. One section is bizarre in that there appears to be east-trending zones of gabbroic and ultramafic rubble, which are at right angles to the regional trend of the contact between these two rock types. It seems unlikely that this actually reflects the distribution of units in the bedrock, and perhaps the pale grey gabbro has been locally transported eastward by glacial processes. Alternatively, perhaps there are larger-scale fold structures in this region, which are difficult to see because the terrain is so broken.

Stop 4.8 consists of more continuous outcrop of deformed gabbro, also containing augen-like clinopyroxene (augite) megacrysts akin to those described at Stop 4.7. However, deformation appears to be much stronger here than at Stops 4.6 and 4.7, and there are several areas of streaky, finely banded gabbro that have a distinctly mylonitic appearance. These fabrics likely result from intense transposition of inter-layered gabbro and dunite.



Figure 31. Large rotated clinopyroxene megacryst in strongly layered and foliated gabbro within the Moho region (Stop 4.7). This same feature was illustrated in the article by Malpas (1987).



Figure 32. Folds defined by leucocratic layers within the mafic–ultramafic cumulate sequence of the Moho region at Stop 4.9.

Stop 4.8 also reveals the first views to the south and southwest. The high hills to the south are part of the Gregory Plateau, consisting of the higher levels of the ophiolite suite, predominantly sheeted dykes and pillow lavas. These rock units are located on the other side of Trout River Pond, across a later fault, and are part of the North Arm Massif of the BOIC. (Further information is provided in the descriptions for Excursion 3). The valley to the southwest is occupied by the lower part of Trout River Pond, and lies mostly within mélanges that sit beneath the ultramafic rocks.

Stop 4.9: Fold Structures in the Moho Region (424488E, 5479579N)

Stop 4.9 is located a short distance south of Stop 4.8, and displays similar rock types and panoramic views. It also displays some folds developed within layered gabbro (Fig. 32). Some of the folds appear to be intrafolial structures, i.e. early folds that have been completely disrupted by later deformation. This is part of the process of transposition, which creates the strongly banded rocks seen at Stop 4.8 and elsewhere.

Stop 4.10: Transition Zone Beneath the Moho (424617E, 5479079N)

Stop 4.10 is a prominent grey area visible downhill from Stop 4.9. The walk from Stop 4.9 to Stop 4.10 passes back through the Moho again, and shows the same transition from layered foliated gabbros containing olivine-rich layers into feldspathic dunites containing plagioclase-rich layers. Towards Stop 4.10, there are also many blocks of harzburgite derived from the east, and the contrast between their yellow-brown weathering and the dark-brown weathering of the dunites becomes more obvious. There is no outcrop at Stop 4.10, which is dominated by large frost-heaved blocks of harzburgite.

The dunitic and harzburgitic rocks that sit beneath the Moho are part of the transition zone, which is believed to be derived from the cumulate crystal mush that developed at the base of the mafic magma chamber at the original spreading

centre. The lower parts of the transition zone consist largely of harzburgites that contain pods and dykes of dunitic and pyroxenitic composition. These are the same rock types that are exposed in the area around Winterhouse Brook (Excursion 2).

This is the turnaround point for the shorter, easier version of Excursion 4. If you do not wish to descend into the rocks of the upper mantle, simply retrace the route described above. The next section of the full hike involves some difficult walking in broken terrain; the smooth brown mountains of ultramafic rock ahead of you look like easy walking, but distance is very deceptive, as much of the surface is chaotically broken.

Stop 4.11: Harzburgite Containing Pyroxenitic and Dunitic Veins (425143E, 5479144N)

From Stop 4.10, the route heads almost due east, downhill at first, across a small stream, and then up a gentle slope on the other side. This walk goes from the dunitic rocks that lie east of the geophysical Moho and into the yellow-brown weathering harzburgites that dominate most of the Tablelands plateau. The petrological Moho, as distinct from the geophysical Moho, is crossed somewhere in this interval, but is a much more subtle feature because it is manifested largely by textures in the ultramafic rocks, rather than being an obvious compositional boundary.

Outcrops at Stop 4.11 consist of coarse-grained harzburgite cut by two generations of younger veins. Fine-grained, brown-weathering veins appear to be of dunitic composition, and are themselves cut by coarse-grained, pale green, pyroxenitic veins. The relationships of these veins to the host harzburgites resemble those observed near Winterhouse Brook (Excursion 2) but here there appears to be little or no superimposed deformation.

Stop 4.12: Harzburgite Hill (425963E, 5479954N)

Stop 4.12, located on the summit of a broad hill, consists entirely of harzburgites. It is almost 1 km from Stop 4.11 to Stop 4.12, and the walk is not exactly pleasant, as most of the route consists of large frost-heaved harzburgite blocks. This type of material is typical of much of the plateau on top of the Tablelands Ophiolite, which looks quite inviting from a distance. However, the landscape is surreal and extraterrestrial in appearance. The outcrops between Stop 4.11 and Stop 4.12 are all harzburgites, essentially identical to those at Stop 4.11. The hilltop is worth exploring for its views. This is the highest point on the hike (just over 650 m), but not the highest point in the Tablelands Ophiolite, which is located at 721 m, east of the head of Winterhouse Brook canyon. The highest point is only about 2 km southeast of here, but is broad and flat-topped, with inferior views, and probably not worth the grueling walk across the broken landscape. Table Mountain itself is a smaller peak of gabbro seen to the west northwest, but its elevation is a mere 520 m.

Stop 4.13: Canyon Access Point (425458E, 5481144N)

From Stop 4.12, head northwest across the seemingly endless plain of frost-heaved harzburgite to this spot, located above a



small canyon that descends the northern slope of Table Mountain. This canyon is also rather extraterrestrial and lifeless in appearance, totally lacking the vegetation that normally marks a watercourse. The canyon walls are loose scree slopes, and must be treated with caution. However, it is possible to descend into the canyon and up the opposite side, with care. After doing so, follow the stream for a few hundred metres. The walking by the stream is difficult, but most of the blocks are stable. Where the gradient of the stream suddenly increases, leave the valley again and follow the contours of the hillside to the northwest. From here, you can see Route 431 below, and can walk towards it directly. The route down is relatively easy if you manage to stay away from several areas dominated by large frost-heaved blocks. Upon reaching the road, walk westwards for about 1 km to return to the parking area (UTM coordinate 424650E, 5283050N) where the hiking route commenced.

ACKNOWLEDGEMENTS

This article, although long, is but a small sample of the scenic and geological attractions of Gros Morne National Park. Much of it started as part of a larger project initiated around 2001 to write a geological guidebook to the Park for visitors, in conjunction with the late Robert K. Stevens. Like many overambitious ideas, it was only partly completed, spending more than a decade in a filing cabinet and on obsolete floppy disks. Adapting this section for Geoscience Canada was far more time-consuming than I ever anticipated, but hopefully other parts of this project may yet see the light of day. Robert (Bob) Stevens played a crucial role in understanding the geology of western Newfoundland, and was the first to correctly assemble all the pieces of the Gros Morne jigsaw puzzle. His work focused on the sedimentary rocks and their fossils, but he noted many other things on many other topics, often preferring new questions to firm answers. His contribution to knowledge is not fully reflected in the number of publications that he authored. Bob was not the one who first introduced me to Gros Morne National Park, but it was through knowing and talking with Bob that I gained a better but still incomplete understanding of this amazing place.

I also acknowledge many useful general and detailed comments from reviewers John Waldron and Peter Cawood, which greatly improved a rather sloppy first draft. Brendan Murphy is also thanked for editorial handling of the paper on behalf of Geoscience Canada.

REFERENCES

- Baker, D.F., 1979, Geology and geochemistry of an alkali volcanic suite (Skinner Cove Formation) in the Humber Arm Allochthon, Newfoundland: Unpublished M.Sc. Thesis, Department of Geology, Memorial University, St. John's, NL, 314 p.
- Bédard, J.H., 2014, Ophiolites: Perspectives from field work in the Appalachians: *Elements*, v. 10, p. 87–88, <https://doi.org/10.2113/gselements.10.2.87>.
- Berger, A.R., Bouchard, A., Brookes, I.A., Grant, D.R., Hay, S.G., and Stevens, R.K., 1992, Geology, topology and vegetation, Gros Morne National Park, Newfoundland: Geological Survey of Canada, Miscellaneous Report 54; scale 1:150,000, <https://doi.org/10.4095/134066>.
- Bernoulli, D., 2001, Where did Gustav Steinmann see the Trinity? (Abstract): Geological Society of America, Annual Meeting, Boston, Massachusetts, November 2001, Abstract 71-0.
- Brongniart, A., 1821, Sur le gisement ou position relative des ophiolites, euphotides, jaspes, etc., dans quelques parties des Apennins: *Annales des Mines (Paris)*, v. 6, p. 177–238.
- Brookes, I.A., 1993, Table Mountain, Gros Morne National Park, Newfoundland: *Canadian Geographer*, v. 39, p. 69–75, <https://doi.org/10.1111/j.1541-0064.1993.tb01545.x>.
- Brookes, I.A., and Deardon, P., 1981, Guidebook for Excursion H: Geomorphology and plant geography, Gros Morne National Park. Canadian Association of Geographers, Annual Meeting 1981, Corner Brook, Newfoundland, Field Trip Guide, 68 p.
- Brown, P.A., 1978, Xonotlite: A new occurrence at Rose Blanche, Newfoundland: *Canadian Mineralogist*, v. 16, p. 671–672.
- Buddington, A.F., and Hess, H.H., 1937, Layered peridotite laccoliths in the Trout River area, Newfoundland: *American Journal of Science*, v. 33, p. 380–388.
- Calon, T.J., Dunsworth, S.D., and Suhr, G., 1988, Trip B8 - The Bay of Islands Ophiolite: Geological Association of Canada - Mineralogical Association of Canada, Annual Meeting 1988, St. John's, Newfoundland, Field Trip Guidebook, 92 p.
- Casey, J.F., Elthon, D.L., Siroky, F.X., Karson, J.A., and Sullivan, J., 1985, Geochemical and geological evidence bearing on the origin of the Bay of Islands and Coastal Complex ophiolites of western Newfoundland: *Tectonophysics*, v. 116, p. 1–40, [https://doi.org/10.1016/0040-1951\(85\)90220-3](https://doi.org/10.1016/0040-1951(85)90220-3).
- Cawood, P.A., 1991, Processes of ophiolite emplacement in Oman and Newfoundland, in Peters, T., Nicolas, A., and Coleman, R.G., eds., *Ophiolite Genesis and Evolution of the Oceanic Lithosphere: Petrology and Structural Geology*, v. 5, p. 501–516, https://doi.org/10.1007/978-94-011-3358-6_25.
- Cawood, P.A., and Suhr, G., 1992, Generation and obduction of ophiolites: Constraints from the Bay of Islands Complex, western Newfoundland: *Tectonics*, v. 11, p. 884–897, <https://doi.org/10.1029/92TC00471>.
- Cawood, P.A., McCausland, P.J.A., and Dunning, G.R., 2001, Opening Iapetus: Constraints from the Laurentian margin in Newfoundland: *Geological Society of America Bulletin*, v. 113, p. 443–453, [https://doi.org/10.1130/0016-7606\(2001\)113<0443:OICFTL>2.0.CO;2](https://doi.org/10.1130/0016-7606(2001)113<0443:OICFTL>2.0.CO;2).
- Church, W.R., and Stevens, R.K., 1971, Early Paleozoic ophiolite complexes of the Newfoundland Appalachians as mantle-oceanic crust sequences: *Journal of Geophysical Research*, v. 76, p. 1460–1466, <https://doi.org/10.1029/JB076i005p01460>.
- Coleman, R.G., 1967, Low-temperature reaction zones and alpine ultramafic rocks of California, Oregon and Washington: *United States Geological Survey, Bulletin* 1247, 49 p.
- Coleman, R.G., 1971, Plate tectonic emplacement of upper mantle peridotites along continental edges: *Journal of Geophysical Research*, v. 76, p. 1212–1222, <https://doi.org/10.1029/JB076i005p01212>.
- Colman-Sadd, S.P., Hayes, J.P., and Knight, I., 1990, Geology of the Island of Newfoundland, 1: 1 million scale: Newfoundland Department of Mines and Energy, Geological Survey, Map 90-01.
- Cooper, J.R., 1936, Geology of the southern half of the Bay of Islands Igneous Complex: Newfoundland Department of Natural Resources, Geological Section, Bulletin 4.
- Cooper, R.A., Williams, S.H., and Nowlan, G., 2001, Global stratotype section and point for the base of the Ordovician System: Episodes, v. 24, p. 19–23.
- Dalziel, I.W.D., Lawver, L.A., and Murphy, J.B., 2000, Plumes, orogenesis, and supercontinental fragmentation: *Earth and Planetary Science Letters*, v. 178, p. 1–11, [https://doi.org/10.1016/S0012-821X\(00\)00061-3](https://doi.org/10.1016/S0012-821X(00)00061-3).
- Desmurs, L., Manatschal, G., and Bernoulli, D., 2001, The Steinmann Trinity revisited: mantle exhumation and magmatism along an ocean-continent transition: the Platta nappe, eastern Switzerland, in Wilson, R.C.L., Whitmarsh, R.B., Taylor, B., and Froitzheim, N., eds., *Non-volcanic Rifting of Continental Margins: A Comparison of Evidence from Land and Sea*: Geological Society, London, Special Publications, v. 187, p. 235–266, <https://doi.org/10.1144/GSL.SP.2001.187.01.12>.
- Dewey, J.F., and Bird, J.M., 1971, Origin and emplacement of the ophiolite suite: Appalachian ophiolites in Newfoundland: *Journal of Geophysical Research*, v. 76, p. 3179–3206, <https://doi.org/10.1029/JB076i014p03179>.
- Dewey, J.F., and Casey, J.F., 2013, The sole of an ophiolite: the Ordovician Bay of Islands Complex, Newfoundland: *Journal of the Geological Society*, v. 170, p. 715–722, <https://doi.org/10.1144/jgs2013-017>.
- Dilek, Y., and Furnes, H., 2011, Ophiolite genesis and global tectonics: Geochemical and tectonic fingerprinting of ancient oceanic lithosphere: *Geological Society of America Bulletin*, v. 123, p. 387–411, <https://doi.org/10.1130/B30446.1>.
- Dilek, Y., and Furnes, H., 2014, Ophiolites and their origins: *Elements*, v. 10, p. 93–100, <https://doi.org/10.2113/gselements.10.2.93>.
- Dilek, Y., Moores, E.M., Elthon, D., and Nicolas, A., editors, 2000, Ophiolites and oceanic crust: New insights from field studies and the Ocean Drilling Program: *Geological Society of America Special Paper*, v. 349, 552 p.
- Dunning, G.R., and Krogh, T.E., 1985, Geochronology of ophiolites of the Newfoundland Appalachians: *Canadian Journal of Earth Sciences*, v. 22, p. 1659–1670, <https://doi.org/10.1139/c85-174>.
- Gass, I.G., 1968, Is the Troodos massif of Cyprus a fragment of Mesozoic ocean floor? *Nature*, v. 220, p. 39–42, <https://doi.org/10.1038/220039a0>.
- Goodenough, K.M., Thomas, R.J., Styles, M.T., Schofield, D.I., and MacLeod, C.J., 2014, Records of ocean growth and destruction in the Oman-UAE ophiolite: *Elements*, v. 10, p. 109–114, <https://doi.org/10.2113/gselements.10.2.109>.
- Harland, W.B., and Gayer, R.A., 1972, The Arctic Caledonides and earlier oceans: *Geological Magazine*, v. 109, p. 289–314, <https://doi.org/10.1017/S0016756800037717>.
- Hess, H.H., 1938, A primary peridotite magma: *American Journal of Science*, v. 35,

- p. 321–344, <https://doi.org/10.2475/ajs.s5-35.209.321>.
- Hild, M.H., 2012, *Geology of Newfoundland: Touring through time at 48 scenic sites*: Boulder Publications, St. John's, NL, 256 p.
- Hodych, J.P., Cox, R.A., and Košler, J., 2004, An equatorial Laurentia at 550 Ma confirmed by Grenvillian inherited zircons dated by LAM ICP-MS in the Skinner Cove volcanics of western Newfoundland: Implications for inertial interchange true polar wander: *Precambrian Research*, v. 129, p. 93–113, <https://doi.org/10.1016/j.precamres.2003.10.012>.
- Ingerson, F.E., 1935, Layered peridotitic laccoliths of the Trout River area, Newfoundland: *American Journal of Science*, v. 29, p. 422–440, <https://doi.org/10.2475/ajs.s5-29.173.422>.
- James, N.P., Stevens, R.K., Barnes, C.R., and Knight, I., 1989, Evolution of a lower Paleozoic continental-margin carbonate platform, northern Canadian Appalachians, in Crevello, P.D., Wilson, J.L., Sarg, J.F., and Read, J.F., eds., *Controls on Carbonate Platforms and Basin Development*: Society of Economic Paleontologists and Mineralogists (SEPM) Special Publication, v. 44, p. 123–146, <https://doi.org/10.2110/pec.89.44.0123>.
- Jébrak, M., and Marcoux, E., 2015, *Geology of Mineral Resources*: Geological Association of Canada, St. John's, NL, 668 p.
- Jenner, G.A., Dunning, G.R., Malpas, J., Brown, M., and Brace, T., 1991, Bay of Islands and Little Port complexes revisited: Age, geochemical and isotopic evidence confirm suprasubduction-zone origin: *Canadian Journal of Earth Sciences*, v. 28, p. 1635–1652, <https://doi.org/10.1139/e91-146>.
- Malpas, J., 1979, Two contrasting trondhjemite associations from transported ophiolites in western Newfoundland: Initial Report, in Barker, F., ed., *Trondhjemites, Dacites and Related Rocks: Developments in Petrology*, v. 6, p. 465–487, <https://doi.org/10.1016/B978-0-444-41765-7.50020-4>.
- Malpas, J., 1987, The Bay of Islands ophiolite; A cross-section through Paleozoic crust and mantle in western Newfoundland, in Roy, D.C., ed., *DNAG Centennial Field Guides: Northeastern Section of the Geological Society of America*, v. 5, <https://doi.org/10.1130/0-8137-5405-4.451>.
- Miyashiro, A., 1973, The Troodos ophiolite was probably formed in an island arc: *Earth and Planetary Science Letters*, v. 19, p. 218–224, [https://doi.org/10.1016/0012-821X\(73\)90118-0](https://doi.org/10.1016/0012-821X(73)90118-0).
- Morrill, P.L., Brazelton, W.J., Kohl, L., Rietze, A., Miles, S.M., Kavanagh, H., Schrenk, M.O., Ziegler, S.E., and Lang, S.Q., 2014, Investigations of potential microbial methanogenic and carbon monoxide utilization pathways in ultrabasic reducing springs associated with present-day continental serpentinization: The Tablelands, NL, Canada: *Frontiers in Microbiology*, <https://doi.org/10.3389/fmicb.2014.00613>.
- Pearce, J.A., 2014, Immobile element fingerprinting of ophiolites: *Elements*, v. 10, p. 101–108, <https://doi.org/10.2113/gselements.10.2.101>.
- Penrose Field Conference Participants, 1972, Ophiolites: *Geotimes*, v. 17, p. 24–25.
- Salisbury, M.H., and Christensen, N.I., 1978, The seismic velocity structure of a traverse through the Bay of Islands Ophiolite Complex, Newfoundland, An exposure of oceanic crust and upper mantle: *Journal of Geophysical Research*, v. 83, p. 805–817, <https://doi.org/10.1029/JB083iB02p00805>.
- Scott, D.J., Helmstaedt, H., and Bickle, M.J., 1992, Purtuniqu ophiolite, Cape Smith belt, northern Quebec, Canada: A reconstructed section of early Proterozoic oceanic crust: *Geology*, v. 20, p. 173–176, [https://doi.org/10.1130/0091-7613\(1992\)020<0173:POCSBN>2.3.CO;2](https://doi.org/10.1130/0091-7613(1992)020<0173:POCSBN>2.3.CO;2).
- Searle, M., and Cox, J., 1999, Tectonic setting, origin and obduction of the Oman ophiolite: *Geological Society of America Bulletin*, v. 111, p. 104–122, [https://doi.org/10.1130/0016-7606\(1999\)111<0104:TSAOO>2.3.CO;2](https://doi.org/10.1130/0016-7606(1999)111<0104:TSAOO>2.3.CO;2).
- Searle, M., and Stevens, R.K., 1984, Obduction processes in ancient, modern and future ophiolites: *Geological Society, London, Special Publications*, v. 13, p. 303–319, <https://doi.org/10.1144/GSL.SP.1984.013.01.24>.
- Smith, C.H., 1958, Bay of Islands igneous complex, western Newfoundland: *Geological Survey of Canada, Memoir 290*, 132 p., <https://doi.org/10.4095/123930>.
- Snelgrove, A.K., 1934, Chromite deposits of Newfoundland: Newfoundland Department of Natural Resources, Geological Section, Bulletin 1.
- Steinmann, G., 1927, Die ophiolitischen zonen in den mediterranen kettengebirgen: 14e Congrès Géologique Internationale, Madrid, 1926, *Graficas Reunidas*, v. 2, p. 637–667.
- Stevens, R.K., 1965, *Geology of the Humber Arm area, west Newfoundland*: Unpublished M.Sc. Thesis, Memorial University, St. John's, NL, 121 p.
- Stevens, R.K., 1970, Cambro–Ordovician flysch sedimentation and tectonics in western Newfoundland and their possible bearing on a proto-Atlantic ocean: *Geological Association of Canada, Special Paper 7*, p. 165–177.
- Stevens, R.K., 1988, Ophiolite oddities: preliminary notices of nephrite jade, Sr-aragonite, troilite and hyperalkaline springs from Newfoundland (Abstract): *Geological Association of Canada, Program with abstracts*, v. 13, p. 118.
- Stevens, R.K., Calon, T., and Hingston, R., 2003, From the intertidal zone to the upper mantle - the amazing geology of Gros Morne National Park: *Geological Association of Canada, Newfoundland and Labrador Section, Fall Field Trip (2003) Guidebook*, 38 p.
- Suhr, G., 1991, *Structural and magmatic history of upper mantle peridotites in the Bay of Islands Complex, Newfoundland*: Unpublished Ph.D. Thesis, Memorial University, St. John's, NL, 512 p.
- Suhr, G., and Cawood, P.A., 1993, Structural history of ophiolite obduction, Bay of Islands, Newfoundland: *Geological Society of America Bulletin*, v. 105, p. 399–410, [https://doi.org/10.1130/0016-7606\(1993\)105<0399:SHOOOB>2.3.CO;2](https://doi.org/10.1130/0016-7606(1993)105<0399:SHOOOB>2.3.CO;2).
- Szponar, N., Brazelton, W.J., Schrenk, M.O., Bower, D.M., Steele, A., and Morrill, P.L., 2013, Geochemistry of a continental site of serpentinization, the Tablelands Ophiolite, Gros Morne National Park: A Mars analogue: *Icarus*, v. 224, p. 286–296, <https://doi.org/10.1016/j.icarus.2012.07.004>.
- Wager, L.R., and Deer, W.A., 1939, The petrology of the Skaergaard Intrusion, Kangerdlugssuaq, East Greenland: København, C.A. Reitzels forlag, Bulletin 105.
- Williams, H., 1964, The Appalachians in northeastern Newfoundland; A two-sided symmetrical system: *American Journal of Science*, v. 262, p. 1137–1158, <https://doi.org/10.2475/ajs.262.10.1137>.
- Williams, H., 1979, Appalachian Orogen in Canada: *Canadian Journal of Earth Sciences*, v. 16, p. 792–807, <https://doi.org/10.1139/e79-070>.
- Williams, H., 1995, editor, *Geology of the Appalachian-Caledonian Orogen in Canada and Greenland*: Geological Survey of Canada, *Geology of Canada Series*, v. 6, 944 p., <https://doi.org/10.1130/DNAG-GNA-F1>.
- Williams, H., and Cawood, P.A., 1989, *Geology, Humber Arm Allochthon, Newfoundland*: Geological Survey of Canada, "A" Series Map 1678A, scale 1:250,000, <https://doi.org/10.4095/126990>.
- Williams, H., and Malpas, J., 1972, Sheeted dykes and brecciated dyke rocks within transported igneous complexes Bay of Islands, western Newfoundland: *Canadian Journal of Earth Sciences*, v. 9, p. 1216–1229, <https://doi.org/10.1139/e72-105>.
- Williams, H., and Smyth, W.R., 1973, Metamorphic aureoles beneath ophiolite suites and alpine peridotites: tectonic implications with west Newfoundland examples: *American Journal of Science*, v. 273, p. 594–621, <https://doi.org/10.2475/ajs.273.7.594>.
- Williams, H., and Stevens, R.K., 1974, The ancient continental margin of eastern North America, in Burk, C.A., and Drake, C.L., eds., *The Geology of Continental Margins*: Springer, Berlin, Heidelberg, p. 781–796, https://doi.org/10.1007/978-3-662-01141-6_58.
- Wilson, J.T., 1966, Did the Atlantic close and then re-open?: *Nature*, v. 211, p. 676–681, <https://doi.org/10.1038/211676a0>.
- Yang, J.-S., Robinson, P.T., and Dilek, Y., 2014, Diamonds in ophiolites: *Elements*, v. 10, p. 127–130, <https://doi.org/10.2113/gselements.10.2.127>.

Received May 2019

Accepted as revised August 2019



APPENDIX: GLOSSARY OF SELECTED TERMS

The following is included to assist readers with limited knowledge of geological terms beyond basic concepts. It is not exhaustive, but any undefined terms in the text can probably be understood with reference to one of several popular geoscience directories.

Accretion (tectonic): a large-scale process in which continental fragments and island arcs are swept in towards larger continents by plate tectonics, and then joined to them. Characteristic of *convergent margins*.

Allochthon, allochthonous: packages of rock units that formed in another place but were then transported by the forces of plate tectonics to the location in which they are now found. Opposite of *autochthonous*.

Anorthositic, anorthosite: igneous rock consisting largely of Ca–Na-rich feldspar (plagioclase) with very small amounts of olivine and/or pyroxene. Often a *cumulate rock*.

Autochthon, autochthonous: packages of rock units found in essentially the same location where they originally formed. Opposite of *allochthonous*.

Convergent margin: a location where tectonic plates are moving toward each other and one plate descends beneath the other by the process of *subduction*.

Cumulate rock: igneous rock formed by the accumulation of minerals that have crystallized from a liquid magma. See also *fractionation*.

Diachronous: adjective describing a process or event that occurs at different times in different places, but often in a rather orderly progressive fashion. For example, the human migration out of Africa to other continents was diachronous. It is the opposite of synchronous.

Dunite: a type of peridotite consisting mostly of the mineral olivine.

Fractionation: the process by which a liquid magma changes composition as different minerals crystallize and separate from the remaining liquid magma by some process of accumulation, to make a cumulate rock.

Global Stratotype Section: the location where a geologic boundary is defined between rock layers or with reference to fossil assemblages in the rock layers, as a reference point for the geological time scale.

Gondwanaland: a large Paleozoic continental assembly including much of Africa, South America, Australia and Antarctica, along with parts of Asia. The Appalachian–Caledonian Orogenic Belt was created as part of the process that joined Gondwanaland to North America to create the supercontinent called Pangaea ('all land') at the end of the Paleozoic Era.

Harzburgite: a type of peridotite consisting mostly of the mineral olivine and orthopyroxene, which is a Mg–Fe-rich pyroxene.

Hydrothermal: processes involving hot waters and aqueous solutions circulating in Earth's subsurface.

Intrafolial fold: an isoclinal fold that is disconnected from adjoining layers; normally an indication of a larger and older fold structure that has been torn apart by continued deformation.

Isoclinal fold: a tight fold structure in which the two fold limbs are essentially parallel; a common product of intense deformation under ductile conditions.

Mélange: a mixed up rock, usually deformed and badly fractured, in which blocks of different rock types of varied sizes are contained within a matrix of sedimentary origin, usually in a rather chaotic fashion.

Metamorphic aureoles: zones of metamorphic rocks produced through thermal effects related to the proximity of high-temperature liquid magmas, or hot solid rocks such as mantle peridotites.

Metasomatism: a process by which the bulk composition of a rock unit is changed, normally through reactions with hydrothermal fluids percolating through it.

Mohorovičić Discontinuity (Moho): a region in the crust where geophysical studies indicate an increase in the velocity of seismic waves, consistent with a sudden increase in the density of rock units at or near the boundary between the Earth's crust and the upper mantle region.

Mylonite zone: a belt of fine-grained and intensely banded rocks that geologists usually consider to indicate relative motion of adjoining rock units under high-temperature conditions that lead to ductile behaviour of minerals in the area caught between those rock units.

Obduction: process that places rocks of the oceanic crust above those of the continental crust; the opposite of *subduction*.

Paleoproterozoic: a part of the vast Precambrian time period, extending from 2500 to 1600 million years ago.

Partial melting: process by which some of the minerals in a silicate rock melt due to high temperatures, creating a liquid magma that has a different composition to the starting material and the residual material left behind. See also *fractionation*.

Peridotite: general term for ultramafic igneous rocks consisting mostly of the minerals olivine and various pyroxenes.

Pillow lava: a submarine volcanic rock in which basalt magma chills rapidly into blob-like structures, much as hot honey would congeal on contact with ice-cold water.

Pyroxenite: a rock consisting largely of pyroxene minerals; orthopyroxene and clinopyroxene minerals may occur alone or in combination.

Retrogressed, retrogression: changes in the mineral assemblage found in an igneous rock to water-bearing minerals characteristic of lower temperatures of formation, but without any significant change in the rock's bulk composition.

Serpentinites, serpentinization: rocks that consist largely of serpentine-group minerals, which are soft flaky silicate minerals that resemble snakeskins when weathered. Serpentine minerals are produced by reactions between primary igneous minerals in ultramafic rocks and percolating waters or *hydrothermal* fluids.

Subduction: when one tectonic plate (usually oceanic) descends beneath another, usually resulting in earthquakes and abundant volcanic activity. A characteristic process at *convergent margins*.

Syncline, synclinal: a gentle bowl-like fold structure of rock units, such as thinly-bedded sedimentary rocks or km-scale units of components in an ophiolite complex.

Tethys Ocean: a largely vanished ocean, which once separated Africa and parts of Asia from Europe. The Mediterranean sea is all that now remains of the Tethys Ocean.

Transposition: a process that creates intensely banded rocks such as mylonites, or the banded ultramafic rocks described from the mantle regions of ophiolite complexes.

Trondhjemite: an unusual type of igneous rock that resembles a granite, but consists largely of quartz and the Na-rich feldspar mineral albite (most granites also contain K-rich feldspar minerals such as orthoclase or microcline).

Wehrlite: a type of peridotite consisting mostly of the mineral olivine and clinopyroxene, which is a Ca–Mg-rich pyroxene variety.