Index to Volume 9

Citation refers to $\underline{\text{Issue}}$ Number and $\underline{\text{Page}}$ Number

<u>Nu</u>	mber	Page	. <u>Þ</u>	Number	Page
Acadian deformation in Maine-New Brunswick, the age of the: DONOHOE	3	78	COK, A., see SWIFT	2	37
Age, the, of the Acadian deformation in	3	70	Continental margins, mesa: interdis- ciplinary approach to environ-		
Maine-New Brunswick: DONOHOE	3	78	mental analysis of: SWIFT	2	37
AKPATI, B.N., Observations on the effects	•	, ,	mental analysis of: Swift	2	37
of islets and shoals on salinity			DALRYMPLE, R.W., preliminary investi-		
distribution in an estuarine regime	2	59	gations of an intertidal sand		
Analysis, environmental, of continental			body, Cobequid Bay, Bay of Fundy	1	21
margins, mesa: interdisciplinary			DALRYMPLE, R.W., sediment texture and		
approach to: SWIFT	2	37	transport studies in an intertidal	-	
Analysis, facies, of the Pennsylvania			environment: A progress report	2	45
Pictou Group in New Brunswick,			Deformation, Acadian, of the, in Maine-		
Stratigraphy, sediment dispersal and	:		New Brunswick, the age: DONOHOE	3	78
VAN DE POLL	3	72	Deposits, stream, braided, Devonian in		
Approach, interdisciplinary, to environ-			the Battery Point Formation, Gaspé		
mental analysis of continental			Est, Quebec: CANT	1	13
margins, mesa: SWIFT	2	37	Devonian braided stream deposits in the		
Differ by the state of the stat			Battery Point Formation, Gaspé Est		
Baffin Bay, a late Quaternary strati-			Quebec: CANT	1	13
graphic marker in the central basin	2	63	Dispersal, sediment, and facies analysi	s	
of: PIPER Basin, central, of Baffin Bay, a late	2	62	of the Pennsylvania Pictou Group		
Quaternary stratigraphic marker in			in New Brunswick, stratigraphy: VAN DE POLL	3	72
the: PIPER	2	62	Distribution, salinity, in an estuarine		12
Battery Point Formation, Gaspé Est,	-	02	regime, observations on the effect		
Quebec, Devonian braided stream			of islets and shoals on: AKPATI	2	59
deposits in the: CANT	1	13	DONOHOE, H.V., Jr., and PAJARIE, G.,	-	3,
Bay, Baffin, a late Quaternary strati-	_		The age of the Acadian deformation		
graphic marker in the central basin			in Maine-New Brunswick	3	78
of: PIPER	2	62	DRAKE, D., see SWIFT	2	37
Bay, Cobequid, Bay of Fundy, preliminary					
investigations of an intertidal sand			Effects of islets and shoals on salinit	У	
body: DALRYMPLE	1	21	distribution in an estuarine regim	e,	
Fay of Fundy, preliminary investigations			observations on the: AKPATI	2	59
of an intertidal sand body, Cobequid			Environmental analysis of continental		
Bay: DALRYMPLE	1	21	margins, mesa: interdisciplinary		
Body, sand, intertidal Cobequid Bay, Bay			approach to: SWIFT	2	37
of Fundy, preliminary investigations	,	21	Environmental intertidal: A progress		
of an: DALRYMPLE	1	21	report, sediment texture and	2	45
Briaded stream deposits, Devonian, in the Battery Point Formation, Gaspé Est,			transport studies in an: DALRYMPLE	. 2	43
Ouebec: CANT	1	13	Environments, sedimentary: A study for students, a re-examination of the		
Brief summary of the Late Precambrian	_	13	silt/clay ratios as indicators of:		
rocks in the Caledonian Highlands of			PELLETIER	1	1
southern New Brunswick, a:			Estuarine regime, observations on the		
RUITENBERG	3	83	effects of islets and shoals on		
			salinity distribution in an: AKPAT	I 2	59
Caledonian Highlands of southeastern New					
Brunswick, a brief summary of the			Facies analysis of the Pennsylvania Pic		
Late Paleozoic in the: RUITENBERG	3	83	Group in New Brunswick, stratigrap	_	
CANT, D.J., Devonian braided stream			sediment dispersal and: VAN DE POL	L 3	72
deposits in the Battery Point For-	-	1.2	Formation, Battery Point, Gaspé Est,		
mation, Gaspé Est, Quebec	1	13	Quebec, Devonian braided stream	,	1.2
Cape Breton Island, Precambrian rocks of:	2	100	deposits in the: CANT	1 2	13 37
WIEBE Cape Breton Island, southeast, Nova	3	100	FREELAND, G., see SWIFT Fundy, Bay of, preliminary investigation		31
Scotia, Pre-Carboniferous structural			of an intertidal sand body,	J	
history of: HELMSTAEDT	3	88	Cobequid Bay: DALRYMPLE	1	21
Central basin of Baffin Bay, a late	5	00	conequia bay. Dankini in	-	-1
Quaternary stratigraphic marker in			Gaspé Est, Quebec, Devonian braided str	eam	
the: PIPER	2	62	deposits in the Battery Point For-		
Cobequid Bay, Bay of Fundy, preliminary	•		mation: CANT	1	13
investigation of an intertidal sand			GILES, P.S., see RUITENBERG, A.A.	3	83
body: DALRYMPLE	1	21	Group, Pictou, Pennsylvania, in New		

Number	Page	Number	Page
Brunswick, stratigraphy, sediment dispersal and facies analysis of		summary of the Late Paleozoic rocks in the Caledonian Highlands	
the: VAN DE POLL 3	72	of: RUITENBERG 3	83
HATCHER, P., see SWIFT, D. 2 HELMSTAEDT, H., and TELLA, S., Pre-	37	New Brunswick, stratigraphy, sediment dispersal and facies analysis of the Pennsylvania Pictou Group in:	
Carboniferous structural history of southeast Cape Breton Island, Nova Scotia 3	00	VAN DE POLL 3 Nova Scotia, Pre-Carboniferous structural	7,2 .
Scotia 3 Highlands, Caledonian, of southeastern New Brunswick, a brief summary of	88	history of southeast Cape Breton Island: HELMSTAEDT 3	88
the Late Paleozoic rocks in the: RUITENBERG 3	83	Observations on the effects of islets and shoals on salinity distribution in	
History, structural, Pre-Carboniferous, of southeast Cape Breton Island,		an estuarine regime: AKPATI 2	59
Nova Scotia: HELMSTAEDT 3	88	PAJARIE, G., see DONOHOE 3 Paleozoic rocks in the Caledonian High-	78
Indicators of sedimentary environments: A study for students, a re-exam-		lands of southeastern New Brunswick, a brief summary of the Late:	
<pre>ination of the use of the silt/clay ratios: PELLETIER 1</pre>	1	RUITENBERG 3 PELLETIER, B.R., A re-examination of the	83
Interdisciplinary approach to environ- mental analysis of continental		use of the silt/clay ratios as indicators of sedimentary environ-	_
margins, mesa: SWIFT 2 Intertidal environment: A progress report, sediment texture and transport	37	ments: A study for students l Pennsylvania Pictou Group in New Brunswick, stratigraphy, sediment dispersal	1
studies in an: DALRYMPLE 2 Intertidal sand body, Cobequid Bay, Bay	45	and facies analysis of the: VAN DE POLL 3	72
of Fundy, preliminary investigations of an: DALRYMPLE	21	Pictou Group, Pennsylvania, in New Brunswick, stratigraphy, sediment	
Investigations, preliminary, of an inter- tidal sand body, Cobequid Bay, Bay of Fundy: DALRYMPLE 1	21	dispersal and facies analysis of the: VAN DE POLL 3 PIPER, D.J.W., A late Quaternary strat-	72
Island, Cape Breton, Precambrian rocks of: WIEBE 3	100	igraphic marker in the central basin of Baffin Bay 2	62
Island, Cape Breton, southeast, Nova Scotia, Pre-Carboniferous structural history of: HELMSTAEDT 3	88	Pre-Carboniferous structural history of southeast Cape Breton Island, Nova Scotia: HELMSTAEDT 3	88
Islets and shoals on salinity distribution in an estuarine regime, observations		Precambrian rocks of Cape Breton Island: WIEBE 3	100
on the effects of: AKPATI 2	59	Preliminary investigations of the inter- tidal sand body, Cobequid Bay, Bay of	100
KELLER, G., see SWIFT 2	37	Fundy: DALRYMPLE 1	21
Late Paleozoic rocks in the Caledonian Highlands of southeastern New		Quaternary, late, a, stratigraphic marker in the central basin of	
Brunswick, a brief summary of the: RUITENBERG 3	83	Baffin Bay: PIPER 2 Quebec, Devonian braided stream deposits	62
Late, a, Quaternary stratigraphic marker in the central basin of Baffin Bay: PIPER 2	62	in the Battery Point Formation, Gaspé Est: CANT l	13
LAVELLE, W., see SWIFT 2 McCUTCHEON, S.R. see RUITENBERG 3	37 83	Ratios, silt/clay, as indicators of sedi-	
McKINNEY, T., see SWIFT 2	37	mentary environments: A study for students, a re-examination of the	
Maine-New Brunswick, the age of the Acadian deformation in: DONOHOE 3 Margins, continental, mesa: interdis-	78	use of the: PELLETIER l Re-examination of the use of the silt/ clay ratios as indicators of sedi-	1
ciplinary approach to environmental analysis of: SWIFT 2	37	mentary environments: A study for students, a: PELLETIER $f 1$	1
Marker, stratigraphic, Quaternary, late, a, in the central basin of Baffin Bay: PIPER 2	62	Regime, estuarine, observations on the effects of islets and shoals on salinity distribution in an: AKPATI 2	59
Mesa: interdisciplinary approach to	02	Rocks, Paleozoic, Late, in the Caldeonian	JJ
environmental analysis of continental margins: SWIFT 2	37	Highlands of southeastern New Brunswick, a brief summary of the: RUITENBERG 3	83
NELSEN, T., see SWIFT 2 New Brunswick, southeastern, a brief	37	Rocks, Precambrian, of Cape Breton Island: WIEBE 3	100

Nu	mber	Page		Number	Page
RUITENBERG, A.A., GILES, P.S., VENUGOPAL, D.V., and McCUTCHEON, S.R., A brief			ciplinary approach to environ- mental analysis of continental		
summary of the Late Precambrian			margins	2	37
rocks in the Caledonian Highlands of southern New Brunswick	3	83	TELLA, S., see HELMSTAEDT, H. Texture, sediment, and transport	3	88
Salinity distribution in an estuarine regime, observations on the effects			studies in an intertidal environ ment: A progress report: DALRYM		45
of islets and shoals on: AKPATI Sand body, intertidal, Cobequid Bay, Bay of Fundy, preliminary investigations	2	59	Transport studies in an intertidal environment: A progress report, sediment texture and: DALRYMPLE	2	45
of an: DALRYMPLE	1	21			
Sedimentary environments: A study for students, a re-examination of the use of the silt/clay ratios as indicators			Use of the silt/clay ratios as indicated of sedimentary environments: A study for students, a re-examinated for students.		
of: PELLETIER	1	1	of the: PELLETIER	1	1
Sediment dispersal and facies analysis of			VIIV DE DOLL II II Charlismanha godi	imant	
the Pennsylvania Pictou Group in New Brunswick, stratigraphy: VAN DE POLL	3	72	VAN DE POLL, H.W., Stratigraphy, sed: dispersal and facies analysis o		
Sediment texture and transport studies in	ŭ	, _	the Pennsylvania Pictou Group in	n	70
an intertidal environment: A progress	_	4.5	New Brunswick	3 3	72 83
report: DALRYMPLE Shoals on salinity distribution in an	2	4 5	VENUGOPAL, D.V., see RUITENBERG	3	03
estuarine regime, observations on the			WIEBE, R.A., Precambrian rocks of Cap		
effects of islets and: AKPATI	2	59	Breton Island	3	100
Silt/clay ratios as indicators of sed- imentary environments: A study for					
students, a re-examination of the use					
of the: PELLETIER	1	1			
Southeast Cape Breton Island, Nova Scotia, Pre-Carboniferous structural history					
of: HELMSTAEDT	3	88			
Southeastern New Brunswick, a brief summary of the Late Paleozoic rocks					
in the Caledonian Highlands of: RUITENBERG	3	83			
Stratigraphic, Quaternary, late, a marker					
in the central basin of Baffin Bay: PIPER	2	62			
Stratigraphy, sediment dispersal and facies		Ü.			
analysis of the Pennsylvania Pictou	2	70			
Group in New Brunswick: VAN DE POLL Stream deposits, braided, Devonian, in	3	72			
the Battery Point Formation, Gaspé					
Est, Quebec: CANT Structural history, Pre-Carboniferous, of	1	13			
southeast Cape Breton Island, Nova					
Scotia: HELMSTAEDT	3	88			
STUBBLEFIELD, W., see SWIFT Students, a re-examination of the use of	2	37			
the silt/clay ratios as indicators					
of sedimentary environments: A study	1	1			
for: PELLETIER Study, a, for students, a re-examination	_	1			
of the use of the silt/clay ratios as					
<pre>indicators of sedimentary environ- ments: PELLETIER</pre>	1	1			
Studies, transport, in an intertidal	_	-			
environment: A progress report,	2	45			
sediment texture and: DALRYMPLE Summary, a brief, of the Late Paleozoic	2	45			
rocks in the Caledonian Highlands of					
southeastern New Brunswick:	3	83		•	
RUITENBERG SWIFT, D., FREELAND, G., DRAKE, D.,	3	03			
HATCHER, P., KELLER, G., LAVELLE, W.,					
NELSEN, T., STUBBLEFIELD, W., COK, A. and McKINNEY, T., Mesa: interdis-	r				