

Maritime Sediments and Atlantic Geology

Vol. 20

APRIL, 1984

No. 1

Systematic ichnology of the Middle Ordovician Trenton Group. St. Lawrence Lowland, eastern Canada

D. Fillion and R.K. Pickerill

Department of Geology, University of New Brunswick, Fredericton, N.B. E3B 5A3

Carbonate sediments of the upper Middle Ordovician Trenton Group between Montreal and Quebec City in the St. Lawrence Lowland, eastern Canada, contain a diverse and abundant trace fossil assemblage consisting of *Arenicolites* sp., ?*Calycraterion* sp., *Chondrites* spp., *Circulichnis montanus*, *Clematischnia* sp., ?*Conostichnus* sp., *Cruziana problematica*, *Cruziana* sp., cf. *Diplichnites* sp., *Furculosus carpathicus*, *Helminthopsis hieroglyphica*, *Helminthopsis* sp., *Oichnus paraboloides*, *Palaeophycus tubularis*, *Palaeophycus* sp., ?*Plagiogmus* sp., *Planolites beverleyensis*, *P. montanus*, *Planolites* sp., ?*Rhizocorallium* cf. *R. irregulare*, ?*Rosselia* sp., *Scalarituba missouriensis*, *Scolicia* sp., *Skolithos linearis*, *Skolithos* sp., *Teichichnus rectus*, *Teichichnus* sp., *Trichichnus* sp., *Trypanites weisei*, *Vermiforichnus clarkel* and *Zoophycos* sp. as well as informally diagnosed loop, oblique and pronged burrows and bryozoan borings. Of these forms, only *Chondrites* spp., *Palaeophycus tubularis*, *Palaeophycus* sp., *Planolites* spp., *Teichichnus* spp. and *Trypanites weisei* are abundant; the remainder are rare to only moderately common. Nevertheless, in this paper we describe all the trace fossils in detail and in doing so attempt to resolve several current and controversial problems of nomenclature regarding certain ichnogenera.

Sediments of the Trenton Group were deposited initially in lagoons followed in turn by offshore "bar", shallow and, finally, deeper offshore shelf environments. The trace fossils do not exhibit significant variation with respect to these broad depositional regimes and, instead, each environment is characterized by assemblages typical of the *Cruziana* ichnofacies as recognized in clastic sequences. The major factor primarily limiting the spatial and temporal trace fossil distribution was substrate and its primary characteristics, in turn a reflectance of food availability and environmental energy levels.

Dans les Basses Terres du St-Laurent de l'Est canadien, les calcaires du Groupe de Trenton (Ordovicien moyen supérieur) renferment, entre Montréal et Québec, un assemblage d'ichnofossiles abondants et variés comprenant: *Arenicolites* sp., ?*Calycraterion* sp., *Chondrites* spp., *Circulichnis montanus*, *Clematischnia* sp., ?*Conostichnus* sp., *Cruziana problematica*, *Cruziana* sp., cf. *Diplichnites* sp., *Furculosus carpathicus*, *Helminthopsis hieroglyphica*, *Helminthopsis* sp., *Oichnus paraboloides*, *Palaeophycus tubularis*, *Palaeophycus* sp., ?*Plagiogmus* sp., *Planolites beverleyensis*, *P. montanus*, *Planolites* sp., ?*Rhizocorallium* cf. *R. irregulare*, ?*Rosselia* sp., *Scalarituba missouriensis*, *Scolicia* sp., *Skolithos linearis*, *Skolithos* sp., *Teichichnus rectus*, *Teichichnus* sp., *Trichichnus* sp., *Trypanites weisei*, *Vermiforichnus clarkel* et *Zoophycos* sp. On y trouve également des galeries en boucle et fourchues, des terriers obliques ainsi que des perforations de bryozoaires, tous identifiés de façon informelle. De toutes ces formes, seules les suivantes sont présentes en abondance: *Chondrites* spp., *Palaeophycus tubularis*, *Palaeophycus* sp., *Planolites* spp., *Teichichnus* spp., et *Trypanites weisei*; les autres sont rares ou peu communes. Néanmoins, nous décrivons en détail toutes les traces fossiles et tentons par le fait même de résoudre plusieurs problèmes de nomenclature chez certains ichnogènes qui font à l'heure actuelle l'objet de débats.

Le dépôt des sédiments du Groupe de Trenton s'est initié dans un milieu de lagune puis de barre, de plate-forme de haut-fond et, finalement, de plate-forme d'eau profonde. Les traces fossiles ne varient pas de façon significative d'un environnement de déposition à l'autre. On remarque plutôt que chaque environnement se caractérise par des assemblages typiques de l'ichnofacies à *Cruziana* tel qu'on l'observe dans les séquences détritiques. Le substrat et ses caractéristiques de base, qui reflètent la disponibilité de la nourriture et les conditions énergétiques du milieu, constituent l'élément majeur qui a, avant tout, limité la distribution spatiale et temporelle des traces fossiles.

[Traduit par le journal]

INTRODUCTION

Detailed and systematic trace fossil studies of carbonate rocks, particularly of Lower Paleozoic age, are far outnumbered by comparable studies in their clastic counterparts. In part this is because paleontologists are generally attracted to carbonates by their normally

abundant and well-preserved body fossils and are generally frustrated by the typically poor preservation, usually as a result of carbonate cementation and recrystallization, of their contained trace fossils. Yet many Lower Paleozoic carbonate sequences abound in trace fossils of various kinds and with careful and detailed analysis their systematics

can be resolved at the ichnogenic and even the ichnospecific level.

In this paper we describe the systematics of a suite of trace fossils contained in carbonates of the upper Middle Ordovician Trenton Group exposed between Montreal and Quebec City in the St. Lawrence Lowland, eastern Canada (Fig. 1). This study is one of the first of its kind in Lower Paleozoic carbonate sequences and highlights several controversial and taxonomical problems regarding certain ichnogenera. A companion paper (Fillion et al. 1984) has evaluated their spatial and temporal distributional patterns and primary controlling parameters within the sequence but, until now, the systematics remain unpublished despite the many recordings by previous authors of 'burrows', 'tracks' and 'trails' (eg. Flower 1945; Clark et al. 1972; Clark and Globensky 1975, 1976a).

As the content of this paper is decidedly taxonomic only a brief summary of the stratigraphic, sedimentologic and environmental setting of the Trenton Group is presented below. More detailed

reviews can be found in the referenced papers.

STRATIGRAPHIC, SEDIMENTOLOGIC AND ENVIRONMENTAL SETTING

In the St. Lawrence Lowland between Montreal in the southwest and Quebec City in the northeast, a distance of approximately 250 km (Fig. 1), the upper Middle Ordovician Trenton Group has been subdivided into many lithostratigraphic units several of which, because of their overall similarity, are candidates for synonymy (Harland and Pickerill 1982). Because such stratigraphic revision has not been attempted to date, the previously established units are retained herein. Figure 2 summarizes the Trenton Group stratigraphy and places it in context with underlying and overlying formations. More detailed lithostratigraphic and biostratigraphic summaries are available in Clark (1972), Clark and Globensky (1975, 1976a,b,c) Riva (1968, 1969, 1972). Figure 3 summarizes the essential characteristics and interpreted environment of deposi-




Fig. 1 - Simplified geological map of the St. Lawrence Lowland between Montreal and Quebec City. Numbered localities are: 1. Saint Constant, 2. Ile Bizard, 3. Montreal, 4. Cap Saint Martin, 5. Saint Vincent de Paul, 6. Terrebonne, 7. L'Epiphanie, 8. Saint Roch, 9. Saint Esprit, 10. Saint Jacques, 11. Highway 158/Rivière Ouareau, 12. Highway 158/Rivière Rouge, 13. Pont Corriveau, 14. Barrage savoie, 15. Pont Bournival, 16. Saint Louis de France, 17. Radnor des Forges, 18. Grondines, 19. Saint Casimir, 20. Pont Saint Alban, 21. Pont Rouge, 22. Neuville.




Fig. 2 - Stratigraphic profile of Middle Ordovician lithostratigraphic units between Montreal and Quebec City showing the relationship of the upper Middle Ordovician Trenton Group to underlying and overlying units. Modified after Harland and Pickerill (1982).

tion of each formation, generalized details of which are given below.

The basal Trenton Group formations comprise, from southwest to northeast, the Mile End, Ouareau, Fontaine, Saint Alban and Pont Rouge Formations. These are generally thin (2-6m) and consist of irregular beds of micritic and argillaceous calcisiltites which alternate with thin shales. The shales and most limestones are thoroughly bioturbated. Coquinas, lenses and layers of shell and skeletal hash and muddy skeletal intra-clastic layers also occur. Each of these formations is similar to the underlying Lowville and Leray Formations of the Black River Group and are interpreted as having originated in broadly similar conditions, that is in a low to moderate energy outer lagoon or sheltered near-shore environment which became narrower and more confined from Montreal to Quebec City (Harland and Pickerill 1982). The common occurrence of shell layers indicates a fairly close proximity to fully marine environments.

The overlying Deschambault Formation, present throughout the St. Lawrence Lowland, is generally 24-28 m thick, but thins to about 5 m at Montreal, and is characterized by 20cm - 1m thick, fine- to coarse-grained, tabular and trough cross-stratified skeletal limestones with numerous coquinas and layers of shell hash and only minor shales. These represent high energy deposits which accumulated in a complex

environment of migrating subtidal shoals, bars and large ripples (Harland and Pickerill 1982). Additionally, the formation contains more continuous parallel-laminated or low-angle cross-stratified beds which represent ephemeral beaches developed when shoals and bars were intermittently built up to sea level. Scouring appears to have been common and subtidal channels and runnels cut through the shoals and bars. From Montreal northeastwards, these skeletal limestones developed progressively closer to shore so that in the Quebec City area they accumulated along rocky shorelines and around islands (Harland and Pickerill 1984).

Overlying the Deschambault Formation is the Montreal Formation in the Montreal-Joliette area and the St. Casimir Member of the Neuville Formation from Trois Rivières to Quebec City (Fig. 2). The former is 110-140m thick and the latter 40-60m but generally thins northeastwards. Both consist of highly variable limestones interbedded with dark shales. The limestones are predominantly thinly and irregularly bedded, bioturbated argillaceous micrites but dark micritic rubbly beds or skeletal calcarenites and coquinas (generally 5-10cm but up to 25 cm thick) may also occur. Their high diversity marine faunas and predominantly fine-grained nature indicate a shallow subtidal, mainly low to moderate energy environment in which skeletal layers were produced by

storm reworking of indigenous shelly faunas or periodically derived, perhaps during storm-ebbs or other unusually strong offshore currents, from shoreward shoaling environments (Harland and Pickerill 1982).

The uppermost units of the Trenton Group are the Tetreauville Formation (150m thick) in the Montreal-Joliette area and the Grondines Member of the Neuville Formation (100-120m thick) from Trois Rivières to Quebec City (Fig. 2). Both consist of 3-15cm thick argillaceous and dark micritic limestones interbedded with dark shales. Infrequent thin shelly or skeletal beds, which are commonly graded, are also present. Nodular or irregularly bedded limestones are common and typically progressively increase in frequency through the sequence. Fully marine faunas dominated by brachiopods and trilobites are sparse but diverse and widely distributed whereas bioturbation is high. The sediments were deposited essentially from suspension in a relatively deep, low energy subtidal environment. The infrequent thin skeletal layers are interpreted as distal storm deposits (Harland and Pickerill 1982).

SYSTEMATIC PALEONTOLOGY

As in Häntzschel (1962, 1975), the trace fossils are described below in alphabetical order. In adopting this scheme, subjective and often overlapping ethological groupings, as utilized by several previous studies (e.g. Osgood 1970, Chamberlain 1971, Fürsich 1974a), are avoided and the procedure enables ease of reference to a single ichnogenus. Informally named trace fossils are listed separately after the formally named ichnogenera. Preservational terminology follows that adopted by Seilacher (1964), Webby (1969) and Häntzschel (1975). For brevity, the geographic and stratigraphic distribution and environmental occurrence of each ichnogenus is omitted from the descriptions. Instead reference should be made to Figures 4, 5 and 6 which summarize this information. Representative examples of the majority of collected specimens are now housed in the Department of Geology, University of New Brunswick, as Collection Sl61 - D. Fillion.

Ichnogenus *Arenicolites* Salter, 1857.

Type species - *Arenicola carbonaria*

| | FORMATION | LITHOLOGY | ENVIRONMENT | |
|---------------|--|----------------------|---|---|
| TRENTON GROUP | TETREAUVILLE NEUVILLE Grondines Member | 150 m 100-120 m | Interbedded argillaceous and micritic limestones and shales. Occasional thin skeletal layers. Rubby or nodular beds increase upwards. | Offshore, deep subtidal, low energy shelf subject to periodic storm influxes. |
| | MONTREAL NEUVILLE Saint Casimir Member | 110-140 m 40-60 m | Interbedded fine-coarse limestones with thin shales. Argillaceous micrites predominant but also thin skeletal limestones and coquinas. | Shallow subtidal, low to moderate energy environment. Periodic influxes of skeletal material from shoreward shoaling environments. |
| | DESCHAMBAULT | 5-28 m | Cross-stratified skeletal limestones coquinas and shell hash. Minor shales. | Shallow subtidal shoals, bars, large-scale migrating ripples and ephemeral beaches. High energy. |
| | MILE END OUAREAU FONTAINE SAINT ALBAN PONT ROUGE | 2-6 m | Micrites, argillaceous limestones and shale interbeds, with skeletal hash layers and lenses. | Low energy outer lagoon or sheltered nearshore environment with periodic influxes of storm and tidal overwashes from offshore shoals. |
| | BLACK RIVER GROUP | — | Mixed tidal flat, lagoonal and nearshore siliciclastics and carbonate muds. | |

Fig. 3 - Facies analysis and inferred depositional environments of the Trenton Group between Montreal and Quebec City (after Harland and Pickerill 1982).




Fig. 5 - Stratigraphic distribution and abundance of trace fossils in the Trenton Group between Montreal and Quebec City. The relative bar widths represent respectively from thinnest to thickest 1<10 specimens, 10<50 specimens, 50<100 specimens and >100 specimens. Questionable occurrences refer to uncertain identifications. Formational nomenclature is P.R. = Pont Rouge, S.A. = Saint Alban, F = Fontaine, O = Ouareau, M.E. = Mile End, D = Deschambault, N(SCM) = Saint Casimir Member (Neuville), M = Montreal, N(GM) = Grondines Member (Neuville), and T = Tetreauville Formations.

pressions compare with *A. sparsus* Salter 1857) and *A. statheri* Bather 1925, particularly with respect to the presence in some specimens of a distinct wall lining. Thus, more than a single ichnospecies may be present but the paucity of material and the nature of its preservation precludes more detailed analysis.

Ichnogenus *Caly craterion* Karaszewski, 1971

Type species - *Caly craterion samsonowiczii* Karaszewski 1971, by monotypy.

Diagnosis - Regular and smooth calyx-shaped depressions (on upper bedding surfaces) or knobs (on both upper and lower bedding surfaces) possessing up to three smaller depressions or knobs (after Karaszewski 1971).

Discussion - Karaszewski (1971) considered the structures to have been pro-

duced by annelids, the smaller depressions or knobs representing burrow outlets. The ichnogenus has, to date, only been recorded in nearshore (Karaszewski 1971), deltaic (Chaplin 1980) and intertidal environments (Fillion and Pickerill, in prep.) from, respectively, Jurassic, Carboniferous and Lower Ordovician strata. The asymmetry of the burrow and the smaller knobs or depressions differentiate *Caly craterion* from the closely related ichnogenera *Bergaueria* Prantl 1945 and *Mammilichnis* Chamberlain 1971.

?*Caly craterion* sp. (Figure 7e)

Material - 1 specimen.

Description - The trace occurs as a slightly elongated and ovoid mound preserved in convex epirelief on the upper

| ICHNOTAXA | OUTER LAGOON | BAR | SHALLOW SHELF | OFFSHORE SHELF |
|--|--------------|-----|---------------|----------------|
| <i>Arenicolites</i> sp. | | | | |
| ? <i>Calycraterion</i> sp. | | | | |
| <i>Chondrites</i> type A | █ | | | |
| <i>Chondrites</i> type B | █ | | | |
| <i>Chondrites</i> spp. | █ | | | |
| <i>Circulichnis montanus</i> | | | | |
| <i>Clematischnia</i> sp. | | | | |
| ? <i>Conostichus</i> sp. | | | | |
| <i>Cruziana problematica</i> | | | | |
| <i>Cruziana</i> sp. | | | | |
| cf. <i>Diplichnites</i> sp. | | | | |
| <i>Furculosus carpathicus</i> | | | | |
| <i>Helminthopsis hieroglyphica</i> | | | | |
| <i>Helminthopsis</i> sp. | | | | |
| <i>Oichnus paraboloides</i> | | | | |
| <i>Palaeophycus tubularis</i> | | | | |
| <i>Palaeophycus</i> sp. | | | | |
| ? <i>Plagiagmus</i> sp. | | | | |
| <i>Planolites beverleyensis</i> | | | | |
| <i>Planolites montanus</i> | | | | |
| <i>Planolites</i> sp. | | | | |
| ? <i>Rhizocorallium</i> cf. <i>R. irregulare</i> | | | | |
| ? <i>Rosselia</i> sp. | | | | |
| <i>Rusophycus</i> sp. | | | | |
| <i>Scalartituba missouriensis</i> | | | ? | |
| <i>Scolicia</i> sp. | | | | |
| <i>Skolithos linearis</i> | | | | |
| <i>Skolithos</i> sp. | | ? | | |
| <i>Teichichnus rectus</i> | | | | |
| <i>Teichichnus</i> sp. | ? | | | |
| <i>Trichichnus</i> sp. | | | | |
| <i>Trypanites weisei</i> | | | | |
| <i>Vermiforichnus clarkei</i> | | | | |
| <i>Zoophycos</i> sp. | | | | |
| Loop burrows | | | | |
| Oblique burrows | | | | |
| Pronged burrows | | | | |
| Bryozoan borings | | | | |

Fig. 6 - Environmental distribution of trace fossils in the Trenton Group between Montreal and Quebec City. Relative bar widths and questionable occurrences are as indicated in Figure 5.

surface of a 25mm thick calcilutite which immediately overlies a 10cm thick skeletal calcarenite. The 'mound' is filled with this bioclastic material, is 7mm in height and has a diameter of 9 - 11mm. A poorly preserved knob is present on the upper extremity of the structure.

Remarks - Since knowledge of the vertical aspect of the structure is unknown, even its ichnogenic assignment is uncertain. However, the asymmetry and the presence of the problematical knob are closely comparable to Ordovician examples of *Calycraterion* we have observed on Bell Island, eastern Newfoundland and we therefore tentatively diag-

nose the Trenton specimen as such. In the Newfoundland examples it is evident that *Calycraterion* can be repeated vertically and its preservation as convex epirelief structures is related to subsequent infilling of calyx-shaped burrow outlets. This style of preservation is similar to that of *Margaritichnus* Bandel 1973 but the Trenton specimen can be differentiated from this ichnogenus because *Margaritichnus* is usually remarkably symmetrical and flattened.

Ichnogenus *Chondrites*
von Sternberg, 1833

Type species - *Fucoides antiquus* Brongniart 1828, by subsequent designation of Miller (1889, p. 114).

Diagnosis - Regularly branching tunnel systems consisting of a small number of mastershafts open to the surface which ramify at depth to form a dendritic network (after Osgood 1970, Fürsich 1974a).

Discussion - Although the trace is easily recognizable at ichnogenic level, distinction between the more than 170 supposed ichnospecies (Chamberlain 1977), many of which are candidates for synonymy, is at present impossible without a thorough and detailed taxonomic study. Many of the ichnospecies proposed in the literature are based on variation in size, preservation and angle of branching and our review of this literature has indicated that the majority of these can probably be placed into synonymy with the 11 chondritid ichnospecies (from a total of 36 *Fucoides* species - see Osgood 1970) originally designated by Brongniart (1823, 1828). Chamberlain (1971, 1977) has in part resolved some of the ichnospecific nomenclatural problems but the taxonomic status is still unsatisfactory. Because of these difficulties we do not attempt to identify our material beyond the ichnogenic level but recognize various form types (cf. Osgood 1970).

Chondrites has been interpreted as a plant (Brongniart 1823), the burrow of a sipunculoid (Simpson 1957), or poly-

chaete annelid (Osgood 1975), or the burrow of unknown tentacle-bearing organisms (Taylor 1967), anthoptiloid seapens (Bradley 1981) and even small arthropods (Ekdale 1977). Simpson (1957) and Osgood (1970) have discussed the manner of production and preservation at some length. It is a marine but eurybathic form, having been reported from marginal marine (e.g. Hakes 1976, 1977; Fisher 1978), shallow subtidal (e.g. Hofmann 1979, Marintsch and Finks 1982), deep sea fan (e.g. Crimes et al. 1981, Hill 1981) and abyssal (e.g. Ekdale 1977, 1978; Larson 1982) environments. It ranges in age from Cambrian-Recent (Häntzschel 1975).

Finally, it is noteworthy that the nomenclatural type of *Chondrites* was designated by Miller (1889, p. 130) as *Fucoides antiquus* and subsequent designations (e.g. Bassler 1915, p. 217; Andrews 1955, p. 130) of *Fucoides targionii* are invalid by rule of priority. Additionally, *Fucoides lycopodioides* as listed in the Treatise by Häntzshel (1975) and adopted by some subsequent authors (e.g. Marintsch and Finks 1982) is in fact an error. This species is the nomenclatural type of *Caulerpites* von Sternberg, 1833.

Chondrites sp. type A
(Figures 8a, b)

Diagnosis - As for ichnogenus.

Material - 64 specimens.

Description - Dense, horizontal networks preserved in convex epirelief or hyporelief, composed of straight cylindrical burrows which exhibit regular symmetrical dichotomous branching at angles between 20 - 65°. Burrow diameter varies from 1.5 - 7mm but is constant within a single specimen. Branches are up to 150mm long and a maximum of 5 orders have been observed although there can be long segments without bifurcations. Burrow cross-sections are ellipsoidal and their contact with host material is sharp and well-defined. Burrow fill is variable but is typically finer than the host rock, is usually structureless but occasionally

exhibits concentric or meniscate structures (cf. Taylor 1967). The largest observed network covers an area of 120 x 150mm .

Remarks - The burrow systems frequently run in close proximity without interpenetration, thus suggesting phobotactic behaviour of the producer. The internal meniscate structures present in some of the specimens suggests, at least for these, an origin as stuffed burrows rather than open burrow systems (for a review on the genesis of *Chondrites* see Osgood 1970, Ekdale 1977). Although *Chondrites* type A has not been formally speciated it bears strong resemblance to *Buthotrephis succulens* Hall (1847, pl. 22, fig. 2a).

Chondrites sp. type B
(Figures 8a, c)

Diagnosis - As for ichnogenus.

Material - Several hundreds of specimens.

Description - Oblique networks preserved in endorelief with burrow diameters varying from 0.6 - 2mm and burrow cross-sections circular or elliptical. Burrow fill is of the same or coarser grain size as the host rock and is ubiquitously calcilutite or calcisiltite. The fill is typically of a different colour than the enclosing sediment and is derived from immediately overlying strata. The tubes, which are up to 25mm long, are often lined with micrite. Individual networks may possess 3 orders of branching and have been observed to penetrate at least 36mm of host sediment.

Remarks - These smaller networks of *Chondrites* sp. type B also exhibit good evidence of phobotactic behaviour. They are frequently observed between *Chondrites* type A networks but never penetrate them or, indeed, each other. Type B networks, although always initially oblique to stratification, may turn to become parallel at their distal extremities. The producer(s) of the trace is responsible for much of the bioturbation observed in the Trenton Group.

Chondrites spp.

(Figures 8a, d, e, f)

Specimens included here exhibit poor and variable preservation or are too irregular to permit assignment to types A and B as described above. Three variants can be recognized:-

(i) Elaborate, moderately dense and now flattened networks preserved at shale-calculutite interfaces of the Grondines Member of the Neuville Formation (Figs. 8a, d). Individual networks are composed of sinuous tunnels which vary in diameter along their length from 2 - 8.5mm. Up to 5 orders of branching are present and the branching is very irregular. Branching angles are equally irregular varying from 20 - 90°. Burrow fill is of the same grain size as the enclosing material (micrite), is typically structureless but occasionally exhibits a crudely developed meniscate appearance. (ii) A loose horizontal network preserved at a calcisiltite - calcarenite bed in the Deschambault Formation (Figs. 8a, f). Tunnels vary in diameter from 2 - 4mm but the diameter of a single tunnel remains constant along its length. The branching is curved, asymmetrically dichotomous but never crossing, occurs at angles of 35 - 60° and up to 5 orders can be observed. Burrow fill is micrite. In many respects, this network is very similar to *C. furcatus* as figured by Marintsch and Finks (1982, pl. 2, fig. 4) and *C. antiquatus* as figured by Hofmann (1979, pl. 15, fig. F). (iii) A widely spaced regularly branching network of asymmetrically dichotomous tunnels (Figs. 8a, e). Individual tunnels are straight and retain a constant diameter of 1 - 3 mm. Branching angles vary from 30 - 40° and up to 2 orders were observed. Burrow fill is calculutite but a dark shale lining can be observed in some specimens. This variety resembles *C. targonii sensu* Marintsch and Finks (1982).

Ichnogenus *Circulichnis* Vialov, 1971

Type species - *Circulichnis montanus*, Vialov, 1971, by original designation.

Diagnosis - "Annular trace, almost cir-

cular (or oval) formed by some cylindrical object" (Vialov 1971, p. 91, trans. litt.).

Discussion - The original diagnosis by Vialov (1971) is somewhat confusing in that it is clear in the remainder of his text that the circular (or oval) trace is in fact a burrow of 0.7 - 1.5mm in diameter. Since the original designation it has only been formally recorded from the Tremadoc of Nova Scotia (Pickerill and Keppie 1981). Similar structures have been reported and, or, figured but not described by various authors (Ulrich 1904, pl. 22, fig. 3; Häntzschel 1975, fig. 44, 2a; Książkiewicz 1977, fig. 32; Nicholson 1978, pl. 3, fig. 2; Crimes et al. 1981, p. 975). Preferably such structures should be included within the ichnogenus *Circulichnis* with which they best conform. Accordingly, *Circulichnis* would range from Tremadoc-Palaeocene age. Although its producer and environmental range are uncertain Pickerill and Keppie (1981) suggested a worm-like originator and a preference for deeper-water regimes. Crimes (pers. comm., 1983) has also recorded it from deep-water turbidites of Carboniferous age in Spain and England.

Circulichnis montanus Vialov, 1971

(Figure 7g)

Diagnosis - As for ichnogenus.

Material - 1 specimen.

Description - Slightly elliptical trace preserved in convex epirelief on the upper surface of a 7cm calcarenite. The trace measures 115 x 90mm and the burrow forming it is 4mm in diameter and consistent in width throughout the ellipse. Evidence of burrow collapse is present locally (Fig. 7g). Burrow fill is identical to the host rock.

Remarks - Since Vialov (1971) only gave size measurements of his holotype, the full variation in ellipse size and burrow diameter of the original material is unknown. In view of the incomplete description and the unavailability of type material for restudy, the Trenton specimen can best be regarded as conspeci-

fic. Its occurrence in the Trenton Group, although rare, suggests that it is possibly more a eurybathic form.

Ichnogenus *Clematischnia* Wilson, 1948

Type species - *Buthotrephis succulens* Hall, 1847, by original designation of Wilson (1948, p. 10).

Diagnosis - Irregularly branched or unbranched cylindrical burrows with prominent external ornament of transverse ribs which completely or partially cross the burrow; in the latter case the burrow assumes a distinctive mammilated appearance (modified after Wilson 1948).

Discussion - Wilson (1948) designated *Buthotrephis succulens* Hall as her type ichnospecies of *Clematischnia* but it is apparent that Hall's (1847) two figured specimens are morphologically separate; his figure 2a, plate XXI is more akin to a chondritid (*Chondrites* type A as described herein) and his figure 2b, plate XXI falls within *Clematischnia* as defined by Wilson (1948). Furthermore, even Wilson's (1948) material is taxonomically variable; her plesiotype G.S.C. 9283 (plate 1, fig. 4) is likewise more akin to *Palaeophycus* sp. or *Chondrites* sp. whereas her plesiotype G.S.C. 13442 (plate 1, fig. 5) falls within *Clematischnia* and is very similar to Hall's (1847) figure 2b. Clearly a detailed systematic review of the ichnogenus is warranted but until this is undertaken we regard it as a useful form to describe these distinctive mammilated burrows. Because of the rarity of actual recordings of *Clematischnia* (for example the ichnogenus was only listed as an unrecognized form in Häntzschel (1975) and briefly discussed by Osgood (1970) in his re-description of *Buthotrephis succulens*) it is difficult to comment on its pro-

ducer and stratigraphic range. Its similarity to chondritids and *Palaeophycus* suggest, however, that it is the feeding and dwelling burrow of an unknown annelid. Similarly, although to date it has only been reported from Ordovician strata, its stratigraphic range is likely to be considerably more extensive.

Clematischnia sp.
(Figure 7i)

Material - 12 specimens.

Description - Unbranched cylindrical burrows up to 7mm in diameter and 10cm in length preserved on upper surfaces and all parallel to stratification. In all of the observed specimens the burrows occur in groups of four to seven, apparently radiating from some common centre which is not preserved. Burrow surfaces are mammilated, this being produced by an alternating series of transversely oriented interlocking pear or cylindrical shaped protruberances, each separated by a narrow depression. Burrow fill (siltstone) is coarser than the surrounding matrix and is structureless.


Remarks - Congeneric material from the Trenton Group immediately around Quebec City was previously identified by Pickerill and Forbes (1979) as cf. *Biformites*. However, both their material and that described herein lacks the initial short and longitudinally furrowed section more typical of *Biformites* and the material is therefore best included within *Clematischnia*. In view of its unresolved taxonomic status it is herein only identified at ichnogenetic level.

Ichnogenus *Conostichus* Lesquereux, 1876

Type species - *Conostichus ornatus* Lesquereux, 1876, by original monotypy.

Opposite page

Fig. 7 - a. *Cruziana* sp. on lower bedding surface in convex hyporelief, Grondines Member, locality 22. b. *Cruziana problematica* on lower bedding surface in convex hyporelief, Deschambault Formation, locality 12. c. *Scolicia* sp. and *Diplichnites* sp. (arrowed) preserved on upper bedding surface of thin chert, Grondines Member, locality 18 (see text for details). d. *Arenicolites* sp. tubes (arrowed) on upper bedding surface, Saint Casimir Member, locality 16. e. ?*Calycraterion* sp. on upper bedding surface in convex epirelief with arrow indicating possible location of burrow outlet, Deschambault Formation, locality 21. Coin diameter 2 cm. f. ?*Conostichus* sp. actually preserved on lower bedding surface in convex hyporelief but here reversed to exhibit faint longitudinal fluting, Grondines Member, locality 18. g. *Circulichnis montanus* on upper bedding surface in convex epirelief, arrow indicating burrow collapse, Deschambault Formation, locality 21. h. *Rusophycus* sp. on lower bedding surface in convex hyporelief, Grondines Member, locality 22. i. *Clematischnia* sp. on upper bedding surface in convex epirelief, Pont Rouge Formation, locality 21.






Fig. 8 - a. Variation in chondritids in the Trenton Group. 1 = *Chondrites* sp. type A, 2 = *Chondrites* sp. type B, 3 = *Chondrites* sp. dense and elaborate networks, 4 = *Chondrites* sp. loose horizontal networks, and 5 = *Chondrites* sp. widely spaced horizontal regularly branching networks (see text for details). All redrawn from actual specimens and all x 0.5. b. *Chondrites* sp. type A on upper bedding surface, Pont Rouge Formation, locality 21. c. *Chondrites* sp. type B in vertical cross-section, Tetreauville Formation, locality 8. d. Dense horizontal network of *Chondrites* sp. on lower bedding surface, Grondines Member, locality 18 (cf. Fig. 8a [3]). e. Loose horizontal network of *Chondrites* sp. on upper bedding surface, Deschambault Formation, locality 21 (cf. Fig. 8a [5]). f. *Chondrites* sp. on upper bedding surface, Deschambault Formation, locality 16 (cf. Fig. 8a [4]).

Diagnosis - Variable but mostly small conical to subconical structures preserved in full relief. Bodies commonly fluted by transverse constrictions and longitudinal furrows and ridges originating from the apical end. Concentric laminae may be present internally (after Häntzschel 1975, Hakes 1976).

Discussion - Following its original interpretation as an alga (Lesquereux 1876) the structure has been interpreted as a sponge, a scyphomedusoid, a burrow of a filter-feeding organism, a funnel-shaped opening of *Arenicola*-like burrows and a trace produced by upward migration of actinian-like organisms (see Pfefferkorn 1971, Chamberlain 1971, Barthel and Barth 1972, Hakes 1976). Its origin still remains enigmatic and, indeed, its producers may be polyphyletic. Many of its numerous species are probably synonymous and it is a prime candidate for serious taxonomic revision (see Branson 1960). It ranges in age from Arenig (pers. observ. - Wabana Group of eastern Newfoundland) to ?Lower Cretaceous (Imlay 1961) and, although exceptions do occur (e.g. Chamberlain 1971), is typically associated with near-shore environments subject to rapid sedimentation.

?*Conostichus* sp.
(Figure 7f)

Material - 1 specimen.

Description - A truncated conical structure 15mm in diameter and 13mm in height preserved in convex hyporelief below a 25mm-thick bioclastic calcilutite. Poorly preserved longitudinal ribs (ca. 6-7) are present.

Remarks - The specimen, although not well enough preserved for definite identification, may be distinguished from the allied ichnogenus *Conichnus* Myanil, 1966 by the presence of longitudinal fluting. Additionally, *Bergaueria* Prantl, 1945 is more flattened and less conical. A more important difficulty is demonstrating that the structure is in fact organic and not inorganic in origin. The former, however, is suggested by the

regularity of the structure, its association on the same bedding surface with abundant trace fossils, particularly *Chondrites* sp. type B and *Planolites beverleyensis*, and a complete absence on that bedding surface of additional physically produced inorganic structures.

Ichnogenus *Cruziana* d'Orbigny, 1842

Type species - *Cruziana rugosa* d'Orbigny 1842, by subsequent designation of Miller (1889, p. 115).

Diagnosis - Elongate band-like furrows covered by herringbone-shaped ridges, with or without two outer sooth or finely longitudinally striated zones outside the V-markings, occasionally with lateral ridges and, or, wisp-like markings when preserved on bedding soles (after Häntzschel 1975).

Discussion - Although Seilacher (1970) united under *Cruziana* both long furrows and short coffee bean-like excavations (= *Rusophycus*) most subsequent authors have, as we do, preferred to retain the two as distinctive ichnogenera (see Crimes et al. 1977). Furthermore, we herein accept the reasoning by Bromley and Asgaard (1979) to include *Isopodichnus* Bornemann, 1899 within *Cruziana* as the only essential differences between the two previously separated ichnogenera are in accessory features (*sensu* Fürsich 1974b) which should be employed only for classification at the ichnospecific level. It is, nevertheless, clear that Bromley and Asgaard (1979) 'lumped' heterogeneous material into a single ichnospecies (*C. problematica*) and revision of their and other material is still necessary.

Undoubtedly, the majority of *Cruziana* furrows were produced by trilobites (Seilacher 1970), although the small Triassic forms described by Bromley and Asgaard (1970) may have been produced by trilobite-like arthropods or notostracan branchiopods and the large Middle Ordovician and Triassic forms described by, respectively, Fisher (1978) and Shone (1978, 1979) may be the result of aglaspids or even vertebrates. Many ichno-

species of *Cruziana* now exist and it is perhaps the most useful of all trace fossils in stratigraphic studies (Seilacher 1970, Crimes 1975). The ichnogenus ranges in age from Cambrian - Triassic and is a typically shallow-water marine form constituting the type ichnogenus of Seilacher's (1967) *Cruziana* ichnofacies, though it has also allegedly been observed in freshwater deposits (e.g. Helwig 1972).

Cruziana problematica (Schindewolf, 1921)
(Figure 7b)

Diagnosis - Straight or curved, often self-crossing, *Cruziana* up to 5mm wide possessing faint, unevenly spaced transverse striations which may reach the margins of the trace in low relief specimens or terminate before reaching the margins in high relief specimens (adapted from Schindewolf 1928, Häntzschel 1975, Trewin 1976, Bromley and Asgaard 1979).

Material - 2 specimens.

Description - Two poorly preserved specimens, one in convex hyporelief at the base of a 30mm thick calcisiltite, the second in concave epirelief on an 80mm thick calcilutite. The former is a short 8mm long straight band-like trace, 4mm wide with lobes 1.5mm deep and the latter a 120mm long slightly sinuous trace, 2mm in width, composed of two furrows each 0.7mm wide separated by a median ridge. Both specimens possess transverse but poorly preserved and faint scratch markings irregularly developed along their exposed length.

Remarks - Although poorly preserved the material lacks the 'Isopodichnus' - type bilobed expanded terminations

(sensu Trewin 1976) typical of *C. stromnessa* and the well-defined scratches, often arranged in a V-pattern, of *C. eutendorfensa*. Conspecific material was described by Pickerill and Forbes (1979) from the Trenton Group northeast of Quebec City.

Cruziana sp.
(Figure 7a)

Material - 8 specimens.

Description - The specimens consist of variably preserved, straight or slightly sinuous bilobate furrows preserved in convex hyporelief. Mean width is 20mm and length is variable, up to a maximum observed of 100mm. Endopodal lobes are poorly to moderately well-preserved giving the trace a depth of 1-8mm. V-markings, when preserved, are moderately deep, closely spaced and V at 160 - 180°. Genal spine markings are absent.






Remarks - Though not well enough preserved for ichnospecific identification the furrows can be clearly diagnosed as *Cruziana*. They are reminiscent of specimens described as *Cruziana* sp. by Alpert (1976, pl. 1, fig. 4) from the Cambrian of California and also resemble *C. pudica* described from Middle Ordovician to Middle Silurian strata of the United States, southern Jordan and Wales by, respectively, Osgood (1970), Seilacher (1970) and Pickerill (1977). Congeneric material has also been figured by Pickerill and Forbes (1979) from the Trenton Group northeast of Quebec City.


Ichnogenus *Diplichnites* Dawson, 1873

Type species - *Diplichnites aenigma* Dawson, 1873, by original monotypy.

Opposite page

Fig. 9 - a. *Furculosus carpathicus* on lower bedding surface in convex hyporelief, Montreal Formation, locality 11. b. Unbranched *Palaeophycus tubularis* on upper bedding surface in convex epirelief, Deschambault Formation, locality 21. Coin diameter 2 cm. c. ?*Plagiogmus* sp. on upper bedding surface in convex epirelief, Deschambault Formation, locality 11. d. *Helminthopsis* sp. on upper bedding surface in convex epirelief, Deschambault Formation, locality 21. Coin diameter 2 cm. e. *Helminthopsis hieroglyphica* on upper bedding surface in convex epirelief, Pont Rouge Formation, locality 21. Pen length 10 cm. f. Lined *Planolites* sp. on upper bedding surface in concave epirelief, Grondines Member, locality 18. Coin diameter 2 cm. g. *Planolites beverleyensis* on upper bedding surface in convex epirelief, Pont Rouge Formation, locality 21. h. *Planolites montanus* on upper bedding surface, Grondines Member, locality 18. i. *Helminthopsis* sp. (arrowed) and *Palaeophycus* sp. (to left of coin) on upper bedding surface in convex epirelief, Deschambault Formation, locality 21. Coin diameter 2 cm.





Diagnosis - Morphologically simple trails up to 36cm wide and consisting of two parallel series of tracks (each up to 9cm wide); individual tracks elongate, closely and regularly spaced up to one per centimeter and roughly normal to trace axis (emend. Briggs et al. 1979).

Discussion - Although Dawson (1873, p. 7) originally interpreted *Diplichnites* as the result of 'a large crustacean or gigantic annelid or myriapod' subsequent workers, following Seilacher (1955), have frequently utilized the ichnogenus to describe morphologically similar but smaller scaled trilobite-produced traces (e.g. Crimes 1970, Osgood and Drennen 1975). Although Briggs et al. (1979) questioned the trilobite origins of originally described specimens and suggested a more restricted usage to exclude trilobite trails, suggesting that those specimens of *Diplichnites* previously interpreted as trilobite trails to be included in one of the junior synonyms listed by Osgood (1970), the taxonomic confusion has still not been resolved. Because of the limited material and its poor preservation and the unresolved taxonomic difficulties, we herein retain *Diplichnites* even though we assume it to have been produced by trilobites. The ichnogenus has been reported from Cambrian-Triassic strata and though typically a shallow marine form (e.g. Crimes 1977) has also been recorded from deep-water sediments (e.g. Pickerill 1980, 1981).

cf. *Diplichnites* sp.
(Figure 7c)

Material - 2 specimens collected and 5 recorded in the field.

Description - Paired(?) elongated dot-like impressions preserved in concave epirelief on the upper surface of a chert bed. Each impression is 2.5mm long,

1mm wide and 1mm deep and set width (sensu Osgood 1970, p. 354) is 7-11mm. Up to 13 sets have been observed in a single track, giving a maximum length of 45mm.

Remarks - The poor preservation precludes positive identification even at ichnogenus level, although of all published 'trackways' they compare most favourably to *Diplichnites*.

Ichnogenus *Furculosus*
Roniewicz and Pieńkowski, 1977

Type species - *Furculosus carpathicus* Roniewicz and Pieńkowski, 1977, by original designation.

Diagnosis - Cylindrical burrows forming tight fork-line loops with endings parallel or diverging.

Discussion - Roniewicz and Pieńkowski (1977) hypothesized that the loops of *F. carpathicus* were formed by a deposit-feeding organism passing through an interface boundary. Although to date no other occurrence of the ichnogenus has formally been made, similar forms are known in the external Carpathians (Książkiewicz, pers. comm. in Roniewicz and Pieńkowski 1977). The occurrence of *Furculosus* in the Trenton Group extends its known stratigraphic range from Middle Ordovician - Oligocene and its environmental range from shallow to deep-water regimes. *Gordia arcuata* Książkiewicz, 1977 is a morphologically similar form but can be distinguished from *Furculosus* by its more arcuate shape and smaller burrow diameter.

Furculosus carpathicus
Roniewicz and Pieńkowski, 1977
(Figure 9a)

Diagnosis - As for ichnogenus.

Material - 1 specimen.

Description - Tight-looped smooth and

Opposite page

Fig. 10 - a. *Trichichnus* sp. (arrowed) in vertical section, Montreal Formation, locality 14. b. *Zoophycos* sp. on upper bedding surface in convex epirelief, Grondines Member, locality 18. Note the peripheral tube (arrowed). c. *Teichichnus rectus* in transverse vertical section with the longitudinal burrow arrowed, Grondines Member, locality 21. d. Thickly lined *Skolithos* sp. burrow (arrowed) in transverse vertical section, Pont Rouge Formation, locality 21. e. *Scalarituba missouriensis* on upper bedding surface in convex epirelief, Ouareau Formation, locality 11. f. g. ?*Rosselia* sp., f. on upper bedding plane surface, and g. in transverse vertical section, Saint Casimir Member, locality 16. h. *Teichichnus* sp. on upper bedding surface, Grondines Member, locality 18.

cylindrical burrow preserved in convex hyporelief on the base of a parallel laminated calcarenite. Burrow diameter is 3.5 - 4mm and the loop is 35mm long and 11mm wide. The burrow terminates as a result of both distal extremities turning up into the overlying calcarenite. Burrow fill is calcarenite.

Remarks - Although Roniewicz and Pieńkowski (1977) discussed the 3-dimensional aspect of *F. carpathicus* they omitted to include it in their diagnosis, according to which, knowledge of the bedding plane expression of the burrow is sufficient for positive identification. The tight fork-like loop is therefore the most characteristic feature of the species and the Trenton material is therefore regarded as conspecific.

Ichnogenus *Helminthopsis* Heer, 1877

Type species - *Helminthopsis abeli*, by subsequent designation of Książkiewicz (1977, p. 116).

Diagnosis - "Unbranched, very long, very much snake-like winding tubes or cylinders of very different sizes. *Helminthopsis magna* was undoubtedly tubular but this is uncertain for the two other species The outer surface is smooth. It is similar to *Helminthoida* Schafh. in which, however, the meanders are closer and the string-like or cylindrical body touches itself" (Heer 1877, p. 116 - trans. litt.).

Discussion - Heer (1877) originally established three species of *Helminthopsis* but did not designate a type. Subsequently, Ulrich (1904, p. 144) designated *H. magna* a somewhat unfortunate choice as this trace is a bilobate structure (see Heer 1877, pl. XLVII, figs. 1,2) resembling *Taphrhelminthopsis* Sacco, 1888 or *Scolicia* de Quatrefages, 1849. In view of this problem Książkiewicz (1977) proposed to utilize a specimen originally figured by Abel (1935, p. 290, fig. 261B) and subsequently illustrated by Häntzschel (1962, p. W197, fig. 4a; 1975, p. W71, fig. 2b) as the type ichnospecies under the name *Helminthopsis abeli*. Compared to *H. magna* this latter ichnospecies conforms best to Heer's (1877) original diagnosis

and is retained herein as the type.

It must be noted that although most authors have utilized the ichnogenus for winding or loosely meandering, unbranched, generally unornamented 'burrows', the original diagnosis did not specify that the traces had necessarily to be of burrow origin. Thus, it seems logical that morphologically and geometrically similar 'trails' (e.g. Hall 1852, pl. 11, fig. 2b; Dörjes 1978, fig. 4 and Swinbanks and Murray 1981, fig. 10A) should also be included within the ichnogenus.

Many authors, too numerous to mention here, have reported the ichnogenus in strata of ?Late Precambrian (Webby 1970, fig. 17, *H. hieroglyphica*) to Oligocene (Książkiewicz 1977) or even Recent (Dörjes 1978, Swinbanks and Murray 1981) age. It is a eurybathic form, though more frequently reported from deep-water flysch successions (Pickerill 1981) and is generally regarded as having been produced by polychaete and possibly priapulid annelids.

Helminthopsis hieroglyphica
Heer in Maillard, 1887
(Figure 9c)

Diagnosis - *Helminthopsis* in which the windings are irregular, low and the course of the trace is often alternately winding and straight (after Książkiewicz 1977).

Material - 7 specimens.

Description - The traces are moderately well-preserved in convex epirelief on the upper surface of a massively bedded crinoidal calcarenite. They consist of burrows, up to 2cm in diameter and 50cm in length which are characterized by long straight segments up to 30cm long, but usually smaller, with a low turn between individual segments. Burrow diameter is constant along the length, burrow fill is identical to the enclosing calcarenite material and the burrows are smooth and unornamented.

Remarks - As noted by Książkiewicz (1977, p. 119), this ichnospecies is not well defined and the material originally figured in Maillard (1887, pl. 2, fig. 4;

pl. 1, fig. 2) resembles both a *Helminthoida* (pl. 1, fig. 2) and a loose irregularly winding trace (pl. 2, fig. 4) which Książkiewicz argued be adopted as the nomenclatural type of the species. To maintain stability this approach is adopted herein.

***Helminthopsis* sp.**
(Figures 9d, i)

Material - 5 specimens and numerous occurrences recorded in the field.

Description - Variably preserved burrows preserved in convex and concave epirelief on upper surfaces of thinly bedded calcilutites or bioclastic calcisiltites. Burrows are smooth, slightly sinuous or meandering but meanders are always incomplete. Burrow fill is coarser or of the same grain size as underlying sediments. They are up to 150cm in length and range in diameter from 1.8 - 30mm (mean 7mm), the diameter remaining constant in a single specimen. Collapse in some specimens has resulted in their preservation as two epichnial ridges separated by a narrow furrow.

Remarks - The most workable scheme for ichnospecific identification of *Helminthopsis* is that proposed by Książkiewicz (1977), who utilized the nature and type of winding, the burrow diameter and the presence or absence of surface ornamentation. Because of their generally poor and incomplete preservation, particularly regarding the meander patterns, the Trenton specimens can therefore best only be identified at ichnogeneric level. Although *Helminthopsis* is generally regarded to be a deposit-feeding burrow (Książkiewicz 1977) the presence of collapsed tunnels in some of the Trenton specimens suggests, at least for these specimens, that they remained open after the passage of the producing organism(s).

Ichnogenus *Oichnus* Bromley, 1981

Type species - *Oichnus simplex* Bromley, 1981, by original designation.

Diagnosis - Circular to subcircular holes

of biogenic origin bored into hard substrates. The hole may pass right through the substrate as a penetration, where the substrate is a thin shell; or end within the substrate as a shallow to deep depression or short, subcylindrical pit (Bromley 1981).

Remarks - The borings are considerably shorter than *Trypanites Mägdefrau*, 1932, range in age from possibly early Cambrian to Recent, and certainly Tertiary and younger examples are produced by predatory snails, as more fully discussed in Bromley (1981).

***Oichnus paraboloides* Bromley, 1981**
(Figure 11a)

Diagnosis - *Oichnus* having a spherical paraboloid form, truncated in those cases where the boring penetrates right through the substrate. Where it does not so penetrate, the paraboloid may be deformed by a slightly raised central boss (Bromley 1981).

Material - 1 specimen.

Description - A round perforation within the pygidium of the trilobite *Isotelus gigas*, 1.2mm in diameter at the initial point of penetration of the dorsal surface narrowing to 1mm at its ventral exit. Walls of the truncated paraboloid are smooth.

Remarks - Although we observed no evidence of etching patterns on the boring it is identical to and diagnosed as *O. paraboloides* Bromley, 1981, as *O. simplex* Bromley, 1981 possesses a cylindrical to subcylindrical form. The producer of the Trenton specimen remains enigmatic.

Ichnogenus *Palaeophycus* Hall, 1847

Type species - *Palaeophycus tubularis* Hall, 1847, by subsequent designation of Miller (1889, p. 130).

Diagnosis (emended) - Branched or unbranched, straight to slightly curved to slightly undulose or flexuous, smooth or ornamented, lined essentially cylin-

drical, predominantly horizontal burrows of variable diameter; infillings typically structureless, of same lithology as host rock (after Pemberton and Frey 1982).

Discussion - There is still much confusion regarding the original designation of the type of the ichnogenus *Palaeophycus*. In their thorough and detailed review of *Palaeophycus* and *Planolites* Pemberton and Frey (1982) followed Häntzschel (1975) and suggested the type was *P. tubularis* Hall, 1847, subsequently designated by Bassler (1915, p. 939). At an earlier date, however, Miller (1889, p. 130), in a frequently overlooked and comprehensive text, had also designated *P. tubularis* Hall, 1847. Unfortunately, both Bassler (1915) and Miller (1889) did not design any type material, and indeed, the first to designate such material was Andrews (1955, p. 202). In the absence of any formal and universal code, however, at present we regard the subsequent designation of Miller (1889) and not Bassler (1915) as the most acceptable.

Minor emendation of the diagnosis of *Palaeophycus* by Pemberton and Frey (1982) is considered necessary to exclude systematic and non-systematic winding and, or, meandering burrows (e.g. *Gordia* Emmons, 1844, *Helminthopsis* Heer, 1877, *Cochlichnus* Hitchcock, 1858). Otherwise, the review of the taxonomy of *Palaeophycus* by Pemberton and Frey (1982), following several decades of inconsistent usage and confusion, is considered the most workable nomenclatural system and is adopted herein, even if problems may arise in the case of concealed bed-junction preservation (Hallam 1975), interface-crossing burrows (Hofmann 1979) or faintly meniscate burrows (Clifton and Thompson 1978).

Palaeophycus is a eurybathic form ranging in age from Proterozoic to Recent (Häntzschel 1975) and is usually attributed to represent the burrows of predaceous or suspension-feeding organisms, commonly but not universally polychaete annelids.

Palaeophycus tubularis Hall, 1847
(Figure 9b)

Diagnosis - Smooth, unornamented *Palaeophycus* of variable diameter, thinly but distinctly lined (Pemberton and Frey 1982).

Material - Several hundreds of specimens.

Description - Straight to slightly sinuous, generally smooth, horizontal to slightly inclined, frequently collapsed cylindrical burrows, 1-24mm in diameter and up to 57cm in length. Burrow diameter is constant along the length of a single burrow. True branching is rare, but when observed is irregular and occurs at 30-70°; cross-cutting is common. Burrow walls are thinly lined, burrow fill is identical to enclosing material and rare backfill structures are present. The burrows are preserved in convex hyporelief and convex and concave epirelief typically on thin beds of fine calcarenite, calcisiltite or calcilutite.

Remarks - Some of Hall's (1847) original material which he described as *P. rugosus* and *P. simplex*, but now both included in *P. tubularis*, was collected from the Trenton Group of New York and therefore the ichnospecies is apparently widespread and abundant throughout the Group. Although we have chosen to adopt the classification scheme of Pemberton and Frey (1982), some of the Trenton *P. tubularis* specimens illustrate some of the difficulties of its universal application. In particular, several specimens were observed to pass from calcilutites and calcisiltites into and even through patchily distributed lenses of calcarenite, though still retaining a calcisiltite or calcilutite fill throughout. Had only the portion of the burrows within the calcarenite been observed then they would have had to be referred to as *Planolites* which, of course, would necessitate, in such cases, erroneous ethological interpretations. Nevertheless, in most cases it was possible to observe the transition and such 'compound' specimens were named according to their dominant component (cf. Bromley and Frey 1974, Frey et al. 1978).




Fig. 11 - a. *Oichnus paraboloides* (upper arrow) and *Vermiforichnus clarkei* (lower arrow) in the pygidium of the isotelenid *Isotelus gigas*, Grondines Member, locality 18. b. Micrite filled borings of *Trypanites weisei* in a hemispherical colony of the bryozoan *Prasopora* sp., Pont Rouge Formation, locality 18. c. Longitudinal section through the bryozoan *Prasopora* sp. illustrating essentially vertical *Trypanites weisei* borings, Grondines Member, locality 18. d. Numerous small borings of *Vermiforichnus clarkei* in the cephalon of the isotelenid *Isotelus gigas*, Grondines Member, locality 18. e. f. Two views of reticulate bryozoan borings on a pygidium of the isotelenid *Isotelus gigas*, with f. an enlargement of the central area exhibited in e. Grondines Member, locality 18.

Palaeophycus sp.
(Figure 9i)

Material - 67 specimens.

Description - Ellipsoidal, thinly lined burrows, 3-7mm in diameter and observed mainly as full relief structures in transverse section. Burrow fill is identical to host rock and is apparently structureless. The few examples observed in semirelief are poorly preserved and incomplete, but essentially straight to slightly sinuous, horizontal to gently inclined, smooth, unbranched and rarely branched at up to 25°.

Remarks - Since the specimens are observed essentially in transverse section and their complete 3-dimensional character is unknown they are only identified at ichnogenic level. Semirelief specimens are apparently smooth and therefore possibly *P. tubularis*. This smoothing, however, could be a result of weathering as the specimens are badly preserved and weathered throughout. *P. heberti* (de Saporta 1872-73) can be dismissed because of the presence of only thin linings, but whether or not the material also represents one or several of the ichnospecies *P. striatus* Hall, 1852, *P. sulcatus* (Miller and Dyer 1878) or *P. alternatus* Pemberton and Frey, 1982, whose differentiation is based on external sculpture, is unknown.

Ichnogenus *Plagiogmus* Roedel, 1929

Type species - *Plagiogmus arcuatus* Roedel, 1929, by subsequent designation of Häntzschel (1962, p. W210).

Diagnosis - Horizontal endichnial burrow, straight or slightly curved 10 - 28mm wide, possessing sausage-shaped or fusiform transverse ventral ridges (as viewed on upper surfaces) and corresponding dorsal furrows (as viewed on lower surfaces), the spacing of which varies considerably. Narrow smooth zones separate the ends of the ridges from the margins of the trace (after Häntzschel 1975, Jaeger and Martinsson 1980).

Discussion - *Plagiogmus* was originally described from Sweden by Nathorst

(1897) but not formally erected until 1929 by Roedel, who defined two species, *P. simplex* and *P. arcuatus*, the latter subsequently being designated as the type by Häntzschel (1962). Roedel (1929) believed the structures were trails produced by polychaetes, but later Glaessner (1969) and more recently Jaeger and Martinsson (1980), who restudied the type material, demonstrated their true burrow nature. Although the nature of the producing organism is enigmatic, molluscs (Glaessner 1969, Crimes et al. 1977) and segmented annelids (Roedel 1929, Whitaker 1979 - described as *Steinsfjordichnus brutoni*) have been considered as likely producers. Jaeger and Martinsson (1980) have demonstrated that irrespective of the geological affinities of the producer, the movement was controlled and locomotory and not peristaltic in any sense.

The ichnogenus commonly occurs in strata of early Cambrian age (e.g. Banks 1970, Alpert 1977, Crimes et al. 1977) though Häntzschel (1975, p. W95) reports it from the Upper Precambrian of Russia. If, as we suspect, specimens of *Steinsfjordichnus brutoni* described by Whitaker (1979, p. 87) are congeneric, its stratigraphic range must be extended to the Lower Devonian. It has previously been recorded from shallow marine (Banks 1970) to possibly fluvial and lacustrine environments (Whitaker 1979).

?*Plagiogmus* sp.
(Figure 9c)

Material - 1 specimen recorded in the field.

Description - Preserved in convex epirelief on the upper surface of a 15cm thick massive bioclastic calcilitite as a series of 13 parallel and equally spaced (4mm) ridges, each 1mm high, 2mm wide and 10.5mm long, oriented transversely to the trace axis. The areas between the ridges are smoother, flatter and slightly wider. Burrow walls are not preserved.

Remarks - Poor preservation and lack of knowledge of the 3-dimensional struc-

ture precludes positive identification even at ichnogeneric level but the specimen does resemble *Plagiogmus* to which it is tentatively referred. The specimen is very similar, both in scale and morphology, to *Steinsfjordichnus brutoni* Whitaker, 1979, which as noted previously, is also best included in *Plagiogmus*.

Ichnogenus *Planolites* Nicholson, 1873

Type species - *Planolites vulgaris* Nicholson and Hinde, 1875, by subsequent designation of Miller (1889, p. 520).

Diagnosis (emended) - Unlined, rarely lined, rarely branched, straight to tortuous, smooth to irregularly walled or annulated burrows, circular to elliptical in cross-section, of variable dimensions and configurations; infillings essentially structureless, differing in lithology from host rock (after Pemberton and Frey 1982).

Discussion - Although Alpert (1975, p. 512) was the first to designate a type specimen of *Planolites*, Miller (1889, p. 520) and later Bassler (1915, p. 982) were the first authors to designate the (same) type species. Like *Palaeophycus*, *Planolites* has been a topic of considerable confusion since its original establishment by Nicholson (1873). Detailed summaries have been presented by Alpert (1975), Benton and Trewin (1978) and more recently Pemberton and Frey (1982). As with *Palaeophycus*, we accept the most useful and workable nomenclatural system as that of Pemberton and Frey (1982) and adopt it herein. Nevertheless, we slightly emend their diagnosis to include lined forms, which, although rare in the Trenton Group are consistent with the overall diagnosis of *Planolites* as originally described by Nicholson (see Benton and Trewin 1978, pp. 5-8).

Planolites is a eurybathic form (Chamberlain 1978b) ranging in age from Late Precambrian-Recent (Häntzschel 1975) and is commonly attributed to the sediment-ingesting activity of vagile deposit feeding vermiform organisms (Alpert 1975, Pemberton and Frey 1982), though several phyla may in fact be responsible.

Planolites beverleyensis Billings, 1862
(Figure 9g)

Diagnosis - Relatively large, smooth, straight to gently curved or undulose cylindrical *Planolites* (after Pemberton and Frey 1982).

Material - 21 specimens and several hundreds observed in the field.

Description - Smooth, straight to slightly curved, horizontal to slightly inclined (up to 20°) regularly cylindrical, unbranched or branched (at varied angles of 20-115°), isolated or cross-cutting burrows. Burrow fill is typically coarser than the host rock but may be finer and of different colour. Burrow fill is typically structureless but rare specimens exhibit thin shale or micritic linings up to 9mm thick which may be composed of a series of concentric laminae, giving the burrows an apparent central core. Specimens are preserved in convex or concave epirelief, convex hyporelief and endorelief in a wide variety of lithologies and vary from 5-20mm in diameter and 2.7-100cm in length.

Remarks - Although the diagnosis proposed by Pemberton and Frey (1982) did not include lined burrows, those observed in the Trenton Group are so rare and are so well mixed within crowded *P. beverleyensis* that the use of another name would obscure the close similarity of the burrows. Arguably, strict adherence to the original definition of *Cylindrichnus* Toots in Howard, 1966, might suggest their inclusion within this ichnogenus. However, the original definition of *Cylindrichnus* encompasses a wide variation in morphology, often overlapping into related forms such as *Asterosoma* von Otto, 1854 and *Rosselia* Dahmer, 1937. Additionally, most authors now adopt an undefined but emended usage of *Cylindrichnus* as vertical or steeply inclined, unbranched concentrically lined structures (e.g. Fürsich 1974b, 1981; McCarthy, 1979). We are therefore hesitant to adopt *Cylindrichnus* as a useful description of these Trenton specimens and include them within *Planolites*, to which they best conform.

Otherwise, the specimens described here exhibit all the attributes of *P. beverleyensis* and are definitively conspecific.

Planolites montanus Richter, 1937
(Figure 9h)

Diagnosis - Relatively small, curved to contorted *Planolites* burrows less than 5mm in diameter (after Pemberton and Frey 1982).

Material - 10 specimens and more than a hundred observed in the field.

Description - Straight to slightly sinuous, smooth, horizontal unbranched and rarely branched (angles of 25-83°), irregularly cylindrical burrows, 1-5mm in diameter and 14-74mm in length. Cross-overs, interpenetration and reburrowed segments are common. Burrow fill is typically structureless but rare examples preserve thin wall linings (0.6mm) and, or, concentric laminae. They occur in a wide variety of lithologies and are preserved mainly in endorelief but also in convex epirelief and concave hyporelief.

Remarks - As the various ichnospecies of *Planolites* are based, according to Pemberton and Frey (1982), on size, curvature and wall sculpture, the Trenton material described here can be confidently assigned to *P. montanus*. Lined and concentrically laminated specimens, otherwise exhibiting the same basic size and morphology as *P. montanus*, are here included within the ichnospecies based on the same reasoning as discussed under *P. beverleyensis*.

Planolites sp.
(Figure 9f)

Material - 57 specimens.

Description - Horizontal, straight to slightly sinuous, unbranched, irregularly cylindrical (where observed) burrows, 3-20mm in diameter and up to 12cm in length. The burrows are most typically preserved in endorelief in calcisiltites and calcilutites, where their coarser fill and rare concentric laminae can be readily observed, and only occasionally

in convex and concave epirelief.

Remarks - Their preservation as endorelief and only occasionally as semi-relief structures, and hence the lack of knowledge of the wall characteristics and detailed 3-dimensional geometry precludes ichnospecific assignment. They are certainly larger than *P. montanus* and although extremely similar to *P. beverleyensis* differ in that they are occasionally slightly sinuous.

Ichnogenus Rhizocorallium
Zenker, 1836

Type species - *Rhizocorallium jenense*
Zenker, 1836, by original monotypy.

Diagnosis - U-shaped spreiten burrows, parallel or oblique to bedding plane, limbs more or less parallel and distinct; tube diameter: diameter of spreite 1.5 (after Fürsich 1974c).

Discussion - Two of the three ichnospecies of *Rhizocorallium*, namely *R. irregulare* Mayer, 1954 and *R. uliarensis* Firtion, 1958, are clearly the result of deposit-feeding organisms, whilst the third, *R. jenense* Zenker, 1836, is the product of a suspension feeder (Fürsich 1974c). In both cases, the most likely producers have been suggested to be crustaceans or even nymphs of ephemerids (Fürsich 1974c, Fürsich and Mayr 1981) or even possibly annelids (Basan and Scott 1979). The ichnogenus is cosmopolitan, Cambrian-Tertiary in age and although commonly reported in marginal or shallow marine strata (e.g. Hakes 1976; Fürsich 1974c, 1981) has also been reported in deep-water (Crimes 1977, Pickerill et al. 1982) and non-marine (Fürsich and Mayr 1981) environments. A full review of its nomenclatural and interpretive history is given in Fürsich (1974c).

?*Rhizocorallium* cf. *R. irregulare*
Mayer, 1954

(Fig. 10, Pickerill and Forbes 1979)

Diagnosis - Long, sinuous, bifurcating or planispiral U-shaped spreiten burrows, in the main horizontal (after Fürsich 1974c).

Material - 5 specimens.

Description - Straight, parallel or slightly oblique to stratification, isolated, U-shaped burrows, with tube diameter (where discernible) 2-3mm, total width 12-15mm and length 60-70mm with irregularly shaped and often discontinuous poorly preserved protrusive spreiten. The burrows are smooth and fill is identical to the enclosing sediment thus making for extremely poor preservation which is in both full relief and concave epirelief.

Remarks - The poor preservation and the extremely small size of the burrows precludes confident assignment to *Rhizocorallium*. However, slightly larger material and with comparable discontinuous or even no spreiten has been recently described by Fürsich and Mayr (1981) from the Miocene of southern Germany. If, indeed, the specimens are true *Rhizocorallium*, then they are conspecific with *R. irregulare* Mayer, 1954, as *R. uliarensis* is a 3-dimensional spiral and *R. jenense* is shorter and typically oblique. Several attempts to photograph the specimens proved futile but a schematic figure is reproduced in Pickerill and Forbes (1979, p. 2032, fig. 10).

Ichnogenus *Rosselia* Dahmer, 1937

Type species - *Rosselia socialis* Dahmer, 1937, by original monotypy.

Diagnosis - "Cylindrical pencil-thick burrows, filled with quartzite, dug in the sediment at various distances and orientations. Most are inclined at an angle of 30° or more to bedding. The lower end is not observed; the opening is expanded and filled with imbricated concentric layers of matrix which as a rule are strongly weathered; these structures are clearly distinctive in their appearance" (Dahmer 1937, pp. 532-533 - trans. litt.).

Discussion - Dahmer (1937) originally erected *Rosselia* for cone or funnel-shaped burrows with a concentric back-fill structure and, as outlined briefly in the discussion of *P. beverleyensis*, there has been considerable confusion regard-

ing its use with respect to the morphologically related forms *Cylindrichnus* Toots in Howard, 1966 and *Asterosoma* von Otto, 1854. *Asterosoma* should only be used for star-shaped, concentrically lined and longitudinally wrinkled burrows and, for example, the specimens figured by Frey and Howard (1970) are not consistent in any sense of the original diagnosis (cf. Häntzschel 1975). Many authors now employ *Cylindrichnus* for vertical or steeply inclined, straight to weakly curved, unbranched *Skolithos*-like structures with multiple layered walls (e.g. Fürsich 1974a, 1981; McCarthy 1979), that is in contradiction to its original and extremely broad diagnosis (see Howard 1966). Strict adherence to this emended usage is to be recommended, because once morphologically similar structures assume a cone or funnel-shape, they should be included within *Rosselia* and *Asterosoma*. Although the three ichnogenera are clearly end members of intergradational burrow systems (e.g. Chamberlain 1971, Fürsich 1974a, McCarthy 1979), providing caution is observed, ichnogenetic assignment should be possible.

Rosselia ranges in age from Lower Cambrian (Seilacher 1955) to Upper Eocene (Książkiewicz 1977) and although most commonly reported from shallow sublittoral environments (Garcia-Ramos 1976, McCarthy 1979, Pemberton et al. 1982) has also been reported in association with deep-water flysch (Książkiewicz 1970, 1977). It may represent the feeding or dwelling burrows of annelids (Chamberlain 1971), crustaceans (Frey 1970) or sea anemones (Chamberlain and Clark 1973, Książkiewicz 1977).

?*Rosselia* sp.
(Figures 10f, g)

Material - 10 specimens.

Description - Vertical to slightly inclined, funnel-shaped burrows preserved only in concave epirelief on upper surfaces of thinly bedded (20-35mm) calcilutites. Burrow diameter is 4-9mm; depth is up to 9mm. In most examples the filling has been eroded but some

exhibit a poorly preserved concentric wall surrounding a central tube up to 3mm in diameter.

Remarks - The material is so poorly preserved even ichnogenic identification is uncertain. However, its vertical attitude and funnel-like cross-section and its poorly preserved concentric layering and central tube are sufficient to tentatively assign it to *Rosselia* Dahmer, 1937.

Ichnogenus *Rusophycus* Hall, 1852

Type species - *Fucoides biloba* Vanuxem, 1842, first? subsequent designation by Seilacher (1955, p. 110) as *R. bilobatus* (= *R. bilobatum*).

Diagnosis - Shallow to deep, short, posteriorly tapering bilobed resting burrows (cubichnia) of trilobites. Lobes may be smooth or exhibit transverse to oblique scratch marks in various arrangements directed anterolaterally. Coxal, exopodal, spinal, cephalic and pygidial markings may be present (after Osgood 1970, Alpert 1976).

Discussion - If the ichnogenera *Cruziana* and *Rusophycus* are to be retained as distinctive forms, the designations by Miller (1889, p. 138), Bassler (1915, p. 1132), Andrews (1955, p. 230) and Hakes (1976, p. 32) of the type species of *Rusophycus* as *R. clavatus* must be rejected, as Hall's (1852) original material should more correctly be diagnosed as *Cruziana clavata* (Hall), as suggested by Osgood and Drennen (1975). Other authors (e.g. Osgood 1970, Häntzschel 1975, Osgood and Drennen 1975, Alpert 1976) quote *Fucoides biloba* (Vanuxem) as the type, even though Hall (1852) did not designate *R. bilobatus* (= *R. bilobatum*) as his nomenclatural type. Arguably, his *R. pudicus* (= *R. pudicum*) could equally be regarded as the type as it was described prior to *R. bilobatus*. To retain some semblance of nomenclatural stability, however, particularly as *R. bilobatum* is more closely representative of typical rusophycids, as originally inferred by Hall himself (1852, p. 23), we accept *F. biloba* Vanuxem, 1842 as the type species.

The morphology, interpretation and detailed taxonomy of *Rusophycus* has been fully discussed by Osgood (1970) and Osgood and Drennen (1975). It ranges in age from Cambrian-Triassic (Crimes 1975, Bromley and Asgaard 1979) and, following Seilacher (1955) and Osgood (1970), was produced by trilobites and is a trace diagnostic of the *Cruziana* ichnofacies of Seilacher (1967).

Rusophycus sp. (Figure 7h)

Material - 9 specimens.

Description - Variably and generally poorly preserved (in convex hyporelief) posteriorly tapering bilobate traces; maximum length 44mm, maximum width 26mm and a mean length/width ratio of 1.7:1. Depth variable, up to 8mm. Lobes are smooth and outer margins grade imperceptibly into surrounding material.

Remarks - Poor preservation and the absence of detailed scratch markings precludes ichnospecific assignment. They are certainly extremely similar to specimens of *Rusophycus* described from the Trenton Group northeast of Quebec City by Pickerill and Forbes (1979), who noted a close similarity to *R. pudicum*. Sectioning reveals the traces were formed at the sediment-water interface.

Ichnogenus *Scalarituba* Weller, 1899

Type species - *Scalarituba missouriensis* Weller, 1899, by original monotypy.

Diagnosis - Mainly horizontal "burrows, subcylindrical in form, never straight for more than a few centimeters, curving in all directions, and marked by transverse ridges situated at distances of one or two millimeters, filled with material similar to that which surrounds them, detected by their slightly different color" (after Weller 1899, p. 12).

Discussion - *Scalarituba* has been the subject of much nomenclatural controversy since Seilacher and Meischner (1965) first considered it to be a behavioural variant but not strictly synonymous with associated *Nereites* and *Neo-*

nerites traces. Chamberlain (1971), with extremely well-preserved material, demonstrated that for a single specimen, preservation in hyporelief resulted in biserially arranged pustules characteristic of *Neonereites biserialis* whereas preservation in epirelief resulted in a median gallery with lateral lobes characteristic of *S. missouriensis*, *Nereites* sp. (= *Phyllodocites* view) and *Neonereites uniserialis*. He placed *N. uniserialis* and *N. biserialis* in synonymy with *S. missouriensis* and expanded *Scalarituba* to include a 'Nereites view' in convex hyporelief and a 'Phyllodocites view' in concave epirelief. Although this approach has subsequently gained a large following (e.g. Chamberlain 1977, 1978b; Chaplin 1980; Jordan 1981; Larson 1982) several authors have preferred to retain the forms as distinctive ichnogenera, particularly because poorer preserved material than that described by Chamberlain (1971) does not illustrate the morphological overlap and, instead, conforms more closely to individual ichnogenera as originally defined (e.g. Häntzschel 1975; Hakes 1976; Brasier and Hewitt 1979; Pickerill 1980, 1981; Marintsch and Finks 1982). We adopt the latter approach herein.

Scalarituba represents the deposit-feeding activity of a worm-like organism burrowing in the sediment and periodically backfilling its burrow (Weller 1899, Conkin and Conkin 1968). It is a marine but eurybathic form (Crimes 1977) having been reported from a wide spectrum of environments ranging from tidal flat (Conkin and Conkin 1968) to deep-water flysch (Chamberlain 1971, Pickerill 1980). It ranges in age from Ordovician-Permian (Häntzschel 1975).

Scalarituba missouriensis Weller, 1899
(Figure 10c)

Diagnosis - As for ichnogenus

Material - 5 specimens.

Description - Horizontal, sinuous meniscate burrows, that straighten over short distances, preserved in endorelief on upper surfaces of calcilutites. Burrow fill is identical to host material but occas-

ionally meniscae possess coarser bioclastic material. Burrow diameter is 2.5-5mm and is consistent within a single specimen; length is variable, a maximum of 18cm having been observed. Meniscae are irregularly distributed but up to 3 per cm may be observed.

Remarks - The Trenton material can undoubtedly be diagnosed as *S. missouriensis* and is extremely similar to conspecific material described by Pickerill (1981) and by Marintsch and Finks (1982) as *S. missouriensis* var. "meniscate form".

Ichnogenus *Scolicia* de Quatrefages, 1849

Type species - *Scolicia prisca* de Quatrefages, 1849, by original monotypy.

Diagnosis - Concave, top-surface meandering trails broadly U-shaped in section with floor and lateral walls. Most of the floor shows a single or biserially alternating series of laminae, but the edges of the floor are smooth tracks, sometimes consisting of discrete sediment strings. Lateral walls with distinct curved laminae, concave surface anterior and steep (after Smith and Crimes 1983).

Discussion - *Scolicia* was erected from the Paleocene flysch of Spain but its original description was inadequate and had no accompanying figure (Smith and Crimes 1983). This led to utter confusion in the literature and the "Scolicia group" now comprises extremely variable traces interpreted as the trails of annelids, gastropods, crustaceans, polychaetes, echinoids, holothuroids, Pennatulacea (as reviewed by Bradley 1980) and more recently by Smith and Crimes (1983) as heart urchins. Following Książkiewicz (1977), Crimes et al. (1981) and Smith and Crimes (1983) used *Scolicia* for either shallow burrows or furrows and *Subphyllochorda* for cylindrical backfilled burrows. Although this approach is followed herein we do not agree with the suggestion of Smith and Crimes (1983) that all Palaeozoic forms referred to *Scolicia* should be transferred to other ichnogenera because they cannot possibly be related to heart urchins. Knowledge of the producer of a trace fossil is not taxonomically significant in ichnology

(see Fillion and Pickerill 1984). Clearly, a full and detailed taxonomic revision of the ichnogenus is still required and until such a study is undertaken we regard *Scolicia* as a variable and facies transgressive form which ranges in age from Lower Cambrian to Recent (Häntzschel 1975).

Scolicia sp.
(Figure 7c)

Material - 3 specimens and several occurrences on a single bedding plane.

Description - All specimens are preserved on the upper surface of a 5mm thick chert bed in concave epirelief and consists of two parallel furrows, each 4mm wide, separated by a slightly raised central lobe 4-4.5mm wide. The latter is smooth and rounded whereas the furrows exhibit crude transverse striations oriented approximately normal to the central lobe, usually less than 2mm apart. Total depth of the furrows is 1.5 mm. The traces are sinuous, often tens of centimetres in length and never self-crossing.

Remarks - Because of the generally poor preservation of the traces and the unsatisfactory taxonomic status of the ichnogenus, the Trenton material is only identified at the ichnogeneric level. Although we know of no identical analogues, they do resemble specimens of *Scolicia* figured by both Garcia-Ramos (1976) from the Devonian of Spain and Chamberlain (1971) from the Carboniferous of Oklahoma.




Fig. 12 - Height: Diameter plot of intact and bored, symmetric and asymmetric colonies of the bryozoan *Prasopora* spp.

Ichnogenus *Skolithos* Haldeman, 1840

Type species - *Fucoides? linearis* Haldeman, 1840, by original monotypy.

Diagnosis - Unbranched, vertical or steeply inclined, cylindrical or subcylindrical, lined or unlined burrows, 1-15mm in diameter and a few centimetres to up to a meter in length. Burrow walls distinct or indistinct, smooth to rough, possibly annulated; burrow diameter may vary slightly along its length. Burrow fill structureless (after Alpert 1974).

Discussion - As with several other commonly occurring ichnogenera, *Skolithos* has received considerable discussion and systematic treatment. Alpert (1974) recently published a systematic review, conducted solely on literature research, and suggested that many of the numerous previously published ichnospecies and even ichnogenera such as *Tigillites*, *Stipsellus* and *Foralites* could confidently be placed within five ichnospecies of *Skolithos*. He later proposed a sixth, *S. bulbus* (Alpert 1975), and more recently Hofmann (1979) erected a new ichnospecies, *S. gyratus*. James (1891), Richter (1920, 1921), Howell (1943) and Osgood (1970) have also extensively discussed the ichnogenus. Undoubtedly, a thorough systematic review still seems necessary, but until this is undertaken we regard Alpert's (1974, 1975) scheme as the most satisfactory.

Skolithos has been reported from a variety of marine environments and is typically found in association with 'high energy' conditions, particularly in near-shore shallow-water conditions but also in deep-water submarine channels and canyons (see Seilacher 1967, Alpert 1974, Crimes 1977, Pickerill 1981 etc.) or even in floodplain and dune environments (e.g. Ratcliffe and Fagerstrom 1980). It constitutes the type ichnogenus of the *Skolithos* ichnofacies of Seilacher (1967) and is generally regarded as the dwelling and feeding burrow of annelids or phoronids (Alpert 1974) and ranges in age from Late Precambrian-Recent (Crimes and Germs 1982).

Skolithos linearis Haldeman, 1840
(Figure 13a)

Diagnosis - Burrows cylindrical to subcylindrical, perfectly straight to slightly curved and inclined, up to a meter in length. Diameter 3-7mm, occasionally up to 12mm. Burrow wall distinct to indistinct, may be annulated (after Alpert 1974, p. 663).

Material - 13 specimens.

Description - Straight, vertical, unbranched burrows 3-5mm in diameter and up to 66mm in length. Burrow fill is typically coarser than host material, rarely finer. Burrow walls are distinct, parallel, apparently smooth and unlined. Preservation is in endorelief.

Remarks - Of the seven existing ichnospecies of *Skolithos*, the Trenton material most closely resembles *S. linearis* Haldeman, 1840 and is diagnosed as such. Although they superficially resemble *S. verticalis* (Hall), this ichnospecies is generally shorter and smaller and more commonly inclined and curved (Alpert 1974).

Skolithos sp.
(Figure 10d)

Material - 4 specimens and several occurrences in the field.

Description - Straight, vertical to steeply inclined (up to 60°) burrows 4-10mm in diameter and up to 35mm in length. Burrow fill is finer or identical to the host rock (calcisiltite). Burrow walls are distinct and irregular; some specimens possess a thick calcilutite wall (2-3mm) around an inner tube of 4mm diameter. Preservation is in endorelief.

Remarks - Though clearly lacking the characteristics of *S. linearis*, the specimens are not well enough preserved for ichnospecific identification. Similar structures have been figured by Pickerill and Forbes (1979, p. 2033, fig. 116).

Ichnogenus Teichichnus Seilacher, 1955

Type species - *Teichichnus rectus*, Seilacher 1955, by original monotypy.

Diagnosis - Long wall-shaped septate

structures consisting of a pile of gutter shaped laminae (Seilacher 1955, p. 378 -trans. litt.).

Discussion - *Teichichnus* was introduced by Seilacher (1955) to describe a type of spreiten trace fossil formed by a series of long horizontal burrows stacked vertical to bedding. He interpreted the burrow as resulting from continual upward-shift by the producing deposit-feeding organism. Since its original description, it has been shown to intergrade with *Phycodes* (Häntzschel and Reineck 1968, Hofmann 1979), *Thalassinoides*, *Ophiomorpha* and *Gyrolithes* (Hester and Pryor 1972, Frey et al. 1978, Frey and Seilacher 1980), *Rhizocorallium* (Chisholm 1970, Sellwood 1970) and even *Cruziana* (Garcia-Ramos 1976). As such, its producers are probably polyphyletic, though annelids and arthropods are usually most favoured (Martinsson 1965, Chisholm 1970, Chamberlain 1977).

It is a strictly marine but eurybathic form, most commonly occurring in shallow subtidal sediments (e.g. Fürsich 1981) but also in association with submarine fans and abyssal environments (Ekdale 1977, Wetzel 1981). It ranges in age from Lower Cambrian-Recent.

Teichichnus rectus Seilacher, 1955
(Figure 10c)

Diagnosis - Straight, unbranched *Teichichnus* with an exclusively retrusive spreite (after Seilacher 1955).

Material - 138 specimens.

Description - Straight, smooth, unbranched burrows, parallel or slightly oblique to stratification possessing retrusive spreiten. Length 8.5-125mm, width 4-35mm and height 14-25mm; individual specimens are consistently deeper than wide, are preserved in full relief and possess a calcilutite or calcisiltite fill of identical grain size to host rock.

Remarks - *T. rectus* is the product of a shallow burrower usually associated with low sedimentation rates (Fürsich 1975, Hakes 1976) but occasionally observed in environments subject to rapid deposition (Chisholm 1970). Specimens from the Trenton Group within Quebec City




Fig. 13 - a. *Skolithus linearis* in vertical section, Grondines Member, locality 18. b. 'Oblique burrow' in transverse section, Grondines Member, locality 18. c. 'Loop burrows' on lower bedding surface in convex hyporelief, Grondines Member, locality 21. d. 'Pronged burrows' with typical prongs arrowed, on upper bedding surface, Grondines Member, locality 21.

have also been figured in Pickerill and Forbes (1979, p. 2031, figs. 9c, 9d).

***Teichichnus* sp.**
(Figure 10h)

Material - 12 specimens.

Description - All specimens preserved in full relief on upper bedding plane surfaces as elongate, unbranched, straight endoreliefs with parallel sides and rounded terminations. Length 50-105mm, width 11-20mm and depth, where observed, 16-21mm. Plan views exhibit bidirectional spreite concave towards the centre of the burrow; transverse sections indicate

spreite are concave-up and retrusive.

Remarks - The total character of this form is not fully preserved and not clearly understood, but the absence of a tube clearly distinguishes them from the similar *Polarichnus* Narbonne (in Narbonne et al. 1979). They most closely accord to *Teichichnus* Seilacher, 1955 and are diagnosed as such but only at ichnogenetic level.

Ichnogenus *Trichichnus* Frey, 1970

Type species - *Trichichnus linearis* Frey, 1970, by original designation.

Diagnosis - Branched or unbranched, hairlike, cylindrical, straight to sinuous burrows distinctly 1mm in diameter, oriented at various angles with respect to bedding. Burrow walls distinct, commonly lined with diagenetic minerals (after Frey 1970).

Discussion - Frey (1970) interpreted *Trichichnus* as the feeding and dwelling burrow of a minute deposit-feeding vermiform or crustacean-like organism, and since its original formulation remains a monospecific form, even though it has been reported from strata of Tremadoc-Recent age (Wetzel 1981, Fillion and Pickerill 1982). It is strictly marine but eurybathic, having been reported from shallow (Frey 1970, 1975; Pickerill and Forbes 1979) and less commonly deep-water (Kennedy 1975) environments.

Trichichnus sp.
(Figure 10a)

Material - 15 specimens and several occurrences recorded in the field.

Description - Unbranched, straight to slightly curved, thin cylindrical burrows less than 1mm in diameter and up to 30mm long. Commonly vertical but may be inclined and rarely horizontal. Burrow walls distinct and fill is typically coarser than enclosing material but may be finer. Preservation is in full relief.

Remarks - The Trenton material is only identified at ichnogenetic level as specimens lack the diagenetic wall linings diagnostic of *T. linearis* (see Frey 1970, p. 22), thus suggesting that the producing organism was not required to reinforce its burrow. Pickerill and Forbes (1979, p. 2031, fig. 9e) have also figured identical material from the Trenton Group at Pont Rouge (see Fig. 1).

Ichnogenus *Trypanites* Mägdefrau, 1932

Type species - *Trypanites weisei* Mägdefrau, 1932, by original monotypy.

Diagnosis - Simple, unbranched borings in hard substrates, non-parallel to the

surface with a single opening to the surface (after Bromley 1972).

Discussion - As recently noted by Bradshaw (1980), Mägdefrau (1932) named *Trypanites weisei*, the type species, for slender more or less straight-sided and long (up to 4cm) but very narrow (1mm) vertical borings in the Middle Triassic Muschelkalk hardgrounds. At a later date, however, Bromley (1972) expanded *Trypanites* to include all single-entrance pouch-shaped borings (e.g. *Teredolites*, *Gastrochaenolites*) resulting from the activities of a variety of organisms but excluding acrothoracican cirripeds. The expanded definition has generally been accepted apart from the inclusion by Bromley (1972) of *Vermiforichnus* Cameron, 1969, which is still adopted by many authors (e.g. Pickerill 1976, Elias 1976, Hofmann 1979) as a useful descriptor of unbranched horizontal or subhorizontal cylindrical borings. In this paper we also exclude borings more closely akin to *Vermiforichnus* and adopt *Trypanites* as originally defined by Mägdefrau (1932).

Trypanites ranges in age from Lower Cambrian (James et al. 1977) to Recent (Kobluk et al. 1978) and is commonly attributed to the activities of polychaetes, sipunculids, phoronids and thoracican cirripeds of which the sipunculids (Pemberton et al. 1980) or polychaetes (Kobluk and Nemcsok 1982) are most favoured. It constitutes the type of the recently defined *Trypanites* ichnofacies of Frey and Seilacher (1980).

Trypanites weisei Mägdefrau, 1932
(Figures 11b, c)

Diagnosis - More or less straight and vertical *Trypanites*, diameter ca. 1mm and reaching over 4cm in length (after Bromley 1972).

Material - 97 specimens and several hundred recordings in the field.

Description - All borings are preserved in colonies of the calcareous red alga *Solenopora compacta* or the bryozoans

Prasopora spp. and **Diplotrypa** spp. Diameter is 0.1-3mm and length up to 17mm; they are straight to gently curved, parallel sided with clearly defined unlined walls and typically possess a micrite fill. In **Prasopora**, walls of the borings are smooth provided they parallel zoecia but as soon as they turn they become ragged.

Remarks - Restriction of the borings to **Solenopora**, **Prasopora** and **Diplotrypa** suggests host specificity as a result of larval selectivity of the producing sipunculids or polychaetes probably as a result of the availability of the large surface area substrates afforded by these genera. As all the examined colonies do not show any reaction to the borings we also conclude that individual colonies were dead prior to boring activity. In a recent study of comparable material from the Middle Ordovician of Ontario, Kobluk and Nemcsok (1982) suggested that the producing polychaetes preferred large and asymmetrical **Prasopora** colonies. However, our data (Fig. 12) suggest that populations of bored and intact colonies thoroughly overlap and symmetrical and asymmetrical colonies were bored indiscriminately.

Ichnogenus **Vermiforichnus**
Cameron, 1969

Type species - **Vermiforichnus clarkei** Cameron, 1969, by monotypy.

Diagnosis - Unbranched, horizontal or subhorizontal, straight to slightly curved, rarely irregular hooked or coiled, non-intersecting borings (after Cameron 1969a, b).

Discussion - Although Bromley (1972) included **Vermiforichnus** within **Trypanites**, we regard it as a useful descriptor of horizontal or subhorizontal unbranched borings, as discussed under the description of **Trypanites** in this paper. The borings have been reported in a wide variety of hosts (Cameron 1969a, b) and range in age from Middle Ordovician-Permian (Kobluk et al. 1978). Cameron (1969a, b) has demon-

strated that at least some **Vermiforichnus** were produced by spionid polychaetes and Pickerill (1976) has demonstrated at least a single example of its host and size specificity.

Vermiforichnus clarkei Cameron, 1969
(Figures 11a, d)

Diagnosis - As for ichnogenus.

Material - 3 specimens.

Description - (i) A single boring preserved in endorelief in a pygidium of the isotelinid **Isotelus gigas**, 4.5mm long and 1-1.3mm wide with slightly expanded terminations. The boring is smooth, slightly sinuous and is filled with calcisiltite (Fig. 11a). (ii) A single boring preserved in endorelief on an undetermined branching bryozoan, 11mm long and 2mm wide. The boring is smooth, straight with rounded terminations and is filled with partially dolomitized sparite. (iii) Numerous smaller unbranched borings preserved in endorelief surrounding the right eye of **Isotelus gigas**, 0.09-0.16mm wide and 0.6-1mm long. The borings are straight to slightly sinuous, crowded but non-intersecting (Fig. 11d).

Remarks - Despite the small number of specimens, the variability exhibited by the borings of **V. clarkei** in the Trenton Group suggests that different groups of organisms were responsible for its production. Type (iii) borings differ from those of **Talpina pungens** Quenstedt, 1849 in being parallel and not oblique to the surface (see Voigt and Soule 1973).

Ichnogenus **Zoophycos**
Massalongo, 1855

Type species - **Zoophycos caputmedusae** Massalongo, 1855, by subsequent designation of Andrews (1955, p. 262).

Diagnosis - Variably shaped spreiten structures comprised of numerous small protrusive more or less U- or J-shaped burrows of variable length and orientation. Spreite tabular or arranged in helicoid spirals giving an overall outline of circular, elliptical or lobate

shape; central vertical tunnel or marginal tube may be present (after Frey 1970, Häntzschel 1975, Wetzel and Werner 1981).

Discussion - The ichnogenus *Zoophycos* has been plagued with excessive taxonomic splitting, different species, many of which are probably synonymous, being created merely on the whorl shape and the width and manner of branching of the rays (Frey, 1970, Häntzschel 1975, Książkiewicz 1977). Furthermore, there is still disagreement on the type species, because following the designation of the type as *Z. caputmedusae* by Andrews (1955, p. 262), Häntzschel (1962, p. 220), Taylor (1967, p. 4) and Plička (1968, p. 840) respectively designated *Fucoides brianteus*, *Zoophycos laminatus* and *Fucoides circinatus*. Häntzschel (1962), however, noted his designation only by an asterisk, and although still adopted as the type by several recent authors (e.g. Marintsch and Finks 1982) we regard such a designation as invalid by rule of priority. The original illustration of *Fucoides circinatus* by Brongniart (1828, p. 83) is too schematic to ascertain its true nature and *Zoophycos laminatus* is a *nomen nudum*. Until resolved, therefore, we accept Andrews' (1955) designation of the nomenclatural type as *Z. caputmedusae* by rule of priority.

Although there is still no general agreement on its origin, *Zoophycos* has been commonly attributed to the deposit-feeding activity of polychaetes (Bischoff 1968) and sipunculids (Wetzel and Werner 1981), to the imprints of prostomia of sabellids (Plička 1970) and even the feeding activity of umbellulids (Pennatulaceids) (Bradley 1973). Though considerable environmental significance has been attached to the ichnogenus, particularly regarding bathymetry, it is now realized to be a eurybathic form ranging in age from Ordovician-Quaternary and is typical of fine-grained substrates characterized by low sedimentation rates (for reviews see Frey 1970, Kennedy 1970, Simpson 1970, Chamberlain 1971, Osgood and

Szmuc 1972, Bradley 1973, Häntzschel 1975, Ekdale 1977, Książkiewicz 1977, Wetzel and Werner 1981, Marintsch and Finks 1982).

Zoophycos sp.
(Figure 10b)

Material - 1 specimen.

Description - Horizontal arcuate burrow 7.5cm in width and preserved in epirelief, possessing disturbed back-filled coarse spreiten which are straight to moderately curved, composed of bioclastic material, ca. 3-6mm wide and separated by furrows up to 10mm wide. A J-shaped tube, 6.5mm wide and filled with bioclastic material bounds the burrow and gives it its particular outline.

Remarks - Because of the poor preservation identification at the ichnospecific level is uncertain, but the structure clearly has a J-form (*sensu* Wetzel and Werner 1981) in which only one side of the spreite is bounded by a tube, and is therefore regarded as *Zoophycos*. It resembles parts of *Zoophycos insignis* as illustrated by Książkiewicz (1977, pl. 10, figs. 1, 7, text-fig. 16) but otherwise we are unaware of any comparable form.

MISCELLANEOUS TRACES

In addition to the morphologically distinctive forms recognized from the Trenton Group several forms are present which, because of their poor preservation, morphological variability and sufficient differences in configuration to already established ichnogenera, or combinations thereof, are not formally diagnosed. Four broad categories may be recognized:

(i) 'Loop burrows' (Fig. 13c) - Irregularly meandering and winding, subcylindrical, unlined horseshoe or more commonly loop-shaped burrows, 4-11mm wide and up to 28cm long, preserved in convex hyporelief on the soles of calcilutites. Burrow fill is coarser than enclosing sediment. Although looped, the burrows never inter-

sect; rather they go under or over previous pathways. They are obviously the product of an unknown deposit feeder and resemble *Lumbricaria* as figured by Gutschick and Rodriguez (1977, p. 206, pl. 1a-c).

(ii) 'Oblique burrows' (Fig. 13b) - Straight, unbranched, inclined (30-50°) cylindrical burrows preserved in endorelief, 4-9mm in diameter and downward tapering to a rounded or pointed termination. Length 20-35mm, walls distinct and regular, lined or more commonly unlined, with a structureless burrow fill of identical grain size to enclosing material. We were initially tempted to include these structures within *Obliquus* Marintsch and Finks, 1982; however, this ichnogenus is consistently lined, inclined at 45° or less and upper portions may approach the horizontal and lower portions the vertical. Furthermore, there are difficulties in distinguishing this recently formed ichnogenus from *Cylindrichnus* as originally defined (Toots in Howard 1966) and inclined specimens of *Palaeophycus*.

(iii) 'Pronged burrows' (Fig. 13d) - Straight, unbranched, elongate burrows preserved in epirelief on the upper surfaces of calcilutites, 4-8mm wide, up to 9cm long. Burrow fill is typically structureless calcilutite but occasionally poorly preserved spreite can be discerned. Irregular prongs up to 4mm in width and length project from the burrows (Fig. 13d). Similar unnamed material has been figured by Ward and Lewis (1975, p. 884, fig. 2) and Pickerill and Forbes (1979, p. 2033, fig. 11a). The presence of a spreite in some specimens suggests the burrows were produced by deposit-feeding organisms and their overall morphology suggests they are akin to teichichnids.

(iv) 'Bryozoan borings' (Figs. 11e, f) - Delicate pinnate or net-like systems of tear-drop shaped zoecial pits, each 0.3-0.5mm in length, 0.07-0.1mm in width, rarely adjoined and more commonly separated by stolomate impressions up to 1mm long and 0.02mm

wide preserved in two pygidia of *Iso-telus gigas*. Similar borings or etching traces have been figured by Bromley and Surlyk (1973, p. 362, fig. 14c), Warme (1975, p. 201, fig. 11-14), Pickerill and Forbes (1979, p. 2033, fig. 11g) and Hofmann (1979, p. 57, pl. 22) and all have been interpreted as bryozoan in origin.

CONCLUSIONS

As outlined previously, sediments of the Trenton Group were deposited initially in lagoons followed in turn by offshore "bar", shallow and, finally, deeper offshore shelf environments. Trace fossils in each of these environments may be summarized as follows (see also Fig. 6):

Lagoon - *Arenicolites* sp., *Chondrites* type A, *Chondrites* type B, *Clematischnia* sp., *Helminthopsis hieroglyphica*, *Helminthopsis* sp., *Palaeophycus tubularis*, *Palaeophycus* sp., *Planolites beverleyensis*, *Planolites montanus*, *Planolites* sp., *Scalarituba missouriensis*, *Skolithos linearis*, *Skolithos* sp., *Teichichnus rectus*, ?*Teichichnus* sp., *Trichichnus* sp., *Trypanites weisei* and oblique burrows.

"Bar" - ?*Calycraterion* sp., *Chondrites* type A, *Chondrites* type B, *Chondrites* spp., *Circulichnis montanus*, *Cruziana problematica*, *Helminthopsis* sp., *Palaeophycus tubularis*, ?*Plagiogmus*, *Planolites beverleyensis*, *Skolithos linearis*, ?*Skolithos* sp. and *Trypanites weisei*.

Shallow shelf - *Arenicolites* sp., *Chondrites* Type A, *Chondrites* type B, *Chondrites* spp., *Cruziana problematica*, *Furculosus carpathicus*, *Helminthopsis hieroglyphica*, *Helminthopsis* sp., *Palaeophycus tubularis*, *Planolites beverleyensis*, *P. montanus*, *Planolites* sp., ?*Rhizocorallium* cf. *R. irregulare*, ?*Rosselia*, *Rusophycus* sp., ?*Scalarituba missouriensis*, *Skolithos linearis*, *Skolithos* sp., *Teichichnus* sp., *Trichichnus* sp., *Trypanites weisei* and *Zoophycos* sp.

Offshore shelf - *Arenicolites* sp., *Chondrites* type B, *Chondrites* spp., *Clem-*

atischnia sp., ?Conostichnus, Cruziana sp. cf. Diplichnites sp., Helminthopsis sp., Oichnus paraboloides, Palaeophycus tubularis, Palaeophycus sp., Planolites beverleyensis, P. montanus, Planolites sp., ?Rosselia, Rusophycus sp., Scalarituba missouriensis, Scolicia sp., Skolithos linearis, Skolithos sp., Teichichnus rectus, Teichichnus sp., Trypanites weisei, Vermiforichnus clarkei and informally diagnosed loop, oblique and pronged burrows and bryozoan borings.

Although there is some obvious variation in the abundance of traces in different environments (Fig. 6), this is believed to reflect preservational and other hazards such as nature and quality of outcrop, burrow density, topographic or diagenetic preservation, grain size and sediment stability as discussed more fully by Fillion et al. (1984). The entire assemblage, irrespective of environment of deposition, can best be equated with the Cruziana ichnofacies of Seilacher (1967), which is characterized by a mixed association of vertical, inclined and particularly horizontal deposit-feeding but also filter feeding traces. The abundant borings, more characteristic of the recently formulated Trypanites ichnofacies of Frey and Seilacher (1980), are no cause for concern as their presence merely reflects the availability of abundant hard substrates, in this case essentially in the form of the bryozoans *Prasopora* spp. and *Diplotrypa* spp., the red alga *Solenopora compacta* and the isotelenid trilobite *Isotelus gigas*.

Finally, and as discussed more fully by Fillion et al. (1984), the most significant environmental parameter governing the production, nature and preservation of the trace fossils was substrate and its overall characteristics, which in turn reflected the prevailing hydrodynamic regime and depositional rate. Substrate not only provided a primary control but also directly influenced diagenetic affects which both enhanced and masked specific traces depending on their original character.

ACKNOWLEDGEMENTS

We wish to thank Dr. Yvon Globensky and Dr. John Riva for their continual help and encouragement in our research in the Trenton Group over the past several years and Dr. Guy Narbonne and Dr. Hans Hofmann for providing us with certain relevant literature and advice on terminology. The manuscript was reviewed by Drs. T.P. Crimes and T.L. Harland and for their constructive comments we are extremely grateful. One of us (R.K.P.) is grateful to Dr. Crimes for permitting access to his abundant trace fossil collections and for permission to quote unpublished data. Dr. Harland accompanied the authors in the field on several occasions and for his astute observations and keen eye we are equally grateful. The work was undertaken during the tenure by R.K. Pickerill of Natural Sciences and Engineering Research Council of Canada Grant A3857 and the tenure of D. Fillion of scholarships from Bourse de l'Enseignement supérieur du ministère de l'Éducation du Québec and Bourse du Fonds F.C.A.C. pour l'aide et le soutien à la recherche, all of which are gratefully acknowledged. The manuscript was prepared by J. Ley and technical assistance provided by R. McCulloch, S. Aitken and J. Lynch.

- ABEL, O. 1935. Vorzeitliche Lebensspuren. Gustav Fisher, Jena, 644p.
- ALPERT, S.P. 1974. Systematic review of the genus *Skolithos*. *Journal of Paleontology* 48, pp. 661-669.
- , 1975. *Planolites* and *Skolithos* from the Upper Precambrian-Lower Cambrian White-Inyo Mountains, California. *Journal of Paleontology* 49, pp. 509-521.
- , 1976. Trilobite and star-like trace fossils from the White-Inyo Mountains, California. *Journal of Paleontology* 50, pp. 226-240.
- , 1977. Trace fossils and the basal Cambrian boundary. In *Trace fossils 2*. Edited by T.P. Crimes and J.C. Harper. Geological Journal Special Issue 9, Seel House Press, Liverpool, pp. 1-8.
- ANDREWS, H.N. Jr. 1955. Index of generic names of fossil plants, 1820-1950. United States Geological Survey, Bulletin 1013, 262p.
- BANDEL, K. 1973. A new name for the ichnogenus *Cylindrichnus* Bandel, 1967. *Journal of Paleontology* 47, p. 1002.
- BARTHEL, K.W. and BARTH, W. 1972. Paleologic specimens from the Devonian of Bolivia. *Neues*

- Jahrbuch für Geologie und Paläontologie, Monatshefte 1972, pp. 573-581.
- BASAN, P.B. and SCOTT, R.W. 1979. Morphology of *Rhizocorallium* and associated traces from the Lower Cretaceous Purgatoire Formation, Colorado. *Palaeogeography, Palaeoclimatology, Palaeoecology* 28, pp. 5-23.
- BASSLER, R.S. 1915. Bibliographic index of American Ordovician and Silurian fossils. *Bulletin of the United States National Museum*, 92, 718p.
- BANKS, N.L. 1970. Trace fossils from the late Precambrian and Lower Cambrian of Finnmark, Norway. In *Trace fossils*. Edited by T.P. Crimes and J.C. Harper. *Geological Journal Special Issue 3*. Seel House Press, Liverpool, pp. 19-34.
- BATHER, F.A. 1925. U-shaped burrows near Blea Wyke. *Proceedings of the Yorkshire Geological Society* 20, pp. 185-199.
- BENTON, M.J. and TREWIN, N.H. 1978. Discussion and comments on Nicholson's 1872 manuscript "Contributions to the study of errant annelids of the older Palaeozoic rocks". Publication of the Department of Geology and Mineralogy, University of Aberdeen, 1, pp. 1-16.
- BERGSTRÖM, J. 1976. Lower Palaeozoic trace fossils from eastern Newfoundland. *Canadian Journal of Earth Sciences* 13, pp. 1613-1633.
- BILLINGS, E. 1862. New species of fossils from different parts of the lower, Middle and Upper Silurian rocks of Canada. In *Palaeozoic Fossils, Volume 1 (1861-1865)*. Geological Survey of Canada, pp. 96-168.
- BINNEY, E.W. 1852. On some trails and holes found in rocks of the Carboniferous strata, with remarks on the *Microconchus carbonarius*. *Manchester Literary and Philosophical Society, Memoirs and Proceedings*, series 2, 10, pp. 181-201.
- BISCHOFF, B. 1968. *Zoophycos*, a polychaete annelid, Eocene of Greece. *Journal of Paleontology* 42, pp. 1439-1443.
- BORNEMANN, J.G. 1889. Über den Buntsandstein in Deutschland und seine Bedeutung für die Trias. *Beiträge Geologie, Paläontologie* 1, 61p.
- BRADLEY, J. 1973. *Zoophycos* and *Umbellula* (Pennatulacea): Their synthesis and identity. *Palaeogeography, Palaeoclimatology, Palaeoecology* 13, pp. 103-128.
- , 1980. *Scolicia* and *Phycodes*, trace fossils of *Renilla* (Pennatulacea). *Pacific Geology* 14, pp. 73-86.
- , 1981. *Radionereites*, *Chondrites* and *Phycodes*; trace fossils of anthoptiloid sea pens. *Pacific Geology* 15, pp. 1-16.
- BRADSHAW, M.A. 1980. Boring bivalves in the New Zealand Cenozoic with a redefinition of *Teredolites*. *Records of the Canterbury Museum* 9(5), pp. 289-294.
- BRANSON, C.C. 1960. *Conostichus*. *Oklahoma Geological Notes* 20, pp. 195-207.
- BRASIER, M.D. and HEWITT, R.A. 1979. Environmental setting of fossiliferous rocks from the uppermost Proterozoic-Lower Cambrian of central England. *Palaeogeography, Palaeoclimatology, Palaeoecology* 27, pp. 35-57.
- BRIGGS, D.E.G., ROLFE, W.D.I. and BRANNAN, J. 1979. A giant myriapod trail from the Namurian of Arran, Scotland. *Palaeontology* 22, pp. 273-291.
- BROMLEY, R.G. 1972. On some ichnotaxa in hard substrates, with a redefinition of *Trypanites* Mägdefrau. *Paläontologische Zeitschrift* 46, pp. 93-98.
- 1981. Concepts in ichnotaxonomy illustrated by small round holes in shells. *Acta Geológica Hispánica, Concept and Method in Paleontology* 16(1-2), pp. 55-64.
- BROMLEY, R.G. and ASGAARD, U. 1979. Triassic freshwater ichnocoenoses from Carlsberg Fjord, east Greenland. *Palaeogeography, Palaeoclimatology, Palaeoecology* 28, pp. 39-80.
- BROMLEY, R.G. and FREY, R.W. 1974. Redescription of the trace fossil *Gyrolithes*, and taxonomic evaluation of *Thalassinoides*, *Ophiomorpha* and *Spongeliomorpha*. *Bulletin of the Geological Society of Denmark* 23, pp. 311-335.
- BROMLEY, R.G. and SURLYK, F. 1973. Borings produced by brachiopod pedicles, fossil and Recent. *Lethaia* 6, pp. 349-365.
- BRONGNIART, A.T. 1823. Observations sur les *Fucoides*. *Société d'Histoire Naturelle de Paris, Mémoires* 1, pp. 301-320.
- 1828. Histoire des végétaux fossiles ou recherches botaniques et géologiques sur les végétaux renfermés dans les diverses couches du globe, volume 1. G. Dufour and E. d'Ocagne, Paris, 136p.
- CAMERON, B. 1969a. New name for *Palaeosabella prisca* (McCoy), a Devonian worm boring, and its preserved probable borer. *Journal of Paleontology* 43, pp. 189-192.
- 1969b. Paleozoic shell-boring annelids and their trace fossils. *American Zoologist* 9, pp. 689-703.
- CHAMBERLAIN, C.K. 1971. Morphology and ethology of trace fossils from the Ouachita Mountains, southeast Oklahoma. *Journal of Paleontology* 45, pp. 212-246.
- 1977. Ordovician and Devonian trace fossils from Nevada. *Bulletin of the Nevada Bureau of Mines and Geology* 90, pp. 1-24.
- 1978a. A guidebook to the trace fossils and paleoecology of the Ouachita Geosyncline. *Society of Economic Paleontologists and Mineralogists, Tulsa (Oklahoma)*, 68p.
- 1978b. Recognition of trace fossils in cores. In *Trace fossil concepts*. Edited by P.B. Basan. *Society of Economic Paleontologists and Mineralogists, Short Course* 5, pp. 119-166.
- CHAMBERLAIN, C.K. and CLARK, D.L. 1973. Trace fossils and conodonts as evidence for deep-water deposits in the Oquirrh Basin of Central Utah. *Journal of Paleontology* 47, pp. 663-682.
- CHAPLIN, J.R. 1980. Stratigraphy, trace fossil associations and depositional environments in the Borden Formation (Mississippian), northeastern Kentucky. *Kentucky Geological Society, Annual Fall Field Trip Guidebook*. Kentucky Geological Survey, Lexington, Kentucky, 114p.
- CHISHOLM, J.L. 1970. *Teichichnus* and related trace-fossils in the Lower Carboniferous of St. Monace, Scotland. *Bulletin of the Great Britain Geological Survey* 31, pp. 19-35.
- CLARK, T.H. 1972. Montreal area. Quebec Department of Natural Resources, *Geological Report* 152, 244p.
- CLARK, T.H. and GLOBENSKY, Y. 1975. Grondines area. Quebec Department of Natural Resources, *Geological Report* 154, 159p.
- 1976a. Sorel area. Quebec Department of Natural Resources, *Geological Report* 155, 151p.

- 1976b. Laurentides area. Quebec Department of Natural Resources, Geological Reprt 157, 112p.
- 1976c. Trois Rivières area. Quebec Department of Natural Resources, Geological Report 164, 87p.
- CLARK, T.H., GLOBENSKY, Y., RIVA, J. and HOFMANN, H.J. 1972. Stratigraphy and structure of the St. Lawrence Lowland of Quebec. 24th International Geological Congress, Guidebook for Excursion C52, 83p.
- CLIFTON, H.E. and THOMPSON, J.K. 1978. *Macaronichnus segregatis*, a feeding structure of shallow marine polychaetes. *Journal of Sedimentary Petrology* 48, pp. 1293-1302.
- CONKIN, T.E. and CONKIN, B.M. 1968. *Scalarituba missouriensis* and its stratigraphic distribution. *The University of Kansas Paleontological Contributions* 31, pp. 1-7.
- CRIMES, T.P. 1970. Trilobite tracks and other trace fossils from the Upper Cambrian of North Wales. *Geological Journal* 7, pp. 47-68.
- 1975. The stratigraphical significance of trace fossils. In *The Study of Trace Fossils*. Edited by R.W. Frey. Springer-Verlag, New York, pp. 109-130.
- 1977. Trace fossils of an Eocene deep-sea fan, northern Spain. In *Trace fossils 2*. Edited by T.P. Crimes and J.C. Harper, Geological Journal Special Issue 9. Seel House Press, Liverpool, pp. 71-90.
- CRIMES, T.P. and GERMS, G.J.B. 1982. Trace fossils from the Nama Group (Precambrian-Cambrian) of southwest Africa (Namibia). *Journal of Paleontology* 56, pp. 890-907.
- CRIMES, T.P., GOLDRING, R., HOMEWOOD, P., STUIVENBERG, J. VAN and WINKLER, W. 1981. Trace fossil assemblages of deep-sea fan deposits, Gurnigel and Schlieren flysch (Cretaceous-Eocene), Switzerland. *Ecologiae Geologicae Helvetiae* 74, pp. 953-995.
- CRIMES, T.P., LEGG, I., MARCOS, A. and ARBOLEYA, M. 1977. ?Late Precambrian-low Lower Cambrian trace fossils from Spain. In *Trace fossils 2*. Edited by T.P. Crimes and J.C. Harper. Geological Journal Special Issue 9. Seel House Press, Liverpool, pp. 91-138.
- DAHMER, G. 1937. Lebensspuren aus dem Taunusquarzit und den Siegener Schichten (Unterdevon). *Preussische Geologische Landesanstalt, Jahrbuch* 57, pp. 523-539.
- DAWSON, J.W. 1873. Impressions and footprints of aquatic animals and imitative markings on Carboniferous rocks. *American Journal of Science, Series 3*, 5, pp. 16-24.
- DÖRJES, J. 1978. Sedimentologische und faunistische Untersuchungen an Watten in Taiwan. II Faunistische und aktuopaläontologische Studien. *Senckenbergiana Maritima* 10, pp. 117-143.
- EICHWALD, E. 1860. *Lethaea Rossica ou paléontologie de la Russie I*. Stuttgart, E. Schweizerbart, 1657p.
- EKDALE, A.A. 1977. Abyssal trace fossils in worldwide Deep Sea Drilling Project cores. In *Trace fossils 2*. Edited by T.P. Crimes and J.C. Harper. Geological Journal Special Issue 9. Seel House Press, Liverpool, pp. 163-182.
- 1978. Trace fossils in Leg 42A cores. Initial Reports of the Deep Sea Drilling Project 42, pp. 821-827.
- ELIAS, R.J. 1976. Solitary rugose corals of the Selkirk Member, Red River Formation (Late Middle or Upper Ordovician), southern Manitoba. Unpublished M.Sc. thesis, University of Cincinnati, Cincinnati, Ohio, 232p.
- EMMONS, E. 1844. *The Taconic System; based on observations in New York, Massachusetts, Maine, Vermont and Rhode Island*. Carroll and Cook, Albany, 68p.
- FILLION, D. and PICKERILL, R.K. 1982. Cambro-Ordovician trace fossils from Bell Island, eastern Newfoundland (abstract). *The Atlantic Geoscience Society Colloquium on Current Research in the Atlantic Provinces. Maritime Sediments and Atlantic Geology*, 18, p. 43.
- 1984. On *Arthraria anti-quata* Billings, 1872 and its relationship to *Diplocraterion Torell*, 1870 and *Bifungites Desio*, 1940. *Journal of Paleontology* 58, (in press).
- (in preparation. Trace fossils from the Lower Ordovician (Tremadoc-Arenig) of Bell Island, eastern Newfoundland.
- FIRTION, F. 1958. Sur la présence d'ichnites dans le Portlandien de l'île d'Oléron (Charente maritime). *Annales Universitatis Saraviensis, (Naturwissenschaftliche)* 7, pp. 107-112.
- FISHER, W.A. 1978. Trace fossils from the lower Harding Formation (Middle Ordovician), Colorado. In *Energy resources of the Denver Basin*. Edited by J.D. Fruit et al. Rocky Mountain Association of Geologists, Field Conference, Guidebook for 1978, pp. 191-197.
- FLOWER, R.H. 1945. Breviconic cephalopods from Pont Rouge, Quebec. *Canadian Field Naturalist* 59, pp. 74-82.
- FREY, R.W. 1970. Trace fossils of Fort Hays Limestone Member, Niobrara Chalk (Upper Cretaceous), west-central Kansas. *The University of Kansas Paleontological Contributions* 53, 41p.
- 1975. The realm of ichnology, its strength and limitations. In *The Study of Trace Fossils*. Edited by R.W. Frey, Springer-Verlag, New York, pp. 13-36.
- FREY, R.W. and HOWARD, J.D. 1970. Comparison of Upper Cretaceous ichnofaunas from siliceous sandstones and chalk, Western Interior Region, U.S.A. In *Trace fossils*. Edited by T.P. Crimes and J.C. Harper. Geological Journal Special Issue 3. Seel House Press, Liverpool, pp. 141-166.
- FREY, R.W., HOWARD, J.D. and PRYOR, W.A. 1978. *Ophiomorpha*; its morphologic, taxonomic and environmental significance. *Palaeogeography, Palaeoclimatology, Palaeoecology* 23, pp. 199-229.
- FREY, R.W. and SEILACHER, A. 1980. Uniformity in marine invertebrate ichnology. *Lethaia* 13, pp. 183-207.
- FÜRSICH, F.T. 1974a. Corallian (Upper Jurassic) trace fossils from England and Normandy. *Stuttgarter Beiträge zur Naturkunde, Serie B (Geologie und Paläontologie)* 13, pp. 1-52.
- 1974b. On *Diplocraterion Torell* 1870 and the significance of morphological features in vertical, spreiten-bearing, U-shaped trace fossils. *Journal of Paleontology* 48, pp. 952-962.
- 1974c. Ichnogenus *Rhizocorallium*. *Paläontologische Zeitschrift* 48, pp. 16-28.
- 1975. Trace fossils as environmental indicators in the Corallian of England and Normandy. *Lethaia* 8, pp. 151-172.

- 1981. Invertebrate trace fossils from the Upper Jurassic of Portugal. *Comunicações dos Serviços Geológicos de Portugal* 67, pp. 153-168.
- FÜRSICH, F.T. and MAYR, H. 1981. Non-marine *Rhizocorallium* (trace fossil) from the Upper Freshwater Molasse (Upper Miocene) of southern Germany. *Neues Jahrbuch für Geologie und Paläontologie Monatshefte* 6, pp. 321-333.
- GARCIA-RAMOS, J.C.M. 1976. Morfologia de trazas fosiles en dos afloramientos de Arenisca de Naranco (Devonico medio) de Asturias (NW. de Espana). *Trabajos de Geologia* 8, pp. 131-171.
- GLAESSNER, M.F. 1969. Trace fossils from the Precambrian and basal Cambrian. *Lethaia* 2, pp. 369-393.
- GOLDRING, R. 1962. The trace fossils of the Baggy Beds (Upper Devonian) of North Devon, England. *Paläontologische Zeitschrift* 36, pp. 232-251.
- GUTSCHICK, R.C. and RODRIGUEZ, J. 1977. Late Devonian-Early Mississippian trace fossils and environments along the Cordilleran Miogeocline, western United States. In *Trace fossils 2*. Edited by T.P. Crimes and J.C. Harper. Geological Journal Special Issue 9. Seel House Press, Liverpool, pp. 195-208.
- HAKES, W.G. 1976. Trace fossils and depositional environment of four clastic units, Upper Pennsylvanian megacyclothems, northeast Kansas. *The University of Kansas Paleontological Contributions* 63, 46p.
- 1977. Trace fossils in Late Pennsylvanian cyclothems, Kansas. In *Trace fossils 2*. Edited by T.P. Crimes and J.C. Harper. Geological Journal Special Issue 9. Seel House Press, Liverpool, pp. 209-226.
- HALDEMAN, S.S. 1840. Supplement to number one of "A monograph of the Limniades, and other freshwater uivalve shells of North America", containing descriptions of apparently new animals in different classes, and the names and characters of the subgenera in *Paludina* and *Anculosa*. Philadelphia, 3p.
- HALL, J. 1847. *Palaeontology of New York*, Vol. 1. C. Van Benthuysen, Albany, 338p.
- 1852. *Palaeontology of New York*. Vol. 2. C. Van Benthuysen, Albany, 362p.
- HALLAM, A. 1975. Preservation of trace fossils. In *The Study of Trace Fossils*. Edited by R.W. Frey. Springer-Verlag, New York, pp. 56-63.
- HÄNTZSCHEL, W. 1962. Trace fossils and Problematica. In *Treatise on Invertebrate Paleontology*, Part W. Edited by R.C. Moore. Geological Society of America and University of Kansas Press, New York and Lawrence, Kansas, pp. W177-W245.
- 1975. Trace fossils and Problematica. In *Treatise on Invertebrate Paleontology*, Part W, *Miscellanea*, Supplement I. Edited by C. Teichert. Geological Society of America and The University of Kansas Press, Boulder, Colorado, Lawrence, Kansas, 269p.
- HÄNTZSCHEL, W. and REINECK, H.-E. 1968. Fazies-Untersuchungen im Hettangium von Helmstedt (Niedersachsen). *Geologische Staatsinstitut in Hamburg, Mitteilungen* 37, pp. 5-39.
- HARLAND, T.L. and PICKERILL, R.K. 1982. A review of Middle Ordovician sedimentation in the St. Lawrence Lowland, eastern Canada. *Geological Journal* 17, pp. 135-156.
- 1984. Ordovician rocky shoreline deposits - the basal Trenton Group around Quebec City, Canada. *Geological Journal* 19, (in press).
- HEER, O. 1877. *Flora Fossilis Helvetiae*. Die vorweltliche Flora der Schweiz. J. Würster & Company, Zürich, 182p.
- HELWIG, J. 1972. Stratigraphy, sedimentation, paleogeography and paleoclimates of Carboniferous ("Gondwana") and Permian of Bolivia. *Bulletin of the American Association of Petroleum Geologists* 56, pp. 1008-1033.
- HESTER, N.C. and PRYOR, W.A. 1972. Blade-shaped crustacean burrows of Eocene age: a composite form of *Ophiomorpha*. *Bulletin of the Geological Society of America* 83, pp. 677-688.
- HILL, G.W. 1981. Ichnocoenoses of a Paleocene submarine-canyon floor. In *Upper Cretaceous and Paleocene turbidites, Central California Coast*. Edited by V. Frizzell. Society of Economic Paleontologists and Mineralogists, Field Guide Book (Trip #6), pp. 93-104.
- HITCHCOCK, E. 1858. *Ichnology of New England*. A report on the sandstone of the Connecticut Valley, especially its footprints. W. White, Boston, 220p.
- HOFMANN, H.J. 1979. Chazy (Middle Ordovician) trace fossils in the Ottawa-St. Lawrence Lowlands. *Bulletin of the Geological Survey of Canada* 321, pp. 27-59.
- HOWARD, J.D. 1966. Characteristic trace fossils in Upper Cretaceous sandstones of the Book Cliffs and Wasatch Plateau. *Utah Geological and Mineralogical Survey, Central Utah Coal Bulletin* 80, pp. 35-53.
- HOWELL, B.F. 1943. Burrows of *Skolithos* and *Planolites* in the Cambrian Hardyston Sandstone at Reading, Pennsylvania. *Bulletin of the Wagner Free Institute of Science of Philadelphia* 3, pp. 1-33.
- IMLAY, R.M. 1961. Characteristic Lower Cretaceous megafossils from northern Alaska. *United States Geological Survey, Professional Paper* 335, 74p.
- JAEGER, H. and MARTINSSON, A. 1980. The Early Cambrian trace fossil *Plagiogmus* in its type area. *Geologiska Föreningens i Stockholm Förhandlingar* 102, pp. 117-126.
- JAMES, J.F. 1891. Studies in problematic organisms - the genus *Scolithus*. *Bulletin of the Geological Society of America* 3, pp. 32-44. (Part of this paper also appears in *American Naturalist* 26, pp. 240-242, 1892).
- JAMES, N.P., KOBLUK, D.R. and PEMBERTON, S.G. 1977. The oldest macroborers: Lower Cambrian of Labrador. *Science* 197, pp. 980-983.
- JORDAN, T.E. 1981. Enigmatic deep-water mechanisms, upper part of the Oquirrh Group, Utah. *Journal of Sedimentary Petrology* 51, pp. 879-894.
- KARASZEWSKI, W. 1971. Some fossil traces from the lower Liassic of the Holy Cross Mts., Central Poland. *Bulletin de l'Académie Polonaise des Sciences, Série des Sciences de la Terre* 19, pp. 101-105.
- KENNEDY, W.J. 1970. Trace fossils in the Chalk environment. In *Trace fossils*. Edited by T.P. Crimes and J.C. Harper. Geological Journal Special Issue 3. Seel House Press, Liverpool, pp. 263-282.
- 1975. Trace fossils in carbonate rocks. In *The Study of Trace Fossils*. Edited by R.W. Frey. Springer-Verlag, New York, pp. 377-398.
- 1984. Ordovician rocky

- KOBLUK, D.R., JAMES, N.P. and PEMBERTON, S.G. 1978. Initial diversification of macroboring ichnofossils and exploitation of the macroboring niche in the Lower Paleozoic. *Paleobiology* 4, pp. 163-170.
- KOBLUK, D.R. and NEMCSOK, S. 1982. The macroboring ichnofossil *Trypanites* in colonies of the Middle Ordovician bryozoan *Prasopora*: population behaviour and reaction to environmental influences. *Canadian Journal of Earth Sciences* 19, pp. 679-688.
- KSIĄŻKIEWICZ, M. 1970. Observations on the ichnofauna of the Polish Carpathians. In *Trace fossils*. Edited by T.P. Crimes and J.C. Harper. *Geological Journal Special Issue 3*. Seel House Press, Liverpool, pp. 283-322.
- 1977. Trace fossils in the flysch of the Polish Carpathians. *Palaeontologica Polonica* 36, pp. 1-208.
- LARSON, J.A. 1982. Redeposited Carbonates of the Upper Oquirrh Formation, Utah. *Brigham Young University, Geology Studies* 29, pp. 65-84.
- LESQUEREUX, L. 1876. Species of fossil marine plants from the Carboniferous measures. *Geological Survey of Indiana, Annual Report* 7, pp. 134-145.
- MÄGDEFRAU, K. 1932. Über einige Bohrgänge aus dem unteren Muschelkalk von Jena. *Paläontologische Zeitschrift* 14, pp. 150-160.
- MAILLARD, G. 1887. Considérations sur les fossiles décrits comme algues. *Société Paléontologique de la Suisse, Mémoire* 14, pp. 1-40.
- MARINTSCH, E.J. and FINKS, R.M. 1982. Lower Devonian ichnofacies at Highland Mills, New York and their gradual replacement across environmental gradients. *Journal of Paleontology* 56, pp. 1050-1078.
- MARTINSSON, A. 1965. Aspects of a Middle Cambrian thanatotope on Oland. *Geologiska Föreningens i Stockholm, Förhandlingar* 62, pp. 181-230.
- MASSALONGO, A. 1855. *Zoophycos, novum genus plantarum fossilium*. Antonelli (Verona), 52p.
- MAYER, G. 1954. Ein neues *Rhizocorallium* aus dem mittleren Hauptmuschelkalk von Bruchsal. *Beiträge zur Naturkundlichen Forschung in Südwestdeutschland* 13, pp. 80-83.
- MCCARTHY, B. 1979. Trace fossils from a Permian shoreface-foreshore environment, eastern Australia. *Journal of Paleontology* 53, pp. 345-366.
- MILLER, S.A. 1889. *North American geology and palaeontology for the use of amateurs, students and scientists*. Western Methodist Book Concern, (Cincinnati, Ohio), 664p.
- MILLER, S.A. and DYER, C.B. 1878. *Contributions to paleontology*, 2, Privately published (Cincinnati, Ohio), 1lp.
- MYANNIL, R.M. 1966. O Vertikalnykh norkakh zaryvaniya v Ordovikskikh izvestiyakh Pribaltiki. In *Organizm i sreda v geologicheskoy proshlom*, Akademiya Nauk SSSR Paleontologicheskii Institut, pp. 200-207.
- NARBONNE, G.M., GIBLING, M.R. and JONES, B. 1979. *Polarichnus garnienensis*, new genus and new species trace fossil from Siluro-Devonian strata of Arctic Canada. *Journal of Paleontology* 53, pp. 133-141.
- NATHORST, A.G. 1897. Ett märkligt spar från Tesinisandstenen på Öland. *Geologiska Föreningens i Stockholm Förhandlingar* 19, pp. 361-365.
- NICHOLSON, H.A. 1873. Contributions to the study of the errant annelides of the Older Palaeozoic rocks. *Proceedings of the Royal Society of London* 21, pp. 288-290 (also *Geological Magazine* 10, pp. 309-310).
- 1978. Contributions to the study of errant annelides of the older Palaeozoic rocks. Publication of the Department of Geology and Mineralogy, University of Aberdeen 1, pp. 17-47 (posthumus).
- NICHOLSON, H.A. and HINDE, G.J. 1875. Notes on the fossils of the Clinton, Niagara and Guelph formations of Ontario, with descriptions of new species. *Canadian Journal of Science, Literature and History, new series* 14, pp. 137-160.
- ORBIGNY, A. d' 1842. *Voyage dans l'Amérique méridionale (le Brésil, la République orientale de l'Uruguay, la République Argentine, la Patagonie, la République du Chili, la République de Bolivie, la République de Pérou) exécuté pendant les années 1826, 1827, 1829, 1830, 1831, 1832, et 1833*, v. 3, pt. 4 (*Paléontologie*), 188p.
- OSGOOD, R.G. Jr. 1970. Trace fossils of the Cincinnati area. *Paleontographica Americana* 6, pp. 281-444.
- 1975. The paleontological significance of trace fossils. In *The Study of Trace Fossils*. Edited by R.W. Frey. Springer-Verlag, New York, pp. 87-108.
- OSGOOD, R.G. Jr. and DRENNEN, W.T. III 1975. Trilobite trace fossils from the Clinton Group (Silurian) of east-central New York State. *Bulletin of American Paleontology* 287, pp. 299-348.
- OSGOOD, R.G. Jr., DRENNEN, W.T. III and SZMUC, E.G. 1972. The trace fossil *Zoophycos* as an indicator of water depth. *Bulletin of American Paleontology* 62, pp. 1-22.
- OTTO, E. von. 1854. *Additamente zur Flora des Quadergeirges in Sachsen*. G. Mayer, Leipzig, Part 2, 53p.
- PEMBERTON, S.G., FLACH, P.D. and MOSSOP, G.D. 1982. Trace Fossils from the Athabasca Oil Sands, Alberta, Canada. *Science* 217, pp. 825-827.
- PEMBERTON, S.G. and FREY, R.W. 1982. Trace fossil nomenclature and the *Planolites*-*Palaeophycus* dilemma. *Journal of Paleontology* 56, pp. 843-881.
- PEMBERTON, S.G., KOBLUK, D.R., YEO, R.K. and RISK, M.J. 1980. The boring *Trypanites* at the Silurian-Devonian unconformity in southern Ontario. *Journal of Paleontology* 54, pp. 1258-1266.
- PFEFFERKORN, H.W. 1971. Note on *Conostichus broadheady* Lesquereux (trace fossil, pp. Pennsylvania). *Journal of Paleontology* 45, pp. 888-892.
- PICKERILL, R.K. 1976. *Vermiforichnus* borings from the Ordovician of central Wales. *Geological Magazine* 113, pp. 154-164.
- 1977. Trace fossils from the Upper Ordovician (Caradoc) of the Berwyn Hills, central Wales. *Geological Journal* 12, pp. 1-16.
- 1980. Phanerozoic flysch trace fossil diversity-observations based on an Ordovician flysch ichnofauna from the Aroostook-Matapedia Carbonate Belt of northern New Brunswick. *Canadian Journal of Earth Sciences* 17, pp. 1259-1270.
- 1981. Trace fossils in a Lower Palaeozoic submarine canyon sequence - the Siegas Formation of northwestern New Brunswick, Canada. *Maritime Sediments and Atlantic Geology* 17, pp. 37-58.
- PICKERILL, R.K., FILLION, D. and HARLAND, T.L. 1984. Middle Ordovician trace fossils in carbonates of the Trenton Group between Quebec City and

- Montreal, St. Lawrence Lowland, eastern Canada. *Journal of Paleontology* 58, (in press).
- PICKERILL, R.K. and FORBES, W.H. 1979. Ichnology of the Trenton Group in the Quebec City area. *Canadian Journal of Earth Sciences* 16, pp. 2022-2039.
- PICKERILL, R.K., HURST, J.M. and SURLYK, F. 1982. Notes on Lower Palaeozoic flysch trace fossils from Hall Land and Peary Land, North Greenland. *Grønlands Geologiske Undersøgelse* 108, pp. 25-29.
- PICKERILL, R.K. and KEPPIE, J.D. 1981. Observations on the ichnology of the Meguma Group (?Cambro-Ordovician) of Nova Scotia. *Maritime Sediments and Atlantic Geology* 17, pp. 130-138.
- PLICKA, M. 1968. Zoophycos, and a proposed classification of sabellid worms. *Journal of Paleontology* 42, pp. 836-849.
- 1970. Zoophycos and similar fossils. In *Trace fossils*. Edited by T.P. Crimes and J.C. Harper. Geological Journal Special Issue 3. Seel House Press, Liverpool, pp. 361-370.
- PRANTL, F. 1945. Two new problematic trails from the Ordovician of Bohemia. *Académie Tchèque des Sciences, Bulletin International, Classe des Sciences Mathématiques, Naturelles et de la Médecine* 46, pp. 49-59.
- QUATREFAGES, M.A. de 1849. Note sur la *Scolicia prisca* (A. de Q.), annélide fossile de la Craie. *Annales des Sciences Naturelles, série 3, Zoologie* 12, pp. 265-266.
- QUENSTEDT, F.A. 1849. *Petrefactenkunde Deutschlands*. I. Abth. v. 1, pp. Cephalopoden. L.F. Fues, Tübingen, 580p.
- RATCLIFFE, B.C. and FAGERSTROM, J.A. 1980. Invertebrate lebensspuren of Holocene floodplains: their morphology, origin and palaeoecological significance. *Journal of Paleontology* 54, pp. 614-630.
- RICHTER, R. 1920. Ein devonischer "Pfeifenquarzit" verglichen mit der heutigen "Sandkoralle" (*Sabellaria*, Annelidae). *Senckenbergiana* 2, pp. 215-235.
- 1921. *Scolithus*, *Sabellarifex* und *Geflecht-quarzite*. *Senckenbergiana* 3, pp. 49-52.
- 1924. *Flachseebeobachtungen zur Paläontologie und Geologie VII-XI*. *Senckenbergiana* 6, pp. 119-165.
- 1937. *Marken und Spuren aus allen Zeiten I-II*. *Senckenbergiana* 19, pp. 150-169.
- RIVA, J. 1968. Graptolite faunas from the Middle Ordovician of the Gaspé north shore. *Naturaliste Canadien* 95, pp. 1379-1400.
- 1969. Middle and Upper Ordovician graptolite faunas of the St. Lawrence Lowland of Quebec and of Anticosti Island. In *North Atlantic - geology and continental drift*. Edited by M. Kay. American Association of Petroleum Geologists, Memoir 12, pp. 513-556.
- 1972. *Geology of the environs of Quebec City*. 24th International Geological Congress, Guidebook for Excursion B19, 53p.
- ROEDEL, H. 1929. Ergänzung zu meiner Mitteilung über ein kambrisches Geschiebe mit problematischen Spuren. *Zeitschrift für Geschiebeforschung* 5, pp. 48-51.
- RONIEWICZ, P. and PIENKOWSKI, G. 1977. Trace fossils of the Podhale Flysch Basin. In *Trace fossils 2*. Edited by T.P. Crimes and J.C. Harper. Geological Journal Special Issue 9. Seel House Press, Liverpool, pp. 273-288.
- SACCO, R. 1888. *Note di Paleoichnologia Italiana*. Società Italiana di Scienze Naturali e del Museo Civile di Storia Naturale, Atti 31, pp. 151-192.
- SALTER, J.W. 1857. On annelide-burrows and surface markings from the Cambrian rocks of the Longmynd. *Quarterly Journal of the Geological Society of London* 13, pp. 199-206.
- SAPORTA, G. MARQUIS de 1872-1873. *Paléontologie française ou description des fossiles de la France, série 2, Végétaux*. Plantes Jurassiques, Volume 1. G. Masson, Paris, 506p.
- SCHINDEWOLF, O.H. 1921. *Studien aus dem Marburger Buntsandstein I, II*. *Senckenbergiana* 3, pp. 33-49.
- 1928. *Studien aus dem Marburger Buntsandstein III-VI*. *IV. Isopodichnus problematicus* (Schindewolf) im Unteren und Mittleren Buntsandstein. *Senckenbergiana* 10, pp. 27-37.
- SEILACHER, A. 1955. *Spuren und Fazies in Unterkambrium*. In *Beiträge zur Kenntnis des Kambriums, in der Salt Range (Pakistan)*. Edited by O.H. Schindewolf and Adolf Seilacher. Akademie der Wissenschaften und der Literatur zu Mainz, mathematisch-naturwissenschaftliche Klasse, Abhandlungen 10, pp. 253-256.
- 1964. Sedimentological classification and nomenclature of trace fossils. *Sedimentology* 3, pp. 253-256.
- 1967. Bathymetry of trace fossils. *Marine Geology* 5, pp. 189-200.
- 1970. *Cruziana stratigraphy of "non-fossiliferous" Palaeozoic sandstones*. In *Trace fossils*. Edited by T.P. Crimes and J.C. Harper. Geological Journal Special Issue 3. Seel House Press, Liverpool, pp. 447-476.
- SEILACHER, A. and MEISCHNER, D. 1965. *Fazies-Analyse im Palaeozoikum des Oslo-Gebietes*. *Geologische Rundschau* 54, pp. 596-619.
- SELLWOOD, B.W. 1970. The relation of trace fossils to small scale sedimentary cycles in the British Lias. In *Trace fossils*. Edited by T.P. Crimes and J.C. Harper. Geological Journal Special Issue 3. Seel House Press, Liverpool, pp. 489-504.
- SHONE, R.W. 1978. Giant *Cruziana* from the Beaufort Group. *Transactions of the Geological Society of South Africa* 81, pp. 327-329.
- 1979. "Giant *Cruziana* from the Beaufort Group". *Transactions of the Geological Society of South Africa* 82, pp. 371-375.
- SIMPSON, S. 1957. On the trace fossil *Chondrites*. *Quarterly Journal of the Geological Society of London* 112, pp. 475-499.
- 1970. Notes on *Zoophycos* and *Spirophyton*. In *Trace fossils*. Edited by T.P. Crimes and J.C. Harper. Geological Journal Special Issue 3. Seel House Press, Liverpool, pp. 505-514.
- SMITH, A.B. and CRIMES, T.P. 1984. Trace fossils formed by heart urchins - a study of *Scolicia* and related traces. *Lethaia* 16, pp. 79-92.
- SOWERBY, J. 1829. *The Mineral Conchology of Great Britain*, Volume 6. London, 230p.
- STERNBERG, K.M. Graf. von 1833. *Versuch einer geognostisch-botanischen Darstellung der Flora der Vorwelt* 5, 6, pp. 1-80.
- SWINBANKS, D.D. and MURRAY, J.W. 1981. *Biosedimentological zonation of Boundary Bay tidal flats, Fraser River Delta, British Columbia*. *Sedimentology* 28, pp. 201-237.
- TAYLOR, B.J. 1967. Trace fossils from the Fossil Bluff Series of Alexander Island. *Bulletin of the British Antarctic Survey* 13, pp. 1-30.

- TREWIN, N.H. 1976. *Isopodichnus* in a trace fossil assemblage from the Old Red Sandstone. *Lethaia* 9, pp. 29-38.
- ULRICH, E.O. 1904. Fossils and age of the Yakutat Formation. Descriptions of collections made chiefly near Kadiak, Alaska. Harriman Alaska Expedition, 4 (Geology and Paleontology), pp. 125-146.
- VANUXEM, L. 1842. Geology of New York, pt. III, comprising the survey of the 3d geological district. W. and A. White and J. Visscher, Albany, 306p.
- VIALOV, O.S. 1971. Redkie problematiki iz mezozoya Pamira i Kavkaza. *Paleontologicheskii Sbornik*, 7, pp. 85-93.
- VOIGT, E. and SOULE, J.D. 1973. Cretaceous burrowing bryozoans. *Journal of Paleontology* 47, pp. 21-33.
- WARD, D.M. and LEWIS, D.W. 1975. Paleoenvironmental implications of storm-scoured, ichnofossiliferous mid-Tertiary limestones, Waihao District, South Canterbury, New Zealand. *New Zealand Journal of Geology and Geophysics* 18, pp. 881-908.
- WARME, J.E. 1975. Borings as trace fossils, and the processes of marine bioerosion. In *The Study of Trace Fossils*. Edited by R.W. Frey. Springer-Verlag, New York, pp. 181-227.
- WEBBY, B.D. 1969. Trace fossils (*Pascichnia*) from the Silurian of New South Wales, Australia. *Paläontologische Zeitschrift* 43, pp. 81-94.
- 1970. Late Precambrian trace fossils from New South Wales. *Lethaia* 3, pp. 79-109.
- WELLER, S. 1899. Kinderhook faunal studies I. The fauna of the vermicular sandstone at Northview, Webster County, Missouri. *Transactions of the Academy of Science of St. Louis* 9, pp. 9-51.
- WETZEL, A. 1981. Ökologische und stratigraphische Bedeutung biogener Gefüge in quartären Sedimenten am NW - afrikanischen Kontinentalrand. "Meteor" Forschungsergebnisse, Reihe C 34, pp. 1-47.
- WETZEL, A. and WERNER, R. 1981. Morphology and ecological significance of *Zoophycos* in deep-sea sediments off NW Africa. *Palaeogeography, Palaeoclimatology, Palaeoecology* 32, pp. 185-212.
- WHITAKER, J.H.M. 1979. A new trace fossil from the Ringerike Group, Southern Norway. *Proceedings of the Geological Association of London* 91, pp. 85-89.
- WILSON, A.E. 1948. Miscellaneous classes of fossils, Ottawa Formation, Ottawa-St. Lawrence Valley. *Bulletin of the Geological Survey of Canada, Department of Mines and Resources* 11, 116p.
- ZENKER, J.C. 1836. *Historisch-topographisches Taschenbuch von Jena und seiner Umgebung besonders in naturwissenschaftlicher und medicinischer Beziehung*. J.C. Zenker (ed.), Wackenholder (Jena), 338p.

Reviewers: T.P. Crimes
T.L. Harland