

Superposed Late Paleozoic thermal events in the southwestern Meguma Terrane, Nova Scotia

R.D. Dallmeyer

*Department of Geology, University of Georgia
Athens, Georgia 30602 U.S.A.*

and

J.D. Keppie

*Department of Mines and Energy, P.O. Box 1087
Halifax, Nova Scotia B3J 2X1*

Date Received December 5, 1987

Date Accepted August 15, 1988

Cambrian-Devonian sedimentary sequences within southwestern portions of the Meguma Terrane were deformed with attendant development of a regionally penetrative cleavage during low-grade regional metamorphism at *ca.* 400-410 Ma. Contact aureoles developed during subsequent emplacement of the Port Mouton, Shelburne and Barrington Passage plutons at *ca.* 370-375 Ma. Amphibole within calc-silicate horizons in the contact zones record internally discordant $^{40}\text{Ar}/^{39}\text{Ar}$ age spectra. However, $^{36}\text{Ar}/^{40}\text{Ar}$ vs. $^{39}\text{Ar}/^{40}\text{Ar}$ isotope correlations yield ordinate intercept values which approximate atmospheric values, and abscissa ratios corresponding to ages ranging between 385.1 ± 2.6 Ma and 366.9 ± 5.6 Ma. These are interpreted to date post-contact metamorphic cooling through temperatures required for intracrystalline retention of argon. Biotite and muscovite within northern portions of the contact zones record 340-350 Ma $^{40}\text{Ar}/^{39}\text{Ar}$ plateau ages which date later post-metamorphic cooling through the lower temperatures required for mica argon closure. Intracrystalline argon systems within biotite and muscovite in southern portions of the contact zone of the Shelburne and Barrington Passage plutons were variably rejuvenated during a later reheating associated with emplacement of plutons at *ca.* 315 Ma (Wedgeport Pluton) and 290 Ma (offshore, subsurface pluton). Temperatures attained during these later thermal events were not sufficient to rejuvenate hornblende argon systems.

Dans les portions sub-ouest de la Lanrière de Méguma, les séries sédimentaires cambro-dévonniennes ont subi, il y a environ 400-410 Ma, une déformation qui s'accompagna du développement concomitant d'une schistosité pénétrative (à l'échelle régionale) au cours d'un métamorphisme régional de faible intensité. L'emplacement subséquent des plutons de Port Mouton, Shelburne et Barrington Passage se manifesta par l'implantation d'aureoles de métamorphisme vers 370-375 Ma. L'amphibole provenant de niveaux calco-silicatés, au sein des zones de contact, présente des patrons d'âge $^{40}\text{Ar}/^{39}\text{Ar}$ marqués par une discordance interne. En revanche, les courbes de corrélation des isotopes $^{36}\text{Ar}/^{40}\text{Ar}$ vs. $^{39}\text{Ar}/^{40}\text{Ar}$ montrent des ordonnées à l'origine se rapprochant des valeurs de l'atmosphère ainsi que des rapports en abscisse correspondant à des âges qui s'étendent de $385,1 \pm 2,6$ Ma à $366,9 \pm 5,6$ Ma. Ces âges dateraient, par le biais des températures nécessaires à la rétention intracrystalline de l'argon, un refroidissement post-métamorphisme de contact. La biotite et la muscovite provenant des portions septentrionales des zones de contact ont fixé des âges limites $^{40}\text{Ar}/^{39}\text{Ar}$ de 340-350 Ma qui datent un refroidissement post-métamorphe ultérieur à l'aide des températures plus basses régissant la fermeture de l'argon dans les micas. L'emplacement par la suite d'autres plutons aux alentours de 315 Ma (Pluton de Wedgeport) et 290 Ma (pluton de sous-surface en offshore) engendra la réjuvenation à divers degrés des systèmes d'argon intra-cristallin chez la biotite et la muscovite provenant des secteurs méridionaux de la zone de contact des plutons de Shelburne et Barrington Passage. Les températures atteintes lors de ces derniers épisodes thermiques restèrent en deçà de celles auxquelles fait appel la réjuvenation des systèmes d'argon de la hornblende.

[Traduit par le journal]

INTRODUCTION

The Meguma Terrane is characterized by an extensive Cambrian-Ordovician turbidite sequence which is conformably overlain by Silurian-Devonian shallow marine and continental sedimentary rocks. Following a regional, generally low-grade metamorphism (M1) and concomitant folding (D1), the metasedi-

mentary sequence was intruded by numerous generally granitic plutons. Dallmeyer and Keppie (1987) presented reconnaissance $^{40}\text{Ar}/^{39}\text{Ar}$ incremental-release mineral ages (mostly for muscovite and biotite) for several granitic stocks and host metasedimentary units within southwestern portions of the Meguma Terrane. These suggested a complex late Paleozoic tectonothermal evolution which included regional D₁ folding and associated

formation of cleavage during greenschist-lower amphibolite facies metamorphism (M_1) at ca. 400-410 Ma. Subsequent intrusion of granitic stocks occurred over a prolonged interval between ca. 375 and 285 Ma. The plutons were emplaced at shallow to intermediate crustal depths and developed contact metamorphic aureoles of variable size and grade. These overprinted M_1 regional metamorphic assemblages and are locally mutually superposed. The regional extent and relative intensities of individual thermal events affecting southwestern portions of the Meguma Terrane were not clearly defined by the reconnaissance mica ages reported by Dallmeyer and Keppie (1988). In an attempt to more clearly resolve thermal overprinting relationships, samples of amphibole-bearing, calc-silicate horizons were collected within the southwestern Meguma Terrane for $^{40}\text{Ar}/^{39}\text{Ar}$ analysis. These results are presented here and when combined with U-Pb ages reported by Keppie and Krogh (1987) for crystallization of zircon within two large plutons in the area (Shelburne and Barrington Passage plutons) allow a much clearer resolution of the regional tectonothermal evolution than was previously possible.

REGIONAL GEOLOGIC SETTING

The Meguma Group and overlying Silurian-Devonian formations were deformed into upright, subhorizontal D_1 folds which display a variably developed, axial planar, spaced cleavage. In pelitic horizons the cleavage is generally penetrative, and defined, in part, by the preferred orientation of chlorite, muscovite and/or biotite (Taylor and Schiller, 1966; Keppie, 1976). This suggests that a regional low-grade metamorphism (M_1) accompanied D_1 deformation. Numerous plutons cross-cut D_1 fabric elements. These are predominantly granodiorite and granite with subordinate quartz diorite and alaskite porphyry.

The plutons typically are surrounded by low-pressure contact metamorphic aureoles characterized by development of porphyroblastic chlorite, biotite, cordierite, staurolite, andalusite and/or sillimanite (Taylor and Schiller, 1966; Keppie and Meucke, 1979) which typically overgrow D_1 fabric elements (Sage, 1984; Keppie *et al.*, 1985). This contact metamorphism is widespread in southwestern Nova Scotia (Fig. 1) and was initially interpreted as a regional metamorphic event (e.g., Taylor and Schiller, 1966). However, recent detailed mapping (Sage, 1984; White, 1984; Bourque, 1985; Ross, 1985; Wentzell, 1985; Keppie *et al.*, 1985; Raeside *et al.*, 1985; Misner, 1986) suggests that the isograds are spatially related to exposed pluton contacts. Superposition of contact aureoles locally has produced complex textural relationships and zoned porphyroblasts; however, no maps presently exist showing these intersecting isograds. Therefore, the isograds shown in Figures 1 and 2 reflect cumulative peak metamorphic assemblages. Detailed mapping of the isograds throughout poorly exposed interior portions of the Meguma Terrane has not been attempted. In these areas only the reconnaissance isograds defined by Keppie and Muecke (1979) are available.

Variably penetrative ductile shear zones are locally developed throughout the Meguma Terrane. These affect both granitic plutons and host metasedimentary units. The shear zones generally dip steeply and trend parallel to regional strike. Preliminary analysis of associated fabric elements suggests that most formed during dextral shear (Keppie *et al.*, 1985). While a portion of this strain occurred in the late Devonian (Dallmeyer and Keppie, 1987), some is also recorded in the 315 ± 3 Ma Wedgeport pluton (Keppie *et al.*, 1983). In places the ductile shear zone fabrics are overgrown by undeformed contact metamorphic porphyroblasts of muscovite and/or biotite while elsewhere micas are dynamically recrystallized.




Fig. 1. Simplified geological map of the Meguma Terrane, Nova Scotia. Area of Figure 2 is outlined. Tick-marks on high-grade side of isograds. Individual plutons are named. Area of Figure 2 is indicated.

PREVIOUS GEOCHRONOLOGY

Isotopic ages previously reported for the Meguma Terrane have been summarized by Dallmeyer and Keppie (1987). In addition, Keppie and Krogh (1988) reported *ca.* 370-375 Ma U-

Pb zircon crystallization ages for the Shelburne and Barrington Passage plutons. The ⁴⁰Ar/³⁹Ar mineral ages presented by Dallmeyer and Keppie (1987) from the present study area are located in Figure 2.




Fig. 2. Location of the three exposures (circled numbers) investigated in this study (refer to Fig. 1 for area and map explanation) compared with the ⁴⁰Ar/³⁹Ar mineral ages presented by Dallmeyer and Keppie (1987): plateau ages shown for hornblende (H), biotite (B), and/or muscovite (M) concentrates recording internally concordant age spectra; all discordant spectra are shown (abscissa and ordinate coordinates similar for all samples). Analytical uncertainties (two sigma, intralaboratory) are represented by vertical width of bars. Experimental temperatures increase from left to right. Plateau and/or total-gas ages listed on each spectrum. Isograd ornamentation identical to that in Figure 1 (gar = garnet, and = andalusite, sill = sillimanite). Shear zones are indicated.

ANALYTICAL METHODS

$^{40}\text{Ar}/^{39}\text{Ar}$

The techniques used during $^{40}\text{Ar}/^{39}\text{Ar}$ analyses of the Meguma samples generally followed those described in detail by Dallmeyer and Keppie (1987). The mineral concentrates were wrapped in aluminum-foil packets, encapsulated in sealed quartz vials, and irradiated for 40 hr at 1000 Kw in the central thimble position of the U.S. Geological Survey TRIGA reactor in Denver, Colorado. Variations in the flux of neutrons along the length of the irradiation assembly were monitored with several mineral standards, including MMhb-1 (Alexander *et al.*, 1978). The samples were incrementally heated until fusion with an RF generator. Each heating step was maintained for 30 minutes. Operational system blanks were routinely measured. Although these were consistently characterized by an atmospheric argon isotopic composition, blank volumes were variable ($0.8\text{--}6.2 \times 10^{-12}$ STP cm^3 for ^{36}Ar) and demonstrated a complex dependence on: (1) crucible temperature; (2) history of the furnace following sample loading; and (3) samples characteristics (weight, material, grain size, density). In view of this uncertainty and because blanks generally represented less than *ca.* 5% of the total measured ^{40}Ar signal, blank corrections have not been applied to the analytical data. Measured isotopic ratios were corrected for the effects of mass discrimination and interfering isotopes produced during irradiation using factors reported by Dalrymple *et al.* (1981) for the reactor used in the present study. Apparent $^{40}\text{Ar}/^{39}\text{Ar}$ ages were calculated from the corrected isotopic ratios using the decay constants and isotopic abundance ratios listed by Steiger and Jäger (1977).

Two categories of uncertainties are encountered in $^{40}\text{Ar}/^{39}\text{Ar}$ incremental-release dating. One group involves intralaboratory uncertainties related to measurement of the isotopic ratios used in the age equation. The other group considers interlaboratory uncertainties in the other parameters used in the age equation (monitor age, J-value determination, etc.), and are the same for each gas increment evolved from a particular sample. Therefore, to evaluate the significance of potential incremental age variations within a sample, only intralaboratory uncertainties should be considered. These are reported here for the incremental gas ages. They have been calculated by statistical propagation of uncertainties associated with measurement of each isotopic ratio (at two standard deviations of the mean) through the age equation. Intralaboratory uncertainties are *ca.* $\pm 1.25\text{--}1.5\%$ of the quoted age. Total-gas ages have been computed for each sample by appropriated weighting of the age and percent ^{39}Ar released within each temperature increment. Analyses of the MMhb-1 monitor indicate that apparent K/Ca ratios may be calculated through the relationship $0.518 (\pm 0.0005) \times 3^9\text{Ar}/^{37}\text{Ar}$ corrected.

The analyses of the amphibole concentrates have been plotted on $^{36}\text{Ar}/^{40}\text{Ar}$ vs. $^{39}\text{Ar}/^{40}\text{Ar}$ isotope correlation diagrams (Roddick *et al.*, 1980; Radicati de Brozolo *et al.*, 1981). Regression techniques followed the methods of York (1969). The mean square of the weighted deviates (MSWD) has been used to evaluate isotopic correlations. Roddick (1978) suggests that an MSWD in excess of *ca.* 2.5 indicates scatter about a correlation

line greater than that which can be explained only by experimental errors.

Electron microprobe

The techniques used during electron microprobe analyses of the amphibole concentrates generally followed those described in detail by Cho (1985). Grains were analyzed in carbon-coated, polished epoxy mounts with a JEOL JXA-733 electron microprobe using an accelerating potential of 15 kV and a beam current of 1×10^{-8} amp (on faraday cage). Na-Fe-K, Si-Mn-Ca, Al-Ti, or Mg were analyzed simultaneously. Ten five-second integrations were performed for each spot analyzed. A $10\ \mu\text{m}$ beam diameter was maintained. Following dead-time corrections, oxide percentages were calculated using a modification of the Bence-Albee method (JEOL program).

RESULTS

Samples of calc-silicate (A) and interlayered metapelitic schist (B) were collected at three locations (1-3) within the Goldenville Formation adjacent to exposed plutons in the southwestern Meguma Terrane. Sample locations are indicated in Figure 2. Locations 1 and 2 occur within the andalusite-staurolite-cordierite metamorphic zone whereas location 3 is situated in the sillimanite zone. Concentrates of amphibole, muscovite, and/or biotite have been analyzed from these samples. The analytical data are listed in Tables 2 and 3, and are portrayed as incremental-release age spectra in Figures 3 and 5. Dallmeyer and Keppie (1987) reported $^{40}\text{Ar}/^{39}\text{Ar}$ ages for concentrates of muscovite and biotite from schist interlayered with calc-silicate at location 1 (numbered 15 on their Fig. 3).

Amphibole

The calc-silicate horizons occur as thin ($<0.5\ \text{m}$), discontinuous, elliptical lithologic intervals with granoblastic textures. They are composed predominantly of amphibole, plagioclase and quartz with subordinate garnet. The amphibole crystals display no petrographic evidence of exsolution or zoning. Electron microprobe analyses of representative constituent grains within the concentrates prepared for $^{40}\text{Ar}/^{39}\text{Ar}$ analysis (Table 1) indicate that the concentrates are compositionally uniform. Following the classifications of Leake (1978) and Hawthorne (1982), those from samples 1A and 2A range between ferro-hornblende and ferro-actinolitic hornblende. The concentrate from sample 3A is slightly more magnesian, with compositions ranging between magnesio-hornblende and actinolitic hornblende.

The amphibole concentrate from sample 1A displays an internally discordant $^{40}\text{Ar}/^{39}\text{Ar}$ age spectrum corresponding to a total-gas age of 392.6 ± 4.5 Ma (Fig. 3). The apparent K/Ca spectrum shows little variation suggesting that the concentrate is characterized by relatively homogenous grains. The remaining 11 increments ($625\text{--}925^\circ\text{C}$) constitute *ca.* 94% of the total gas evolved from the concentrate and generally contain $>60\%$ radiogenic ^{40}Ar . These increments define a $^{36}\text{Ar}/^{40}\text{Ar}$ vs. $^{39}\text{Ar}/^{40}\text{Ar}$ correlation line (MSWD = 2.9) with intercepts corresponding to

Table 1. Representative electron microprobe analyses of constituent grains within amphibole concentrates from calc-silicate horizons in the Goldenville Formation, southwestern Meguma Terrane.

	Sample 1A								
	GRAIN 1			GRAIN 2			GRAIN 3		
	1+	2	3	1	2	3	1	2	3
SiO ₂	49.60	49.31	48.82	50.24	47.22	46.73	46.41	46.33	44.39
TiO ₂	0.20	0.25	0.27	0.18	0.47	0.40	0.49	0.28	0.42
Al ₂ O ₃	4.98	5.24	5.86	4.62	8.78	8.90	9.82	10.01	12.54
FeO*	19.38	19.70	19.72	18.59	19.01	19.84	17.98	18.38	18.69
MnO	0.65	0.60	0.59	0.61	0.56	0.53	0.45	0.62	0.47
MgO	9.96	10.13	9.34	10.77	8.93	8.92	9.37	9.34	8.06
CaO	12.35	11.75	12.06	12.09	11.97	11.99	12.03	11.84	11.80
Na ₂ O	0.24	0.28	0.27	0.20	0.48	0.42	0.53	0.65	0.63
K ₂ O	0.06	0.07	0.04	0.05	0.09	0.11	0.07	0.09	0.17
Total	97.42	97.33	96.97	97.35	97.51	97.84	97.15	97.54	97.17
Numbers of Ions on the Basis of 23(O)									
Si	7.436	7.400	7.361	7.493	7.066	7.004	6.942	6.921	6.673
Al ^{IV}	0.564	0.600	0.639	0.507	0.934	0.996	1.058	1.079	1.327
Al ^{VI}	0.316	0.327	0.403	0.305	0.615	0.576	0.673	0.683	0.895
Ti	0.023	0.028	0.031	0.020	0.053	0.045	0.055	0.031	0.047
Fe ⁺²	2.430	2.473	2.487	2.319	2.379	2.487	2.249	2.296	2.350
Mn	0.083	0.076	0.075	0.077	0.071	0.067	0.057	0.078	0.060
Mg	2.226	2.266	2.099	2.394	1.992	1.993	2.089	2.080	1.806
XM 1-3	0.078	0.170	0.095	0.115	0.110	0.168	0.123	0.168	0.158
Ca	1.984	1.890	1.949	1.932	1.919	1.926	1.928	1.895	1.901
	2.062	2.060	2.044	2.047	2.029	2.094	2.051	2.063	2.059
N	0.070	0.081	0.079	0.058	0.139	0.122	0.154	0.188	0.184
K	0.011	0.013	0.008	0.010	0.017	0.021	0.013	0.017	0.033
	0.081	0.094	0.087	0.068	0.156	0.143	0.167	0.205	0.217
Mg/Fe*+Mg	0.443	0.435	0.446	0.434	0.464	0.467	0.473	0.471	0.495
	Sample 2A								
	GRAIN 1			GRAIN 2			GRAIN 3		
	1	2	3	1	2	3	1	2	3
SiO ₂	50.25	48.66	45.03	49.10	48.85	46.76	46.70	45.97	45.54
TiO ₂	0.13	0.14	0.36	0.20	0.17	0.32	0.23	0.30	0.28
Al ₂ O ₃	4.34	5.79	9.93	4.75	6.12	8.16	8.23	8.35	8.76
FeO*	18.18	19.82	19.98	19.87	19.54	20.73	21.15	22.32	21.98
MnO	0.68	0.80	0.64	0.68	0.55	0.59	0.45	0.63	0.58
MgO	11.13	9.46	8.23	10.31	9.87	8.46	8.20	7.40	7.52
CaO	12.26	12.12	12.37	12.15	12.09	12.07	11.75	11.93	11.94

Table 1. Representative electron microprobe analyses of constituent grains within amphibole concentrates from calc-silicate horizons in the Goldenville Formation, southwestern Meguma Terrane.

Na ₂ O	0.26	0.32	0.59	0.33	0.35	0.49	0.44	0.43	0.58
K ₂ O	0.03	0.04	0.11	0.02	0.06	0.08	0.08	0.07	0.08
Total	97.26	97.15	97.24	97.41	97.60	97.66	97.23	97.40	97.26
Numbers of Ions on the Basis of 23(O)									
Si	7.498	7.342	6.833	7.390	7.313	7.060	7.082	1.019	6.961
Al ^{IV}	0.502	0.658	1.167	0.610	0.687	0.940	0.918	0.981	1.039
Al ^{VI}	0.261	0.372	0.609	0.233	0.393	0.512	0.553	0.522	0.539
Ti	0.015	0.016	0.041	0.023	0.019	0.036	0.026	0.034	0.032
Fe ⁺²	2.269	2.501	2.536	2.501	2.447	2.618	2.683	2.850	2.810
Mn	0.086	0.102	0.082	0.087	0.070	0.075	0.058	0.081	0.075
Mg	2.476	2.128	1.862	2.313	2.203	1.904	1.854	1.684	1.713
XM 1-3	0.107	0.119	0.130	0.157	0.132	0.145	0.174	0.171	0.169
Ca	1.960	1.959	2.011	1.959	1.939	1.953	1.909	1.952	1.955
	2.067	2.078	2.141	2.116	2.071	2.098	2.083	2.123	2.124
N	0.075	0.094	0.174	0.096	0.102	0.143	0.129	0.127	0.172
K	0.006	0.008	0.021	0.004	0.011	0.015	0.015	0.014	0.016
	0.081	0.102	0.195	0.100	0.113	0.158	0.144	0.141	0.188
Mg/Fe ⁺ +Mg	0.522	0.460	0.423	0.480	0.474	0.421	0.409	0.371	0.379
Sample 3A									
	GRAIN 1			GRAIN 2			GRAIN 3		
	1	2	3	1	2	3	1	2	3
SiO ₂	46.41	46.10	46.11	44.34	44.15	43.46	43.24	42.83	40.74
TiO ₂	0.63	0.62	0.67	0.59	0.61	0.53	0.67	0.70	0.48
Al ₂ O ₃	9.63	9.30	9.79	12.19	12.41	12.27	11.78	11.47	14.40
FeO*	15.50	15.38	15.80	17.70	18.09	18.26	19.50	19.86	20.83
MnO	0.93	1.25	0.91	0.37	0.42	0.52	0.16	0.63	0.55
MgO	11.55	11.95	11.40	9.11	8.64	8.69	8.16	8.08	6.52
CaO	11.31	11.36	11.31	11.35	11.44	11.85	11.74	12.17	11.93
Na ₂ O	1.05	1.18	1.05	1.24	1.25	1.41	1.24	1.25	1.43
K ₂ O	0.24	0.18	0.21	0.61	0.33	0.27	0.33	0.40	0.37
Total	97.25	97.32	97.25	97.50	97.34	97.26	96.82	97.39	97.25
Numbers of Ions on the Basis of 23(O)									
Si	6.884	6.852	6.852	6.641	6.628	6.563	6.590	6.537	6.260
Al ^{IV}	1.116	1.148	1.148	1.359	1.372	1.437	1.410	1.463	1.740
Al ^{VI}	0.568	0.481	0.567	0.793	0.824	0.747	0.706	0.601	0.868
Ti	0.070	0.069	0.075	0.066	0.069	0.060	0.077	0.080	0.055
Fe ⁺²	1.923	1.912	1.963	2.217	2.271	2.306	2.485	2.535	2.677
Mn	0.117	0.157	0.115	0.047	0.053	0.067	0.021	0.081	0.072
Mg	2.554	2.648	2.525	2.034	1.934	1.956	1.854	1.838	1.494

Table 1. Representative electron microprobe analyses of constituent grains within amphibole concentrates from calc-silicate horizons in the Goldenville Formation, southwestern Meguma Terrane.

XM 1-3	0.232	0.267	0.245	0.157	0.151	0.136	0.143	0.135	0.166
Ca	1.798	1.809	1.801	1.821	1.840	1.917	1.917	1.990	1.964
	2.030	2.076	2.046	1.978	1.991	2.053	2.060	2.125	2.130
N	0.302	0.340	0.303	0.360	0.364	0.413	0.366	0.370	0.426
K	0.045	0.034	0.040	0.117	0.063	0.052	0.064	0.078	0.073
	0.347	0.374	0.343	0.477	0.427	0.465	0.430	0.448	0.499
Mg/Fe*+Mg	0.570	0.547	0.563	0.478	0.460	0.459	0.427	0.420	0.358

+ spot 1, 2, and 3

* Fe as total FeO

an age of 385.1 ± 2.6 Ma and a $^{40}\text{Ar}/^{36}\text{Ar}$ ratio of 295.7 ± 6.3 (Fig. 4).

The amphibole concentrate prepared from sample 2A records an internally discordant $^{40}\text{Ar}/^{39}\text{Ar}$ age spectrum with a total-gas age of 381.6 ± 6.6 Ma (Fig. 3). Apparent K/Ca ratios are variable in the four low-temperature increments (550–650°C) whereas the remaining 12 increments (700°C–fusion) record constant ratios. These increments are therefore interpreted to represent gas evolved from a chemically homogeneous mineral phase. Each of the 8 increments evolved between 700 and 815°C contains >60% non-atmospheric ^{40}Ar components. These increments define a $^{36}\text{Ar}/^{40}\text{Ar}$ vs. $^{39}\text{Ar}/^{40}\text{Ar}$ correlation line (MSWD = 3.2) yielding intercepts corresponding to a $^{40}\text{Ar}/^{36}\text{Ar}$ ratio of 307.5 ± 10.4 and an age of 369.1 ± 2.9 Ma.

An internally discordant age spectrum is also displayed by the amphibole concentrate prepared from sample 3A (Fig. 3). A total-gas age of 376.5 ± 5.2 Ma is defined. The five low-temperature increments (500–700°C) record variable apparent K/Ca ratios. The remaining 10 increments (720°C–fusion) are characterized by a K/Ca ratio of c. 0.050, suggesting experimental evolution of gas from a chemically homogeneous phase. Each of the increments evolved between 720 and 845°C contains >80% non-atmospheric ^{40}Ar . These define a $^{36}\text{Ar}/^{40}\text{Ar}$ vs. $^{39}\text{Ar}/^{40}\text{Ar}$ correlation line (MSWD = 4.7) with intercepts corresponding to a $^{40}\text{Ar}/^{36}\text{Ar}$ ratio of 319.1 ± 9.8 and an age of 366.9 ± 5.6 Ma.

Biotite and muscovite

Dallmeyer and Keppie (1987) previously reported $^{40}\text{Ar}/^{39}\text{Ar}$ plateau ages of 341.1 ± 0.9 Ma and 346.9 ± 0.9 Ma for biotite and muscovite within schist from the exposure where calc-silicate sample 1A was collected (location 15 in Fig. 3 of Dallmeyer and Keppie, 1987). A muscovite concentrate has been prepared from schist at location 2. This displays an internally concordant $^{40}\text{Ar}/^{39}\text{Ar}$ age spectrum corresponding to a plateau age of 346.6 ± 0.9 Ma (Table 3, Fig. 5). Apparent K/Ca ratios are very large and display no significant or systematic variations throughout the analysis. Biotite and muscovite concentrates were also prepared from schist at location 3 (Fig. 6). These display minor spectra discordance with slightly younger ages recorded in the lowest

temperature increments; however, plateau ages of 285.4 ± 0.7 Ma (biotite) and 309.9 ± 0.8 Ma (muscovite) are defined by >90% of the total gas evolved from each sample (Fig. 5). Apparent K/Ca ratios are very large and display no significant or systematic variations in either analysis.

INTERPRETATION

The three amphibole concentrates record variably discordant $^{40}\text{Ar}/^{39}\text{Ar}$ age spectra. The larger volume increments within those portions of each analysis characterized by internally consistent K/Ca ratios yield $^{36}\text{Ar}/^{40}\text{Ar}$ vs. $^{39}\text{Ar}/^{40}\text{Ar}$ correlation lines defining ordinate intercepts corresponding to an atmospheric $^{40}\text{Ar}/^{36}\text{Ar}$ ratio. Abcissa intercepts yield $^{40}\text{Ar}/^{39}\text{Ar}$ ratios corresponding to ages which range between 366.9 ± 5.6 Ma and 385.1 ± 2.6 Ma. These are interpreted to date the last cooling through temperatures required for intra-crystalline retention of argon. Harrison (1981) suggested that argon retention occurs at ca. $500 \pm 25^\circ\text{C}$ in hornblende at the cooling rates likely to be encountered in most geologic settings. It is uncertain if similar temperatures characterize the amphibole compositions examined in the present study.

Lack of internal variations in apparent K/Ca ratios indicate that the biotite and muscovite concentrates are homogeneous and characterized by chemically uniform grains. The concentrates record plateau ages ranging between 285.4 ± 0.7 Ma and 346.6 ± 0.9 Ma. These are interpreted to date the last cooling through argon retention temperatures. Harrison *et al.* (1985) have demonstrated that the temperatures required for intracrystalline retention of argon in biotite are dependent upon composition and increase with decreasing Fe/Mg ratio. They suggest that closure occurs at ca. $300 \pm 25^\circ\text{C}$ in the range of cooling rates likely to be encountered in most geologic environments. Detailed experimental evaluation of the temperatures required for intracrystalline retention of argon in muscovite have not been carried out. Using the preliminary experimental data of Robbins (1972) in the diffusion equations of Dodson (1973) suggests that temperatures of ca. 350°C may be appropriate. These are similar to closure temperatures suggested for muscovite on the basis of empirical comparison with argon systems within other mineral species (e.g., Wagner *et al.*, 1977; Jäger, 1979).

Table 2. $^{40}\text{Ar}/^{39}\text{Ar}$ Analytical Data for Incremental Heating Experiments on Amphibole Concentrates From Calc-Silicate Horizons Within the Goldenville Formation, Southwestern Meguma Terrane.

Release temp (°C)	$(^{40}\text{Ar}/^{39}\text{Ar})^*$	$(^{36}\text{Ar}/^{39}\text{Ar})^*$	$(^{37}\text{Ar}/^{39}\text{Ar})^c$	^{39}Ar % of total	% ^{40}Ar non-atmos. ⁺	^{36}Ar %Ca	Apparent Age (Ma)**
Sample 1A; J = 0.008225							
550	133.84	0.36783	41.792	3.93	21.29	3.09	389.2±10.6
625	53.61	0.10027	56.464	4.64	53.18	15.32	392.7± 6.4
700	58.33	0.11255	39.897	5.11	48.47	9.64	386.0± 5.5
750	51.59	0.08837	45.593	7.98	56.48	14.03	397.9± 4.2
775	39.76	0.04845	40.397	14.93	72.14	22.68	391.2± 3.1
785	41.78	0.05330	34.211	8.64	68.87	17.46	390.8± 3.9
800	45.08	0.06464	31.785	5.12	63.28	13.38	387.2± 7.7
825	38.20	0.04482	25.887	6.12	70.76	15.71	367.7± 5.3
835	34.82	0.03488	27.542	7.67	76.74	21.48	364.2± 3.7
865	35.77	0.03380	41.556	16.36	81.40	33.44	396.7± 2.2
885	38.26	0.03916	41.743	9.54	78.50	28.99	407.9± 3.3
925	42.12	0.05408	52.004	7.81	71.97	26.15	413.8± 3.5
Fusion	96.06	0.23355	48.619	2.15	32.22	5.66	420.7±21.8
Total	46.57	0.07100	40.137	100.00	67.59	21.03	392.6± 4.5
Sample 2A; J = 0.008375							
500	91.38	0.20519	11.781	1.96	34.67	1.56	427.4± 8.0
550	60.54	0.11749	16.448	1.85	44.82	3.81	372.9±17.1
600	73.86	0.16904	22.661	2.10	34.83	3.65	356.6±24.7
650	51.92	0.09551	28.483	5.29	50.04	8.11	361.0±10.5
700	34.82	0.03243	23.752	20.79	77.94	19.92	374.6± 2.3
720	34.25	0.03013	22.011	12.33	79.16	19.87	373.9± 1.9
740	34.13	0.03176	21.793	6.33	77.62	18.67	366.0± 5.6
755	31.83	0.02107	23.536	8.18	86.36	30.38	378.8± 5.4
770	30.64	0.01739	22.961	11.91	89.23	35.91	376.9± 3.6
785	32.88	0.02327	22.878	9.80	84.66	26.74	383.1± 2.3
800	42.01	0.05418	22.190	5.90	66.12	11.14	382.2± 6.2
815	42.78	0.05373	22.196	5.33	67.04	11.24	393.4± 3.5
830	54.16	0.08548	21.637	3.06	56.57	6.89	417.2±15.5
850	57.31	0.11344	22.681	2.10	44.67	5.44	355.1±23.5
870	53.52	0.09871	22.237	1.63	48.83	6.13	361.7±13.7
Fusion	631.70	1.96877	22.790	1.45	8.19	0.31	657.5±81.5
Total	47.96	0.07456	22.773	100.00	72.80	19.45	381.6± 6.6
Sample 3A; J = 0.007322							
500	172.34	0.22589	12.904	0.67	61.87	1.55	1047.4±30.9
550	106.96	0.22034	16.178	0.55	40.34	2.00	499.7±70.9
600	94.65	0.26071	8.363	0.54	19.31	0.87	227.7±72.6
650	110.97	0.28123	12.703	1.40	26.03	1.23	348.7±23.8
700	39.46	0.02921	13.386	6.25	80.84	12.46	381.6± 5.3
720	35.23	0.01664	10.706	11.82	88.47	17.50	373.1± 3.3
740	34.84	0.01680	10.303	14.56	88.11	16.68	367.9± 2.3
760	33.01	0.01158	10.329	9.55	92.13	24.26	364.9± 5.8
780	31.99	0.00914	9.792	10.48	94.00	29.15	361.0± 1.9
800	33.15	0.01007	9.634	7.79	93.34	26.02	370.4± 4.2
820	32.51	0.00595	9.621	17.96	96.95	43.96	376.6± 2.0

Table 2. $^{40}\text{Ar}/^{39}\text{Ar}$ Analytical Data for Incremental Heating Experiments on Amphibole Concentrates From Calc-Silicate Horizons Within the Goldenville Formation, Southwestern Meguma Terrane.

Release temp (°C)	$(^{40}\text{Ar}/^{39}\text{Ar})^*$	$(^{36}\text{Ar}/^{39}\text{Ar})^*$	$(^{37}\text{Ar}/^{39}\text{Ar})^c$	^{39}Ar % of total	% ^{40}Ar non-atmos. ⁺	^{36}Ar %Ca	Apparent Age (Ma)**
845	32.86	0.00685	9.985	15.81	96.26	39.62	377.9± 1.9
865	46.16	0.05103	9.782	0.87	69.02	5.21	380.4±26.8
890	64.92	0.11974	9.562	0.55	46.67	2.17	363.5±28.7
Fusion	188.51	0.52375	9.613	1.19	18.31	0.50	408.6±34.3
Total	38.64	0.02667	10.323	100.00	89.07	26.94	376.5± 5.2

*measured

^ccorrected for post-irradiation decay of ^{37}Ar (35.1 day half-life).⁺ $[(^{40}\text{Ar}_{\text{tot.}} - (^{36}\text{Ar}_{\text{atmos.}})(295.5))] / ^{40}\text{Ar}_{\text{tot.}}$ **calculated using correction factors of Dalrymple *et al.* (1981); two sigma, intralaboratory errors.Table 3. $^{40}\text{Ar}/^{39}\text{Ar}$ Analytical Data for Incremental Heating Experiments on Muscovite and Biotite Concentrates from Schist Within the Goldenville Formation, Southwestern Meguma Terrane.

Release temp (°C)	$(^{40}\text{Ar}/^{39}\text{Ar})^*$	$(^{36}\text{Ar}/^{39}\text{Ar})^*$	^{39}Ar % of total	% ^{40}Ar non-atmos. ⁺	Apparent Age (Ma)**
<u>Muscovite</u>					
Sample 3B; J = 0.007837					
490	29.57	0.02538	1.35	74.62	287.7±9.5
525	24.67	0.00521	3.65	93.73	300.5±3.1
575	25.27	0.00512	10.06	93.98	307.9±1.3
610	24.93	0.00373	25.90	95.56	308.8±0.7
640	24.84	0.00323	7.92	96.13	309.5±1.0
710	25.26	0.00436	10.42	94.88	310.5±1.4
775	24.79	0.00253	34.18	96.96	311.4±0.7
Fusion	27.60	0.01091	6.52	88.29	315.3±2.3
Total	25.17	0.00430	100.00	95.03	309.6±1.1
Total without 490, 525°C and fusion			88.48		309.9±0.8
Sample 2B; J = 0.008572					
480	27.86	0.01815	1.76	80.73	317.9±4.9
510	25.55	0.00335	8.95	96.10	344.6±1.1
540	25.23	0.00183	42.54	97.83	346.2±0.6
580	25.25	0.00177	12.27	97.91	346.6±0.9
630	25.41	0.00246	6.72	97.11	346.1±1.9
690	25.42	0.00215	8.16	97.49	347.4±1.8
740	25.35	0.00145	16.68	98.28	349.1±1.0
Fusion	35.94	0.03381	2.91	72.18	362.1±4.2
Total	25.67	0.00318	100.00	96.64	346.6±1.1
Total without 480°C and fusion			95.33		346.6±0.9

Table 3. $^{40}\text{Ar}/^{39}\text{Ar}$ Analytical Data for Incremental Heating Experiments on Muscovite and Biotite Concentrates from Schist Within the Goldenville Formation, Southwestern Meguma Terrane.

Release temp (°C)	$(^{40}\text{Ar}/^{39}\text{Ar})^*$	$(^{36}\text{Ar}/^{39}\text{Ar})^*$	^{39}Ar % of total	% ^{40}Ar non-atmos. ⁺	Apparent Age (Ma)**
<u>Biotite</u>					
Sample 3B; J = 0.007961					
475	24.46	0.01542	6.46	81.35	265.2±1.6
500	21.98	0.00182	22.42	97.52	284.2±0.4
525	21.86	0.00124	16.22	98.30	284.9±0.8
600	23.71	0.00708	8.82	91.14	286.3±1.3
675	22.36	0.00284	12.46	96.23	285.2±0.9
750	22.07	0.00179	23.14	97.58	285.5±0.6
825	24.44	0.00885	8.93	89.27	288.9±1.3
Fusion	94.14	0.23772	1.55	25.37	314.0±6.6
Total	23.68	0.00747	100.00	94.04	284.5±0.8
Total without 475°C and fusion			91.99		285.4±0.7

*measured

+ $[(^{40}\text{Ar}_{\text{tot.}} - (^{36}\text{Ar}_{\text{atmos.}})(295.5)) / ^{40}\text{Ar}_{\text{tot.}}]$

**calculated using correction factors of Dalrymple *et al.* (1981); $^{37}\text{Ar}/^{39}\text{Ar}$ corrected ratios < 0.02 in all analyses; two sigma, intralaboratory errors.




Fig. 3. $^{40}\text{Ar}/^{39}\text{Ar}$ incremental-release age spectra of amphibole from calc-silicate in the southwestern Meguma Terrane, Nova Scotia. Data plotted as in Figure 2. All spectra have similar abscissa and ordinate coordinates. Total-gas ages listed on each spectra.

GEOLOGIC SIGNIFICANCE

Amphibole, muscovite, and biotite at location 1 immediately north of the Shelburne pluton record post-metamorphic cooling ages of 385.1±2.6 Ma, 346.9±0.9 Ma and 341.1±0.9 Ma. Generally similar mica ages were reported by Dallmeyer and Keppie (1987) for other locations within the andalusite-staurolite-cordierite zone north of the Shelburne pluton. These likely date diachronous cooling following a contact thermal overprint associated with emplacement of the two plutons at *ca.* 370-375 Ma. Relative variations in the cooling ages suggest that the amphibole composition examined in this study has a higher closure temperature than muscovite, and may therefore approximate the *ca.* 500°C value suggested by Harrison (1981) for hornblende. The biotite and muscovite spectra are internally concordant and provide no record for a later thermal disturbance of intracrystalline argon systems.

Amphibole and muscovite south of the Port Mouton pluton (location 2) record post-metamorphic cooling ages of 369.1±2.9 Ma and 346.9±0.9 Ma. These are interpreted to date diachronous cooling through appropriate argon retention temperatures following emplacement of the Port Mouton pluton. Amphibole, muscovite and biotite south of the Barrington Passage pluton (location 3) record cooling ages of 366.9±5.6 Ma, 309.9±0.8 Ma, and 285±0.7 Ma. The amphibole age probably dates cooling following a thermal overprint associated with the intrusion of the Barrington Passage pluton; however, the *ca.* 310 Ma plateau age recorded by coexisting muscovite is similar to plateau ages recorded by Dallmeyer and Keppie (1987) for muscovite and biotite north of Sable Island. The ages are inferred to date cooling following reheating adjacent to an unexposed pluton of approximately the same age as the *ca.* 315 Ma Wedgeport pluton. The *ca.* 285 Ma plateau age of the coexisting biotite could be inter-




Fig. 4. $^{36}\text{Ar}/^{40}\text{Ar}$ vs. $^{39}\text{Ar}/^{40}\text{Ar}$ isotope correlation diagrams for amphibole from calc-silicate horizons in the southwestern Meguma Terrane, Nova Scotia: atm. = ratio in present-day atmosphere.




Fig. 5. $^{40}\text{Ar}/^{39}\text{Ar}$ incremental-release age spectrum of muscovite from schist interbedded with calc-silicate at location 2. Data plotted as in Figure 2. Plateau age is listed.




Fig. 6. $^{40}\text{Ar}/^{39}\text{Ar}$ incremental-release age spectra of muscovite and biotite from schist interbedded with calc-silicate at location 3. Data plotted as in Figure 2. Plateau ages are listed.

preted to date either slow cooling through the biotite blocking temperature following the *ca.* 315 Ma thermal event of a complete rejuvenation following a younger thermal event. The latter alternative finds support in: (1) the previously reported muscovite plateau age of *ca.* 287 Ma recorded in southern Cape Sable Island (Dallmeyer and Keppie, 1987); and (2) the low-temperature spectrum discordance in the muscovite at location 3 and at other adjacent locations, which are consistent with slight volume-diffusive loss of argon during a low grade, *ca.* 290 Ma thermal event. Collectively these data at location 3 suggest that: (1) contact metamorphic temperatures during the *ca.* 315 Ma event exceeded those required to totally reset muscovite and biotite but remained below those required to reset amphibole (i.e., between *ca.* 500-400°C); and (2) during the *ca.* 290 Ma event, contact metamorphic temperatures were sufficient to totally rejuvenate biotite but remained below those required to totally reset muscovite (i.e., between *ca.* 350-325°C).

CONCLUSIONS

Extensive contact aureoles appear to have developed during emplacement of the Port Mouton, Shelburne and Barrington Passage plutons at *ca.* 370-375 Ma. This thermal overprint completely rejuvenated intracrystalline argon systems within all minerals which had grown during an earlier (*ca.* 400-410 Ma) M_1 regional metamorphism. Post-metamorphic cooling through calc-silicate amphibole closure temperatures (*ca.* 500°C?) occurred shortly after pluton emplacement. Cooling through the lower temperatures required for intracrystalline retention of argon within muscovite and biotite occurred between 340 and 350 Ma.

Southern portions of the contact metamorphic aureoles of the Shelburne and Barrington Passage plutons were affected by later reheating associated with emplacement of plutons at *ca.* 315 and 290 Ma. Argon systems within muscovite and biotite were

variably rejuvenated during these events, but were not reset in hornblende. Therefore, the amphibolite facies metamorphism recorded in southwestern portions of the Meguma Terrane appears to largely reflect the results of an *ca.* 370 Ma contact thermal event.

ACKNOWLEDGEMENTS

Funds for this project were provided by the Nova Scotia Provincial Government as part of a joint Federal-Provincial Mineral development Agreement 1984-89. The Director of the Mineral Resources Division of the Nova Scotia Department of Mines and Energy has authorized publication. This paper represents a contribution of I.G.C.P. Project 233, "Terranes in the Circum-Atlantic Paleozoic Orogens." Mr. J. L. Cho and the Department of Geology, Yonsei University, Seoul, Korea, are thanked for their assistance during microprobe analysis.

- ALEXANDER, E.C., JR., MICHELSON, G.M., and LANPHERE, M.A. 1978. MMhb-1: A new $^{40}\text{Ar}/^{39}\text{Ar}$ dating standard. *In* Short papers of the Fourth International Conference, Geochronology, Cosmochronology, isotope Geology. Edited by R.E. Zartman. United States Geological Survey, Open File Report 78-701, pp. 6-8.
- BORQUE, A.D. 1985. Migmatization and metamorphism associated with the Barrington Passage Pluton, Shelburne and Yarmouth Counties, Nova Scotia. B.Sc. Honours thesis, Acadia University, Wolfville, Nova Scotia.
- CHO, D.L. 1985. Petrography and Rb-Sr ages of the Imgye Granite in northeastern Ockcheon Belt, Korea. Unpublished Ms. thesis, Yonsei University.
- DALLMEYER, R.D. and KEPPIE, J.D. 1987. Polyphase late Paleozoic tectonothermal evolution of the southwestern Meguma Terrane, Nova Scotia: Evidence from $^{40}\text{Ar}/^{39}\text{Ar}$ mineral ages. *Canadian Journal of Earth Sciences*, 24, pp. 1242-1254.
- DALRYMPLE, G.B., ALEXANDER, E.C., LANPHERE, M.A., and KRAKER, G.P. 1981. Irradiation of samples for $^{40}\text{Ar}/^{39}\text{Ar}$ dating using the Geological Survey TRIGA reactor. *United States Geological Survey Professional Paper 1176*, 55 p.
- DODSON, M.H. 1973. Closure temperature in cooling geochronological and petrological systems. *Contributions to Mineralogy and Petrology*, 40, pp. 259-274.
- HARRISON, T.M. 1981. Diffusion of ^{40}Ar in hornblende. *Contributions to Mineralogy and Petrology*, 78, pp. 324-331.
- HARRISON, T.M., DUNCAN, I., and MCDUGALL, I. 1985. Diffusion of ^{40}Ar in biotite: Temperature, pressure and compositional effects. *Geochimica et Cosmochimica Acta*, 49, pp. 2461-2468.
- HAWTHORNE, F.C. 1982. Crystal chemistry of the amphiboles. *In* Amphiboles and other hydrous pyriboles Mineralogy. Edited by D.R. Veblen. *Reviews in Mineralogy*, 9a, Mineralogical Society of America, pp. 1-9.
- JÄGER, E. 1979. Introduction to geochronology. *In* Lectures in isotope geology. Edited by E. Jäger and J.C. Hunziker. Berlin, Springer-Verlag, pp. 1-12.
- KEPPIE, J.D. 1976. Structural model for the saddle reef and associated gold veins in the Meguma Group, Nova Scotia. Nova Scotia Department of Mines, Paper 76-1, 34 p.
- KEPPIE, J.D., CURRIE, K., MURPHY, J.B., PICKERILL, R.K., FYFFE, L.R., and ST. JULIEN, P. 1985. Appalachian geotraverse (Canadian Mainland). Geological Association of Canada Excursion 1, 181 p.
- KEPPIE, J.D. and MUECKE, G.K. 1979. Metamorphic map of Nova Scotia. Nova Scotia Department of Mines and Energy. Scale 1:1,000,000.
- KEPPIE, J.D. and KROGH, T. 1988. (*In press.*) U-Pb crystallization ages for zircon and monazite within granitic plutons from the southwestern Meguma Terrane, Nova Scotia. *Bulletin of the Geological Society of America*.
- KEPPIE, J.D., ODOM, L., and CORMIER, R.F. 1983. Tectonothermal evolution of the Meguma Terrane: radiometric controls. *Geological Society of America, Abstracts With Programs*, 1983, 15, p. 136.
- LEAKE, B. 1978. Nomenclature of amphiboles. *Canadian Mineralogist*, 16, pp. 501-520.
- MISNER, A.R. 1986. Metamorphism of the northern part of the Shelburne Metamorphic Complex, Shelburne and Yarmouth Counties, Nova Scotia. B.Sc. Honours thesis, Acadia University, Wolfville, Nova Scotia.
- RADICATI DI BROZOLO, F., HUNEKE, J.C., PAPANSTASSIOU, D.A., and WASSERBURG, J. 1981. $^{40}\text{Ar}/^{39}\text{Ar}$ and Rb-Sr age determinations on Quaternary volcanic rocks. *Earth and Planetary Science Letters*, 53, pp. 233-244.
- RAESIDE, R.P., WHITE, C.E., and WENTZELL, B.D. 1985. The metamorphic development of the Shelburne Complex, southwest Nova Scotia. Geological Association of Canada, Program with Abstracts, 10, p. A50.
- ROBBINS, G.A. 1972. Radiogenic argon diffusion of muscovite under hydrothermal conditions. M.Sc. thesis, Brown University, Providence, Rhode Island.
- RODDICK, J.C. 1978. The application of isochron diagrams in $^{40}\text{Ar}/^{39}\text{Ar}$ dating: A discussion. *Earth and Planetary Science Letters*, 41, pp. 233-244.
- RODDICK, J.C., CLIFF, R.A., and REX, D.C. 1980. The evolution of excess argon in alpine biotite - $^{40}\text{Ar}/^{39}\text{Ar}$ analysis. *Earth and Planetary Science Letters*, 48, pp. 185-208.
- ROSS, D.M. 1985. Structure and metamorphism of the Pubnico area, Yarmouth County, Nova Scotia. B.Sc. Honours thesis, Acadia University, Wolfville, Nova Scotia.
- SAGE, J.D. 1984. Variable water pressure metamorphic assemblages in the Meguma Group, Nova Scotia. M.Sc. thesis, Virginia Polytechnic Institute and State University, Blacksburg, Virginia.
- STEIGER, R.H. and JÄGER, E. 1977. Subcommission on geochronology: convention on the use of decay constants in geo- and cosmochronology. *Earth and Planetary Sciences*, 36, pp. 359-362.
- TAYLOR, F.C. and SCHILLER, E.A. 1966. Metamorphism of the Meguma Group of Nova Scotia. *Canadian Journal of Earth Sciences*, 3, pp. 959-974.
- WAGNER, G.A., REIMER, G.M., and JÄGER, E. 1977. Cooling ages derived from apatite fission, mica Rb-Sr and K-Ar dating: the uplift and cooling history of the central Alps. *Padova University Institute of Geology Mineral Memoir*, 30, pp. 1-27.
- WENTZELL, B.D. 1985. The transition from staurolite to sillimanite zone, Port LaTour, Nova Scotia. B.Sc. Honours thesis, Acadia University, Wolfville, Nova Scotia.
- WHITE, C.E. 1984. Structure and metamorphism of the Jordon River Valley, Shelburne County, Nova Scotia. B.Sc. Honours thesis, Acadia University, Wolfville, Nova Scotia.
- YORK, D. 1969. Least squares fitting of a straight line with correlated errors. *Earth and Planetary Science Letters*, 5, pp. 320-324.