Citation refers to <u>Issue</u> Number and <u>Page</u> Number.

Abrasion, particle, a small circular flume for laboratory studies on: GEES Activities, resource, minerals, in the Canadian offshore: CROSBY Are basal transgressive sands chiefly inlet-filling sands? KUMAR Atlantic, inner, shelf between Cape Henry and Cape Hatteras, Quaternary	1 1 1	15-16 30-36 12-14
sedimentation on the: SWIFT AVIGNONE, N.F., see SWIFT et al	1 1	5-11 5-11
Bank, Browns, sand waves on, observed from the Shelf Diver: DRAPEAU Bar, tidal and channel, river deposits Gaspé Nord, Quebec: COSTELLO	3 2	90-101 68-71
Basal transgressive sands chiefly inlet-filling sands? KUMAR	1	12-14
Basin, Kane, sedimentological reconnaissance survey of: KRAVITZ Bay, Covehead, Prince Edward Island - A preliminary study, benthonic foraminiferal	1	17-20
ecology in: SLESSOR Bay, Fundy of, Nova Scotia, sedimentological sampling and results from the diver lock-out facility of the submersible Shelf Diver: PELLETIER	2 3	48-64 102-109
Bay, Kouchibouguac, New Brunswick, beach processes and shoreline changes: McCANN		102-109
Beach processes and shoreline changes, Kouchibouguac Bay, New Brunswick: McCANN Benthonic foraminiferal ecology in Covehead Bay, Prince Edward Island - A		116-117
preliminary study: SLESSOR	2,	48-64
Benthonic foraminifera in the Restigouche Estuary, studies of: Part 1. Faunal distribution patterns near pollution sources: SCHAFER Benthonic foraminifera, modern, in the northwest Gulf of St. Lawrence, the	3	121-134
distribution of: HOOPER	2	74-78
Bermuda, Part II, Harrington Sound, continuous seismic survey of: GEES	3	118-120
Bermuda Platform, Part I: Castle Harbour a continuous seismic survey of the: GEES Bight, New York and vicinity, integrated continental-shelf and marginal marine environmental research program: sediments, organisms, and waters in :	1	21-25
FRIEDMAN	1	26-29
Browns Bank, sand waves on, observed from the Shelf Diver: DRAPEAU BRYANT, E.A., see McCANN, S.B.	3 3	90-101 116-117
	5	110-111
Canadian offshore, mineral resources activities in the: CROSBY	1	30-36
Castle Harbour, Part I, a continuous seismic survey of the Bermuda Platform: GEES Changes, shoreline, Kouchibouguac Bay, New Brunswick, beach processes and: McCANN Channel, Providence, Bahamas, Eocene deep water sediments from the northeast:	1 3	21-25 116-117
STEHMAN	2	65-67
Channel, river and tidal bar deposits Gaspé Nord, Quebec: COSTELLO	2	68-71
Circular, small, flume for laboratory studies on particle abrasion: GEES	1	15-16
Clay flake orientation in flocculated illite - an electron microscope study: O'BRIEN CROSBY, D.G., Mineral resource activities in the Canadian offshore	2 1	79-80 30-36
Continuous seismic survey of Bermuda, Part II, Harrington Sound: GEES	3	118-120
Continental-shelf, integrated, and marginal marine environmental research program: sediments, organisms, and waters in New York Bight and Vicinity: FRIEDMAN	1	26-29
COSTELLO, W.R., River channel and tidal bar deposits Gaspé Nord, Quebec	2	68-71
Covehead Bay, Prince Edward Island - A preliminary study, benthonic foraminiferal ecology in: SLESSOR	2	48-64
Deposits, bar tidal, and channel river, Gaspé Nord, Quebec: COSTELLO Deposits, Quaternary, from the south coast of the St. Lawrence Estuary, Quebec,	2	68-71
exotic pebbles in: DIONNE	3	110-112
Diapiric structure near Sable Island ~ Scotian Shelf: KING	1	1-4
DILL, C.E. Jr., see SWIFT et al DIONNE, J-C., Exotic pebbles in Quaternary deposits from the south coast of the	1	5-11
St. Lawrence Estuary, Quebec Distribution of modern benthonic foraminifera in the northwest Gulf of St.	3	110-112
Lawrence: HOOPER Distribution patterns, faunal, near pollution sources: Part I of studies of	2	74-78
benthonic foraminifera in the Restigouche Estuary: SCHAFER Diver, lock-out facility of the submersible Shelf Diver, Bay of Fundy, Nova Scotia,	3	121-134
sedimentological sampling and results from: PELLETIER	3	102-109
Diver, Shelf, sand waves on Browns Bank observed from: DRAPEAU Diver, Shelf, submersible, Bay of Fundy, Nova Scotia, sedimentological sampling and	3	90-101
results from the diver lock-out facility of: PELLETIER DRAPEAU, G., Sand waves on Browns Bank observed from the Shelf Diver	3 3	102-109 90-101

.

Ecology, foraminiferal, benthonic, in Covehead Bay, Prince Edward Island - A preliminary study: SLESSOR	2	48-64
Electron microscope study, an, - clay flake orientation in flocculated illite:		
O'BRIEN Environmental, marine, marginal research program, integrated continental-shelf	2	
and: sediments, organisms, and waters in New York Bight and vicinity: FRIEDMAN Eocene deep water sediment from the northeast Providence Channel, Bahamas: STEHMAN	1 2	
Estuary, Restigouche, studies of benthonic foraminifera in: Part I. Faunal distribution patterns near pollution sources: SCHAFER	3	121-13
Estuary, St. Lawrence, Quebec, exotic pebbles in Quaternary deposits from the south coast of: DIONNE		110-1
	3	110-1.
Facility, diver lock-out, of the submersible Shelf Diver, Bay of Fundy, Nova Scotia, sedimentological sampling and results from: PELLETIER	3	90-10
Faunal distribution patterns near pollution sources: Part I of studies of benthonic foraminifera in the Restigouche Estuary: SCHAFER	3	121-1
Flake, clay, orientation in flocculated illite - an electron microscope study: O'BRIEN	è2	79-8
Flocculated illite - an electron microscope study, clay flake orientation in: O'BRIEN	2	79 - 8
Flume, circular, small, for laboratory studies on particle abrasion: GEES	ĩ	
Foraminifera, benthonic, in the Restigouche Estuary, studies of: Part 1. Faunal distribution patterns near pollution sources: SCHAFER	3	121-1
Foraminifera, benthonic modern, in the northwest Gulf of St. Lawrence, the distribution of: HOOPER	2	74-7
Foraminiféral, ecology, benthonic in Covehead Bay, Prince Edward Island - A preliminary study: SLESSOR	2	48-6
FRIEDMAN, G.M., Integrated continental-shelf and marginal marine environmental research program: sediments, organisms, and waters in New York Bight and	2	40 0
vicinity Fundy, bay of, Nova Scotia, sedimentological sampling and results from the diver	1	26-2
lock-out facility of the submersible Shelf Diver: PELLETIER	3	102-1
Gaspé Nord, Quebec, river channel and tidal bar deposits: COSTELLO GEES, R.A., LANGHUS, B., and MEDIOLI, F., small circular flume for laboratory	2	68-7
studies on particle abrasion GEES, R.A. and MEDIOLI, F., continuous seismic survey of the Bermuda Platform,	1	-15-1
Part I: Castle Harbour GEES, R.A. and MEDIOLI, F., Continuous seismic survey of Bermuda, Part II,	1	21-2
Harrington Sound	3	118-1
Gulf of St. Lawrence, northwest, distribution of modern benthonic foraminifera in the: HOOPER	2	74-7
Harbour, Castle, Part I, continuous seismic survey of the Bermuda Platform: GEES	1	21-2
Harrington Sound, continuous seismic survey of Bermuda, Part II: GEES Hatteras, Cape, Quaternary sedimentation on the inner Atlantic shelf between: SWIFT	3 1	118-1 5-1
Henry, Cape, and Cape Hatteras, Quaternary sedimentation on the Inner Atlantic Shelf between: SWIFT	1	5-1
HOBBS, C.H. III, Shoreline orientation and storm surge	3	113-1
HOLLIDAY, B.W., see SWIFT et al HOOPER, K., The distribution of modern benthonic foraminifera in the northwest	1	5-1
Gulf of t. Lawrence	2	74-7
Illite, flocculated - an electron microscope study, clay flake orientation in: O'BRIEN	2	79-8
Inlet-filling sands, chiefly? Are basal transgressive sands: KUMAR	1	12-1
Integrated continental-shelf and marginal marine environmental research program: sediments, organisms, and waters in New York Bight and vicinity: FRIEDMAN	1	26-2
Island, Prince Edward - A preliminary study, benthonic foraminiferal ecology in Covehead Bay: SLESSOR	2	48-6
Kane Basin, sedimentological reconnaissance survey of: KRAVITZ	1	17-2
······································	1 3	1-4 116-1
KRAVITZ, J.H. and SORENSEN, F.H., Sedimentological reconnaissance survey of Kane Basin	1	17-20
KUMAR, N. and SANDERS, J.E., Are basal transgressive sands chiefly inlet-filling	1	12-1
,,, _,, _	1	15-16 15-16
LANGHUS, B., see GEES, R.A. et al Lock-out facility, the diver, of the submersible Shelf Diver, Bay of Fundy, Nova	Ŧ	
Scotia, sedimentological sampling and results from: PELLETIER	3	102-10

MacLEAN, B., see KING, L.H.	1	1-4
McCANN, S.B. and BRYANT, E.A., Beach processes and shoreline changes, Kouchibouguac Bay, New Brunswick	3	116-117
Marginal marine environmental research program, integrated continental-shelf and: sediments, organisms, and waters in New York Bight and vicinity: FRIEDMAN	1	26-29
Marine, marginal, environmental research program, integrated continental-shelf and:	т	20-29
sediments, organisms and waters in New York Bight and vicinity: FRIEDMAN MEDIOLI, F., see GEES, R.A., et al	1 1	26-29 15-16
MEDIOLI, F., see GEES, R.A.	î	21-25
MEDIOLI, F., see GEES, R.A.	1 3	118-120
Method, new, for separation of skeletal material from a plankton sample: VILKS	2	72-73
Microscope study, an electron - clay flake orientation in flocculated illite: O'BRIEN	_	
Mineral resource activities in the Canadian offshore: CROSBY	2 1	79 - 80 30-36
Modern benthonic foraminifera in the northwest Gulf of St. Lawrence, the distribution	_	
of: HOOPER	2	74-78
New Brunswick, Kouchibouguac Bay, beach processes and shoreline changes: McCANN	3	116-117
New method for separation of skeletal material from a plankton sample: VILKS New York Bight and vicinity, integrated continental-shelf and marginal marine environmental research program: sediments, organisms, and waters in:	2	72-73
FRIEDMAN	1	26-29
Nova Scotia, Bay of Fundy, sedimentological sampling and results from the diver		
lock-out facility of the submersible Shelf Diver: PELLETIER	3	102-109
O'BRIEN, N.R., Clay flake orientation in flocculated illite - an electron microscope		
study	2	79-80
Offshore, Canadian, mineral resources activities in the: CROSBY Organisms, sediments, and waters in New York Bight and vicinity, integrated	1	30-36
continental-shelf and marginal marine environmental research program:		
FRIEDMAN	1	26-29
Orientation, flake, clay, in flocculated illite - an electron microscope study: O'BRIEN	2	79-80
Part I: Castle Harbour, a continuous seismic survey of the Bermuda Platform: GEES	1	21-25
	1	21-25 15-16
Patterns, distribution, faunal, near pollution sources: Part I of studies of	-	10 10
benthonic foraminifera in the Restigouche Estuary: SCHAFER	3	121-134
Pebbles, exotic in Quaternary deposits from the south coast of the St. Lawrence	_	
Estuary, Quebec: DIONNE PELLETIER, B.R., Sedimentological sampling and results from the diver lock-out	3	110-112
facility of the submersible Shelf Diver, Bay of Fundy, Nova Scotia	3	102-109
	1	21-25
	2	72-73
Prince Edward Island - A preliminary study, benthonic foraminiferal ecology in		
	2	48-64
Pollution sources, faunal distribution patterns near: Part I of studies of benthonic	•	
foraminifera in the Restigouche Estuary: SCHAFER Program, research, environmental, integrated continental-shelf and marginal marine:	د	121-134
	1	26-29
	3	116-117
Providence Channel, Bahamas, Eocene deep water sediments from the northeast:		
STEHMAN	2	65-67
Quaternary deposits from the south coast of the St. Lawrence Estuary, Quebec,		
	3	110-112
Quaternary sedimentation on the inner Atlantic shelf between Cape Henry and Cape		
	1	5-11
guonee, subpo nord, river shaller and stade sar aspessed or	2	68-71
Quebec, St. Lawrence Estuary, exotic pebbles in Quaternary deposits from the south coast of: DIONNE	3	110-112
never are a set and a	1	17-20
Research program, environmental, integrated continental-shelf and marginal marine:	_	
	1	26-29
Nooveroof Ministrar, accivitation in the conduction terrore terror	1	30-36
Results from the diver lock-out facility of the submersible Shelf Diver, Bay of	2	102-109
randy, nota bootra, bearmontorojeour remprend anno reserve a	3	102-103
Restigouche Estuary, studies of benthonic foraminifera in: Part I. Faunal distribution patterns near pollution sources: SCHAFER	3	121-134
River channel and tidal bar deposits Gaspé Nord, Quebec: COSTELLO	2	68-71
When the start was achested to the start of		

Sable Island - Scotian Shelf, a diapiric structure near: KING	1	4
Sample, plankton, new method for separation of skeletal material from: VILKS	2	72-73
Sampling, sedimentological, and results from the diver lock-out facility of the	_	
submersible Shelf Diver, Bay of Fundy, Nova Scotia: PELLETIER	3	-
SANDERS, J.E., see KUMAR, N.	1	
Sands, basal transgressive, chiefly inlet-filling sands? Are: KUMAR Sands, inlet-filling? Are basal transgressive sands: KUMAR	1	
Sand waves on Browns Bank observed from the Shelf Diver: DRAPEAU	1 3	
SCHAFER, C.T., Studies of benthonic foraminifera in the Restigouche Estuary: 1.	3	90-101
Faunal distribution patterns near pollution sources	2	121-134
Scotian Shelf, a diapiric structure near Sable Island: KING	1	
Sediment, deep water, Eocene from the northeast Providence Channel, Bahamas: STEHMAN	2	
Sedimentation, Quaternary, on the inner Atlantic shelf between Cape Henry and	-	05 07
Cape Hatteras: SWIFT	1	5-11
Sedimentological reconnaissance survey of Kane Basin: KRAVITZ	ī	
Sedimentological sampling and results from the diver lock-out facility of the	-	
submersible Shelf Diver, Bay of Fundy, Nova Scotia: PELLETIER	3	102-109
Sediments, organisms, and waters in New York Bight and vicinity, integrated	-	
continental-shelf and marginal marine environmental research program: FRIEDMAN	1	26-29
Seismic, continuous, survey of the Bermuda Platform, Part I: Castle Harbour: GEES	1	21-25
Seismic survey, continuous, of Bermuda, Part II, Harrington Sound: GEES	3	118-120
Separation, skeletal material of, from a plankton sample, new method for: VILKS	2	72-73
Shelf, continental-, integrated, and marginal marine environmental research		
program: sediments, organisms, and waters in New York Bight and vicinity:		
FRIEDMAN	1	26-29
Shelf Diver, sand waves on Browns Bank observed from: DRAPEAU	3	90-101
Shelf Diver, submersible, Bay of Fundy, Nova Scotia, sedimentological sampling and		
results from the diver lock-out facility of: PELLETIER	3	102-109
Shelf, inner Atlantic, between Cape Henry and Cape Hatteras, Quaternary		
sedimentation on: SWIFT	1	5-11
Shoreline changes, Kouchibouguac Bay, New Brunswick, beach processes and: McCANN	3	116 -117
Shoreline orientation and storm surve: HOBBS	3	116-117
SHIDELER, G.L., see SWIFT et al	1	5-11
Skeletal material, separation of, from a plankton sample, new method for: VILKS	2	72-73
SLESSOR, D.K., Benthonic foraminiferal ecology in Covehead Bay, Prince Edward	2	10 64
Island - A preliminary study	2	48-64
SORENSEN, F.H., see KRAVITZ, J.H.	1 3	17-20
Sound, Harrington, continuous seismic survey of Bermuda, Part II: GEES	2	118-120
Sources, pollution, faunal distribution patterns near, Part 1 of Studies of benthonic		
foraminifora in the Bostigoushe Estuary, SCUARER	2	121 - 134
foraminifera in the Restigouche Estuary: SCHAFER	3	121 -134
STEHMAN, C.F., Eocene deep water sediment from the northeast Providence Channel,		
STEHMAN, C.F., Eocene deep water sediment from the northeast Providence Channel, Bahamas	3 2	121 -134 65-67
STEHMAN, C.F., Eocene deep water sediment from the northeast Providence Channel, Bahamas St. Lawrence Estuary, Quebec, exotic pebbles in Quaternary deposits from the south	2	65-67
STEHMAN, C.F., Eocene deep water sediment from the northeast Providence Channel, Bahamas St. Lawrence Estuary, Quebec, exotic pebbles in Quaternary deposits from the south coast of: DIONNE		
STEHMAN, C.F., Eocene deep water sediment from the northeast Providence Channel, Bahamas St. Lawrence Estuary, Quebec, exotic pebbles in Quaternary deposits from the south coast of: DIONNE St. Lawrence, Gulf of, northwest, distribution of modern benthonic foraminifera in	2 3	65-67 110-112
<pre>STEHMAN, C.F., Eocene deep water sediment from the northeast Providence Channel, Bahamas St. Lawrence Estuary, Quebec, exotic pebbles in Quaternary deposits from the south coast of: DIONNE St. Lawrence, Gulf of, northwest, distribution of modern benthonic foraminifera in the: HOOPER</pre>	2 3 2	65-67
<pre>STEHMAN, C.F., Eocene deep water sediment from the northeast Providence Channel, Bahamas St. Lawrence Estuary, Quebec, exotic pebbles in Quaternary deposits from the south coast of: DIONNE St. Lawrence, Gulf of, northwest, distribution of modern benthonic foraminifera in the: HOOPER Storm surge, shoreline orientation and: HOBBS</pre>	2 3 2	65-67 110-112 74-78
<pre>STEHMAN, C.F., Eocene deep water sediment from the northeast Providence Channel, Bahamas St. Lawrence Estuary, Quebec, exotic pebbles in Quaternary deposits from the south coast of: DIONNE St. Lawrence, Gulf of, northwest, distribution of modern benthonic foraminifera in the: HOOPER Storm surge, shoreline orientation and: HOBBS Structure, diapiric, near Sable Island - Nova Scotia: KING</pre>	2 3 2 3 1	65-67 110-112 74-78 113-115
<pre>STEHMAN, C.F., Eocene deep water sediment from the northeast Providence Channel, Bahamas St. Lawrence Estuary, Quebec, exotic pebbles in Quaternary deposits from the south coast of: DIONNE St. Lawrence, Gulf of, northwest, distribution of modern benthonic foraminifera in the: HOOPER Storm surge, shoreline orientation and: HOBBS</pre>	2 3 2 3 1	65-67 110-112 74-78 113-115
<pre>STEHMAN, C.F., Eocene deep water sediment from the northeast Providence Channel, Bahamas St. Lawrence Estuary, Quebec, exotic pebbles in Quaternary deposits from the south coast of: DIONNE St. Lawrence, Gulf of, northwest, distribution of modern benthonic foraminifera in the: HOOPER Storm surge, shoreline orientation and: HOBBS Structure, diapiric, near Sable Island - Nova Scotia: KING Studies, benthonic foraminifera of, in the Restigouche Estuary: 1. Faunal distribution</pre>	2 3 2 3 1	65-67 110-112 74-78 113-115 1-4
<pre>STEHMAN, C.F., Eocene deep water sediment from the northeast Providence Channel, Bahamas St. Lawrence Estuary, Quebec, exotic pebbles in Quaternary deposits from the south coast of: DIONNE St. Lawrence, Gulf of, northwest, distribution of modern benthonic foraminifera in the: HOOPER Storm surge, shoreline orientation and: HOBBS Structure, diapiric, near Sable Island - Nova Scotia: KING Studies, benthonic foraminifera of, in the Restigouche Estuary: 1. Faunal distribution patterns near pollution sources: SCHAFER</pre>	2 3 2 3 1 3	65-67 110-112 74-78 113-115 1-4 12 ¹ -134
<pre>STEHMAN, C.F., Eocene deep water sediment from the northeast Providence Channel, Bahamas St. Lawrence Estuary, Quebec, exotic pebbles in Quaternary deposits from the south coast of: DIONNE St. Lawrence, Gulf of, northwest, distribution of modern benthonic foraminifera in the: HOOPER Storm surge, shoreline orientation and: HOBBS Structure, diapiric, near Sable Island - Nova Scotia: KING Studies, benthonic foraminifera of, in the Restigouche Estuary: 1. Faunal distribution patterns near pollution sources: SCHAFER Studies, laboratory, on particle abrasion, small circular flume for: GEES</pre>	2 3 2 3 1 3 1 2	65-67 110-112 74-78 113-115 1-4 121-134 15-16
<pre>STEHMAN, C.F., Eocene deep water sediment from the northeast Providence Channel, Bahamas St. Lawrence Estuary, Quebec, exotic pebbles in Quaternary deposits from the south coast of: DIONNE St. Lawrence, Gulf of, northwest, distribution of modern benthonic foraminifera in the: HOOPER Storm surge, shoreline orientation and: HOBBS Structure, diapiric, near Sable Island - Nova Scotia: KING Studies, benthonic foraminifera of, in the Restigouche Estuary: 1. Faunal distribution patterns near pollution sources: SCHAFER Studies, laboratory, on particle abrasion, small circular flume for: GEES Study, microscope, an electron-clay flake orientation in flocculated illite O'BRIEN</pre>	2 3 2 3 1 1 3 1	65-67 110-112 74-78 113-115 1-4 121-134 15-16
<pre>STEHMAN, C.F., Eocene deep water sediment from the northeast Providence Channel, Bahamas St. Lawrence Estuary, Quebec, exotic pebbles in Quaternary deposits from the south coast of: DIONNE St. Lawrence, Gulf of, northwest, distribution of modern benthonic foraminifera in the: HOOPER Storm surge, shoreline orientation and: HOBBS Structure, diapiric, near Sable Island - Nova Scotia: KING Studies, benthonic foraminifera of, in the Restigouche Estuary: 1. Faunal distribution patterns near pollution sources: SCHAFER Studies, laboratory, on particle abrasion, small circular flume for: GEES Study, microscope, an electron-clay flake orientation in flocculated illite O'BRIEN Submersible Shelf Diver, Bay of Fundy, Nova Scotia, sedimentological sampling and results from the diver lock-out facility of: PELLETIER Surge, storm, shoreline orientation and: HOBBS</pre>	2 3 2 3 1 3 1 2	65-67 110-112 74-78 113-115 1-4 121-134 15-16 79-80
<pre>STEHMAN, C.F., Eocene deep water sediment from the northeast Providence Channel, Bahamas St. Lawrence Estuary, Quebec, exotic pebbles in Quaternary deposits from the south coast of: DIONNE St. Lawrence, Gulf of, northwest, distribution of modern benthonic foraminifera in the: HOOPER Storm surge, shoreline orientation and: HOBBS Structure, diapiric, near Sable Island - Nova Scotia: KING Studies, benthonic foraminifera of, in the Restigouche Estuary: 1. Faunal distribution patterns near pollution sources: SCHAFER Studies, laboratory, on particle abrasion, small circular flume for: GEES Study, microscope, an electron-clay flake orientation in flocculated illite O'BRIEN Submersible Shelf Diver, Bay of Fundy, Nova Scotia, sedimentological sampling and results from the diver lock-out facility of: PELLETIER Surge, storm, shoreline orientation and: HOBBS Survey, reconnaissance, sedimentological, of Kane Basin: KRAVITZ</pre>	2 3 2 3 1 3 1 2 3 1 2 3 1	65-67 110-112 74-78 113-115 1-4 121-134 15-16 79-80 102-109 113-115 17-20
<pre>STEHMAN, C.F., Eocene deep water sediment from the northeast Providence Channel, Bahamas St. Lawrence Estuary, Quebec, exotic pebbles in Quaternary deposits from the south coast of: DIONNE St. Lawrence, Gulf of, northwest, distribution of modern benthonic foraminifera in the: HOOPER Storm surge, shoreline orientation and: HOBBS Structure, diapiric, near Sable Island - Nova Scotia: KING Studies, benthonic foraminifera of, in the Restigouche Estuary: 1. Faunal distribution patterns near pollution sources: SCHAFER Studies, laboratory, on particle abrasion, small circular flume for: GEES Study, microscope, an electron-clay flake orientation in flocculated illite O'BRIEN Submersible Shelf Diver, Bay of Fundy, Nova Scotia, sedimentological sampling and results from the diver lock-out facility of: PELLETIER Surge, storm, shoreline orientation and: HOBBS Survey, reconnaissance, sedimentological, of Kane Basin: KRAVITZ Survey, seismic, continuous, of the Bermuda Platform Part I: Castle Harbour: GEES</pre>	2 3 2 3 1 3 1 2 3 1 1	65-67 110-112 74-78 113-115 1-4 121-134 15-16 79-80 102-109 113-115 17-20 21-25
<pre>STEHMAN, C.F., Eocene deep water sediment from the northeast Providence Channel, Bahamas St. Lawrence Estuary, Quebec, exotic pebbles in Quaternary deposits from the south coast of: DIONNE St. Lawrence, Gulf of, northwest, distribution of modern benthonic foraminifera in the: HOOPER Storm surge, shoreline orientation and: HOBBS Structure, diapiric, near Sable Island - Nova Scotia: KING Studies, benthonic foraminifera of, in the Restigouche Estuary: 1. Faunal distribution patterns near pollution sources: SCHAFER Studies, laboratory, on particle abrasion, small circular flume for: GEES Study, microscope, an electron-clay flake orientation in flocculated illite O'BRIEN Submersible Shelf Diver, Bay of Fundy, Nova Scotia, sedimentological sampling and results from the diver lock-out facility of: PELLETIER Surge, storm, shoreline orientation and: HOBBS Survey, reconnaissance, sedimentological, of Kane Basin: KRAVITZ Survey, seismic, continuous, of the Bermuda Platform Part I: Castle Harbour: GEES Survey, seismic continuous of Bermuda, Part II, Harrington Sound: GEES</pre>	2 3 2 3 1 3 1 2 3 1 2 3 1	65-67 110-112 74-78 113-115 1-4 121-134 15-16 79-80 102-109 113-115 17-20
<pre>STEHMAN, C.F., Eocene deep water sediment from the northeast Providence Channel, Bahamas St. Lawrence Estuary, Quebec, exotic pebbles in Quaternary deposits from the south coast of: DIONNE St. Lawrence, Gulf of, northwest, distribution of modern benthonic foraminifera in the: HOOPER Storm surge, shoreline orientation and: HOBBS Structure, diapiric, near Sable Island - Nova Scotia: KING Studies, benthonic foraminifera of, in the Restigouche Estuary: 1. Faunal distribution patterns near pollution sources: SCHAFER Studies, laboratory, on particle abrasion, small circular flume for: GEES Study, microscope, an electron-clay flake orientation in flocculated illite O'BRIEN Submersible Shelf Diver, Bay of Fundy, Nova Scotia, sedimentological sampling and results from the diver lock-out facility of: PELLETIER Surge, storm, shoreline orientation and: HOBBS Survey, reconnaissance, sedimentological, of Kane Basin: KRAVITZ Survey, seismic, continuous, of the Bermuda Platform Part I: Castle Harbour: GEES Survey, seismic continuous of Bermuda, Part II, Harrington Sound: GEES SWIFT, D.J.P., SHIDELER, G.L., AVIGNONE, N.F., HOLLIDAY, B.W., DILL, C.E. Jr.,</pre>	2 3 2 3 1 3 1 2 3 1 1	65-67 110-112 74-78 113-115 1-4 121-134 15-16 79-80 102-109 113-115 17-20 21-25
<pre>STEHMAN, C.F., Eocene deep water sediment from the northeast Providence Channel, Bahamas St. Lawrence Estuary, Quebec, exotic pebbles in Quaternary deposits from the south coast of: DIONNE St. Lawrence, Gulf of, northwest, distribution of modern benthonic foraminifera in the: HOOPER Storm surge, shoreline orientation and: HOBBS Structure, diapiric, near Sable Island - Nova Scotia: KING Studies, benthonic foraminifera of, in the Restigouche Estuary: 1. Faunal distribution patterns near pollution sources: SCHAFER Studies, laboratory, on particle abrasion, small circular flume for: GEES Study, microscope, an electron-clay flake orientation in flocculated illite O'BRIEN Submersible Shelf Diver, Bay of Fundy, Nova Scotia, sedimentological sampling and results from the diver lock-out facility of: PELLETIER Surge, storm, shoreline orientation and: HOBBS Survey, reconnaissance, sedimentological, of Kane Basin: KRAVITZ Survey, seismic, continuous, of the Bermuda Platform Part I: Castle Harbour: GEES Survey, seismic continuous of Bermuda, Part II, Harrington Sound: GEES SWIFT, D.J.P., SHIDELER, G.L., AVIGNONE, N.F., HOLLIDAY, B.W., DILL, C.E. Jr., Quaternary sedimentation on the inner Atlantic shelf between Cape Henry and</pre>	2 3 2 3 1 3 1 2 3 3 1 2 3 1 1 3	65-67 110-112 74-78 113-115 1-4 121-134 15-16 79-80 102-109 113-115 17-20 21-25 118-120
<pre>STEHMAN, C.F., Eocene deep water sediment from the northeast Providence Channel, Bahamas St. Lawrence Estuary, Quebec, exotic pebbles in Quaternary deposits from the south coast of: DIONNE St. Lawrence, Gulf of, northwest, distribution of modern benthonic foraminifera in the: HOOPER Storm surge, shoreline orientation and: HOBBS Structure, diapiric, near Sable Island - Nova Scotia: KING Studies, benthonic foraminifera of, in the Restigouche Estuary: 1. Faunal distribution patterns near pollution sources: SCHAFER Studies, laboratory, on particle abrasion, small circular flume for: GEES Study, microscope, an electron-clay flake orientation in flocculated illite O'BRIEN Submersible Shelf Diver, Bay of Fundy, Nova Scotia, sedimentological sampling and results from the diver lock-out facility of: PELLETIER Surge, storm, shoreline orientation and: HOBBS Survey, reconnaissance, sedimentological, of Kane Basin: KRAVITZ Survey, seismic, continuous, of the Bermuda Platform Part I: Castle Harbour: GEES Survey, seismic continuous of Bermuda, Part II, Harrington Sound: GEES SWIFT, D.J.P., SHIDELER, G.L., AVIGNONE, N.F., HOLLIDAY, B.W., DILL, C.E. Jr.,</pre>	2 3 2 3 1 3 1 2 3 1 1	65-67 110-112 74-78 113-115 1-4 121-134 15-16 79-80 102-109 113-115 17-20 21-25
<pre>STEHMAN, C.F., Eocene deep water sediment from the northeast Providence Channel, Bahamas St. Lawrence Estuary, Quebec, exotic pebbles in Quaternary deposits from the south coast of: DIONNE St. Lawrence, Gulf of, northwest, distribution of modern benthonic foraminifera in the: HOOPER Storm surge, shoreline orientation and: HOBBS Structure, diapiric, near Sable Island - Nova Scotia: KING Studies, benthonic foraminifera of, in the Restigouche Estuary: 1. Faunal distribution patterns near pollution sources: SCHAFER Studies, laboratory, on particle abrasion, small circular flume for: GEES Study, microscope, an electron-clay flake orientation in flocculated illite O'BRIEN Submersible Shelf Diver, Bay of Fundy, Nova Scotia, sedimentological sampling and results from the diver lock-out facility of: PELLETIER Surge, storm, shoreline orientation and: HOBBS Survey, reconnaissance, sedimentological, of Kane Basin: KRAVITZ Survey, seismic, continuous, of the Bermuda Platform Part I: Castle Harbour: GEES Survey, seismic continuous of Bermuda, Part II, Harrington Sound: GEES SWIFT, D.J.P., SHIDELER, G.L., AVIGNONE, N.F., HOLLIDAY, B.W., DILL, C.E. Jr., Quaternary sedimentation on the inner Atlantic shelf between Cape Henry and Cape Hatteras</pre>	2 3 2 3 1 3 1 2 3 3 1 1 3 1 1 3	65-67 110-112 74-78 113-115 1-4 121-134 15-16 79-80 102-109 113-115 17-20 21-25 118-120 5-11
 STEHMAN, C.F., Eocene deep water sediment from the northeast Providence Channel, Bahamas St. Lawrence Estuary, Quebec, exotic pebbles in Quaternary deposits from the south coast of: DIONNE St. Lawrence, Gulf of, northwest, distribution of modern benthonic foraminifera in the: HOOPER Storm surge, shoreline orientation and: HOBBS Structure, diapiric, near Sable Island - Nova Scotia: KING Studies, benthonic foraminifera of, in the Restigouche Estuary: 1. Faunal distribution patterns near pollution sources: SCHAFER Studies, laboratory, on particle abrasion, small circular flume for: GEES Study, microscope, an electron-clay flake orientation in flocculated illite O'BRIEN Submersible Shelf Diver, Bay of Fundy, Nova Scotia, sedimentological sampling and results from the diver lock-out facility of: PELLETIER Surge, storm, shoreline orientation and: HOBBS Survey, reconnaissance, sedimentological, of Kane Basin: KRAVITZ Survey, seismic continuous of Bermuda, Part II, Harrington Sound: GEES SWIFT, D.J.P., SHIDELER, G.L., AVIGNONE, N.F., HOLLIDAY, B.W., DILL, C.E. Jr., Quaternary sedimentation on the inner Atlantic shelf between Cape Henry and Cape Hatteras Tidal bar, and channel, river deposits Gaspé Nord, Quebec: COSTELLO 	2 3 2 3 1 3 1 2 3 3 1 1 3 1 2 3 1 1 2	65-67 110-112 74-78 113-115 1-4 121-134 15-16 79-80 102-109 113-115 17-20 21-25 118-120 5-11 68-71
<pre>STEHMAN, C.F., Eocene deep water sediment from the northeast Providence Channel, Bahamas St. Lawrence Estuary, Quebec, exotic pebbles in Quaternary deposits from the south coast of: DIONNE St. Lawrence, Gulf of, northwest, distribution of modern benthonic foraminifera in the: HOOPER Storm surge, shoreline orientation and: HOBBS Structure, diapiric, near Sable Island - Nova Scotia: KING Studies, benthonic foraminifera of, in the Restigouche Estuary: 1. Faunal distribution patterns near pollution sources: SCHAFER Studies, laboratory, on particle abrasion, small circular flume for: GEES Study, microscope, an electron-clay flake orientation in flocculated illite O'BRIEN Submersible Shelf Diver, Bay of Fundy, Nova Scotia, sedimentological sampling and results from the diver lock-out facility of: PELLETIER Surge, storm, shoreline orientation and: HOBBS Survey, reconnaissance, sedimentological, of Kane Basin: KRAVITZ Survey, seismic, continuous, of the Bermuda Platform Part I: Castle Harbour: GEES Survey, seismic continuous of Bermuda, Part II, Harrington Sound: GEES SWIFT, D.J.P., SHIDELER, G.L., AVIGNONE, N.F., HOLLIDAY, B.W., DILL, C.E. Jr., Quaternary sedimentation on the inner Atlantic shelf between Cape Henry and Cape Hatteras</pre>	2 3 2 3 1 3 1 2 3 3 1 1 3 1 1 3	65-67 110-112 74-78 113-115 1-4 121-134 15-16 79-80 102-109 113-115 17-20 21-25 118-120 5-11
 STEHMAN, C.F., Eocene deep water sediment from the northeast Providence Channel, Bahamas St. Lawrence Estuary, Quebec, exotic pebbles in Quaternary deposits from the south coast of: DIONNE St. Lawrence, Gulf of, northwest, distribution of modern benthonic foraminifera in the: HOOPER Storm surge, shoreline orientation and: HOBBS Structure, diapiric, near Sable Island - Nova Scotia: KING Studies, benthonic foraminifera of, in the Restigouche Estuary: 1. Faunal distribution patterns near pollution sources: SCHAFER Studies, laboratory, on particle abrasion, small circular flume for: GEES Study, microscope, an electron-clay flake orientation in flocculated illite O'BRIEN Submersible Shelf Diver, Bay of Fundy, Nova Scotia, sedimentological sampling and results from the diver lock-out facility of: PELLETIER Surge, storm, shoreline orientation and: HOBBS Survey, reconnaissance, sedimentological, of Kane Basin: KRAVITZ Survey, seismic continuous of Bermuda, Part II, Harrington Sound: GEES SWIFT, D.J.P., SHIDELER, G.L., AVIGNONE, N.F., HOLLIDAY, B.W., DILL, C.E. Jr., Quaternary sedimentation on the inner Atlantic shelf between Cape Henry and Cape Hatteras Tidal bar, and channel, river deposits Gaspé Nord, Quebec: COSTELLO 	2 3 2 3 1 3 1 2 3 3 1 1 3 1 2 3 1 1 2	65-67 110-112 74-78 113-115 1-4 121-134 15-16 79-80 102-109 113-115 17-20 21-25 118-120 5-11 68-71
STEHMAN, C.F., Eocene deep water sediment from the northeast Providence Channel, Bahamas St. Lawrence Estuary, Quebec, exotic pebbles in Quaternary deposits from the south coast of: DIONNE St. Lawrence, Gulf of, northwest, distribution of modern benthonic foraminifera in the: HOOPER Storm surge, shoreline orientation and: HOBBS Structure, diapiric, near Sable Island - Nova Scotia: KING Studies, benthonic foraminifera of, in the Restiguche Estuary: 1. Faunal distribution patterns near pollution sources: SCHAFER Studies, laboratory, on particle abrasion, small circular flume for: GEES Study, microscope, an electron-clay flake orientation in flocculated illite O'BRIEN Submersible Shelf Diver, Bay of Fundy, Nova Scotia, sedimentological sampling and results from the diver lock-out facility of: PELLETIER Surge, storm, shoreline orientation and: HOBBS Survey, reconnaissance, sedimentological, of Kane Basin: KRAVITZ Survey, seismic, continuous, of the Bermuda Platform Part I: Castle Harbour: GEES SWIFT, D.J.P., SHIDELER, G.L., AVIGNONE, N.F., HOLLIDAY, B.W., DILL, C.E. Jr., Quaternary sedimentation on the inner Atlantic shelf between Cape Henry and Cape Hatteras Tidal bar, and channel, river deposits Gaspé Nord, Quebec: COSTELLO Transgressive, basal, sands chiefly inlet-filling sands? Are: KUMAR	2 3 2 3 1 2 3 1 2 3 3 1 1 3 1 2 1 2 1	65-67 110-112 74-78 113-115 1-4 121-134 15-16 79-80 102-109 113-115 17-20 21-25 118-120 5-11 68-71 12-14
 STEHMAN, C.F., Eocene deep water sediment from the northeast Providence Channel, Bahamas St. Lawrence Estuary, Quebec, exotic pebbles in Quaternary deposits from the south coast of: DIONNE St. Lawrence, Gulf of, northwest, distribution of modern benthonic foraminifera in the: HOOPER Storm surge, shoreline orientation and: HOBBS Structure, diapiric, near Sable Island - Nova Scotia: KING Studies, benthonic foraminifera of, in the Restigouche Estuary: 1. Faunal distribution patterns near pollution sources: SCHAFER Studies, laboratory, on particle abrasion, small circular flume for: GEES Study, microscope, an electron-clay flake orientation in flocculated illite O'BRIEN Submersible Shelf Diver, Bay of Fundy, Nova Scotia, sedimentological sampling and results from the diver lock-out facility of: PELLETIER Surge, storm, shoreline orientation and: HOBBS Survey, reconnaissance, sedimentological, of Kane Basin: KRAVITZ Survey, seismic continuous of Bermuda, Part II, Harrington Sound: GEES SWIFT, D.J.P., SHIDELER, G.L., AVIGNONE, N.F., HOLLIDAY, B.W., DILL, C.E. Jr., Quaternary sedimentation on the inner Atlantic shelf between Cape Henry and Cape Hatteras Tidal bar, and channel, river deposits Gaspé Nord, Quebec: COSTELLO 	2 3 2 3 1 3 1 2 3 3 1 1 3 1 2 3 1 1 2	65-67 110-112 74-78 113-115 1-4 121-134 15-16 79-80 102-109 113-115 17-20 21-25 118-120 5-11 68-71
 STEHMAN, C.F., Eocene deep water sediment from the northeast Providence Channel, Bahamas St. Lawrence Estuary, Quebec, exotic pebbles in Quaternary deposits from the south coast of: DIONNE St. Lawrence, Gulf of, northwest, distribution of modern benthonic foraminifera in the: HOOPER Storm surge, shoreline orientation and: HOBBS Structure, diapiric, near Sable Island - Nova Scotia: KING Studies, benthonic foraminifera of, in the Restigouche Estuary: 1. Faunal distribution patterns near pollution sources: SCHAFER Studies, laboratory, on particle abrasion, small circular flume for: GEES Study, microscope, an electron-clay flake orientation in flocculated illite O'BRIEN Submersible Shelf Diver, Bay of Fundy, Nova Scotia, sedimentological sampling and results from the diver lock-out facility of: PELLETIER Surge, storm, shoreline orientation and: HOBBS Survey, reconnaissance, sedimentological, of Kane Basin: KRAVITZ Survey, seismic, continuous, of the Bermuda Platform Part I: Castle Harbour: GEES SWIFT, D.J.P., SHIDELER, G.L., AVIGNONE, N.F., HOLLIDAY, B.W., DILL, C.E. Jr., Quaternary sedimentation on the inner Atlantic shelf between Cape Henry and Cape Hatteras Tidal bar, and channel, river deposits Gaspé Nord, Quebec: COSTELLO Transgressive, basal, sands chiefly inlet-filling sands? Are: KUMAR VILKS, G., New method for separation of skeletal material from a plankton sample 	2 3 2 3 1 2 3 1 2 3 3 1 1 3 1 2 1 2 1	65-67 110-112 74-78 113-115 1-4 121-134 15-16 79-80 102-109 113-115 17-20 21-25 118-120 5-11 68-71 12-14
<pre>STEHMAN, C.F., Eocene deep water sediment from the northeast Providence Channel, Bahamas</pre> St. Lawrence Estuary, Quebec, exotic pebbles in Quaternary deposits from the south coast of: DIONNESt. Lawrence, Gulf of, northwest, distribution of modern benthonic foraminifera in the: HOOPERStorm surge, shoreline orientation and: HOBBS Structure, diapiric, near Sable Island - Nova Scotia: KING Studies, benthonic foraminifera of, in the Restigouche Estuary: 1. Faunal distribution patterns near pollution sources: SCHAPERStudies, laboratory, on particle abrasion, small circular flume for: GEES Study, microscope, an electron-clay flake orientation in flocculated illite O'BRIEN Submersible Shelf Diver, Bay of Fundy, Nova Scotia, sedimentological sampling and results from the diver lock-out facility of: PELLETIER Surge, storm, shoreline orientation and: HOBBS Survey, reconnaissance, sedimentological, of Kane Basin: KRAVITZ Survey, seismic, continuous, of the Bermuda Platform Part I: Castle Harbour: GEES SWIFT, D.J.P., SHIDELER, G.L., AVIGNONE, N.F., HOLLIDAY, B.W., DILL, C.E. Jr., Quaternary sedimentation on the inner Atlantic shelf between Cape Henry and Cape HatterasTidal bar, and channel, river deposits Gaspé Nord, Quebec: COSTELLO Transgressive, basal, sands chiefly inlet-filling sands? Are: KUMARVILKS, G., New method for separation of skeletal material from a plankton sample Waters, sediments, organisms and, in New York Bight and vicinity, integrated	2 3 2 3 1 2 3 1 2 3 3 1 1 3 1 2 1 2 1	65-67 110-112 74-78 113-115 1-4 121-134 15-16 79-80 102-109 113-115 17-20 21-25 118-120 5-11 68-71 12-14
 STEHMAN, C.F., Eocene deep water sediment from the northeast Providence Channel, Bahamas St. Lawrence Estuary, Quebec, exotic pebbles in Quaternary deposits from the south coast of: DIONNE St. Lawrence, Gulf of, northwest, distribution of modern benthonic foraminifera in the: HOOPER Storm surge, shoreline orientation and: HOBBS Structure, diapiric, near Sable Island - Nova Scotia: KING Studies, benthonic foraminifera of, in the Restigouche Estuary: 1. Faunal distribution patterns near pollution sources: SCHAFER Studies, laboratory, on particle abrasion, small circular flume for: GEES Study, microscope, an electron-clay flake orientation in flocculated illite O'BRIEN Submersible Shelf Diver, Bay of Fundy, Nova Scotia, sedimentological sampling and results from the diver lock-out facility of: PELLETIER Surge, storm, shoreline orientation and: HOBBS Survey, reconnaissance, sedimentological, of Kane Basin: KRAVITZ Survey, seismic, continuous, of the Bermuda Platform Part I: Castle Harbour: GEES SWIFT, D.J.P., SHIDELER, G.L., AVIGNONE, N.F., HOLLIDAY, B.W., DILL, C.E. Jr., Quaternary sedimentation on the inner Atlantic shelf between Cape Henry and Cape Hatteras Tidal bar, and channel, river deposits Gaspé Nord, Quebec: COSTELLO Transgressive, basal, sands chiefly inlet-filling sands? Are: KUMAR VILKS, G., New method for separation of skeletal material from a plankton sample 	2 3 2 3 1 2 3 1 2 3 3 1 1 3 1 2 1 2 1	65-67 110-112 74-78 113-115 1-4 121-134 15-16 79-80 102-109 113-115 17-20 21-25 118-120 5-11 68-71 12-14
<pre>STEHMAN, C.F., Eocene deep water sediment from the northeast Providence Channel, Bahamas St. Lawrence Estuary, Quebec, exotic pebbles in Quaternary deposits from the south coast of: DIONNE St. Lawrence, Gulf of, northwest, distribution of modern benthonic foraminifera in the: HOOPER Storm surge, shoreline orientation and: HOBBS Structure, diapiric, near Sable Island - Nova Scotia: KING Studies, benthonic foraminifera of, in the Restigouche Estuary: 1. Faunal distribution patterns near pollution sources: SCHAFER Studies, laboratory, on particle abrasion, small circular flume for: GEES Study, microscope, an electron-clay flake orientation in flocculated illite O'BRIEN Submersible Shelf Diver, Bay of Fundy, Nova Scotia, sedimentological sampling and results from the diver lock-out facility of: PELLETIER Surge, storm, shoreline orientation and: HOBBS Survey, reconnaissance, sedimentological, of Kane Basin: KRAVITZ Survey, seismic, continuous, of the Bermuda Platform Part I: Castle Harbour: GEES SWIFT, D.J.P., SHIDELER, G.L., AVIGNONE, N.F., HOLLIDAY, B.W., DILL, C.E. Jr., Quaternary sedimentation on the inner Atlantic shelf between Cape Henry and Cape Hatteras Tidal bar, and channel, river deposits Gaspé Nord, Quebec: COSTELLO Transgressive, basal, sands chiefly inlet-filling sands? Are: KUMAR VILKS, G., New method for separation of skeletal material from a plankton sample Waters, sediments, organisms and, in New York Bight and vicinity, integrated continental-shelf and marginal marine environmental research program:</pre>	2 3 2 3 1 3 1 2 3 3 1 1 3 1 2 1 2	65-67 110-112 74-78 113-115 1-4 121-134 15-16 79-80 102-109 113-115 17-20 21-25 118-120 5-11 68-71 12-14 72-73