Index

Citation refers to Issue Number and Page Number

Number Page

Anthropical strauctures in coastal sediments: LEATHERMAN	2	69
Barrier Island morphology, changing, Kouchibouguac Bay, New Brunswick, wave climate effects upon: BRYANT Bay, Buctouche, the, complex, physical	2	49
parameter distribution patterns in bottom sediments of: THIBAULT Bay, Kouchibouguac, New Brunswick,	3	95
wave climate effects upon changing barrier island morphology: BRYANT Beach deposits, ice-formed, degrad-	2	49
ation of: ROSEN Bottom sediments of the Buctouche Bay	2	63
complex, physical parameter distri- bution patterns in: THIBAULT Brunswick, New, Wave climate effects	3	95
ology, Kouchibouguac Bay: BRYANT BRYANT, E., Wave climate effects upon	2	49
changing barrier island morphology, Kouchibouguac Bay, New Brunswick Buctouche Bay, the, complex, physical parameter distribution patterns in	2	49
bottom sediments of: THIBAULT	3	95
Canadian, eastern, Holocene foramini- fera, magnesium in 19 species of: HOOPER Changing barrier island morphology	2	73
Kouchibouguac Bay, New Brunswick, wave climate effects upon: BRYANT	2	49
refuted: SMITH	l	11
barrier island morphology, Kouchi- bouguac Bay, New Brunswick: BRYANT Coastal sediments, anthropical	2	49
structures in: LEATHERMAN Complex, Buctouche Bay, the, physical parameter distribution patterns	2	69
in bottom sediments of: THIBAULT	3	95
Degradation of ice-formed beach deposit ROSEN	s: 2	63
ation of: ROSEN	2	63
of Sherbrooke, Nova Scotia: HOOPER Distribution patterns, parameter,	3	83
the Buctouche Bay complex: THIBAULT	3	95
Eastern Canadian Holocene foraminifera, magnesium in 19 species of: HOOPER	2	73
barrier island morphology, Kouchi- bouguac Bay, New Brunswick: BRYANT	2	49
Fiord, Pangnirtung, observations on oceanography and sedimentation at; GILBERT	1	1
Foraminifera, Holocene, Canadian, eastern, magnesium in 19 species of: HOOPER	2	73

	Number	Page
Foraminiferida south of Sherbrooke,		
Nova Scotia, distribution of: HOOPER	3	83
Formation, Presumpscot, the, Portland, Maine, Late Wisconsinan wood and other tree remains in: HYLAND	3	103
CILBERT D. Obcomunitions on occorregener	.h	
and sedimentation at Pangnirtung	ony	
Fiord, Baffin Island.	1	1
a claim of: SMITH	1	11
High level granitization refuted, a		
Claim of: SMITH Holocene foraminifera, Canadian, easter	'n	11
magnesium in 19 species of: HOOPER HOOPER, K., Distribution of Foramini-	2	73
ferida south of Sherbrooke, Nova Scotia	3	83
HOOPER, K., Magnesium in 19 species of		
eastern Canadian Holocene foraminife HUGHES, C.J. and SMITH, M.M., A claim o	ra 2 f	73
high level granitization refuted HYLAND, F., THOMPSON, W.B. and	1	11
STUCKENRATH, R. JR., Late Wisconsina	n	
Presumpscot Formation, Portland, Mai	ne 3	103
<pre>Ice-formed beach deposits, degradation of: ROSEN Island morphology, barrier, changing,</pre>	2	63
Kouchibouguac Bay, New Brunswick, wave climate effects upon: BRYANT	2	49
Kouchibouguac Bay, New Brunswick, wave climate effects upon changing barrie island morphology: BRYANT	r 2	49
Lamination sampling technique for uncon	- ,	זב
Late Wisconsinan wood and other tree	T	12
Portland, Maine: HYLAND	3	103
LEATHERMAN, S.P., Anthropical structure in coastal sediments	s 2	69
LEATHERMAN, S.P. and WILLIAMS, A.T.,	-	
Lamination sampling technique for un consolidated sediments	- 1	15
Level, high, granitization refuted, a	_	
claim of: SMITH	1	11
Magnesium in 19 species of eastern Canadian Holocene foraminifera: HOO Maine, Late Wisconsinan wood and other	PER 2	73
tree remains in the Presumpscot Formation, Portland: HYLAND	3	103
Margin, outer, Nova Scotian, regional review of: STOW	1	17
Morphology, island, barrier, changing,		
wave climate effects upon: BRYANT	2	49
New Brunswick, wave climate effects upon	n	
changing barrier island morphology, Kouchibouguac Bay: BRYANT	2	49

Number Page

Nova Scotia, distribution of Foramini- ferida south of Sherbrooke:		
HOOPER	3	83
Nova Scotlan outer margin, regional review of: STOW	1	17
Observations on oceanography and sedimentation at Pangnirtung Fiord, Baffin Island: GILBERT	1	1
Oceanography and sedimentation at Pangnirtung Fiord, observations on: GILBERT	1	1
Formation, Portland, Maine, Late Wisconsinan wood and: HYLAND Outer margin, Nova Scotian, regional	3	103
review of: STOW	1	17
Pangnirtung Fiord, observations on oceanography and sedimentation at: GILBERT	1	1
Parameter, physical, distribution patterns in bottom sediments of	-	-
the Buctouche Bay complex: THIBAULT Patterns, distribution, parameter,	3	95
the Buctouche Bay complex: THIBAULT Physical parameter distribution patterns	3	95
in bottom sediments of the Bactouche Bay complex: THIBAULT Portland, Maine, Late Wisconsinan wood	3	95
and other tree remains in the Presump- scot Formation: HYLAND Presumpscot Formation, the, Portland	3	103
Maine, Late Wisconsinan wood and other tree remains in: HYLAND	r 3	-103
Refuted, a claim of high level granit- ization: HUGHES	1	11
Regional review of the Nova Scotlan outer margin: STOW Remains, tree, other, in the Presumpscot	1	17
Formation, Portland, Maine, Late Wisconsinan wood and: HYLAND	3	103
Review, regional, of the Nova Scotian outer margin: STOW ROSEN, P.S., Degradation of ice-	1	17
formed beach deposits.	2	63
Sampling technique, lamination, for unconsolidated sediments: LEATHERMAN Scotian, Nova, outer margin, regional	1	15
review of: STOW	1	17

Sedimentation at Pangnirtung Fiord, observations on oceanography and:		
GILBERT	1	1
Sediments, bottom, of the Buctouche Bay complex, physical parameter distri-		
bution patterns in: THIBAULT	3	95
Sediments, coastal, anthropical structur	es	6.0
in: LEATHERMAN	2	69
Sediments, unconsolidated, lamination	1	1 6
Sherbrooke Nova Scotia distribution of	т	10
Foraminiferida south of. HOOPER	з	83
SMITH, M.M., See HUGHES	ĩ	11
South of Sherbrooke Nova Scotia	-	
distribution of Foraminiferida:		
HOOPER	3	83
Species, 19. of eastern Canadian	5	00
Holocene foraminifera, magnesium in:		
HOOPER	2	73
STOW, A.V.D., Regional review of the		
Nova Scotian outer margin	1	17
Structures, anthropical, in coastal		
sediments: LEATHERMAN	2	69
STUCKENRATH, R., JR., see HYLAND	3	103
Technique, sampling, lamination, for		
unconsolidated sediments: LEATHERMAN	1	15
THIBAULT, J., Physical parameter dis-		
tribution patterns in bottom sediment	S	0.5
of the Buctouche Bay complex	3	95
THOMPSON, W.B., See HYLAND	3	103
Formation Portland Maine Late		
Wisconsinan wood and, HVIAND	2	102
WISCONSINAN WOOD AND. MILAND	5	103
Unconsolidated sediments, lamination		
•		
sampling techniques for:		
sampling techniques for: LEATHERMAN	1	15
sampling techniques for: LEATHERMAN	1	15
sampling techniques for: LEATHERMAN Wave climate effects upon changing	1	15
sampling techniques for: LEATHERMAN Wave climate effects upon changing barrier island morphology, Kouchi-	1	15
sampling techniques for: LEATHERMAN Wave climate effects upon changing barrier island morphology, Kouchi- bouguac Bay, New Brunswick:	l	15
sampling techniques for: LEATHERMAN Wave climate effects upon changing barrier island morphology, Kouchi- bouguac Bay, New Brunswick: BRYANT	1 2	15 49
sampling techniques for: LEATHERMAN Wave climate effects upon changing barrier island morphology, Kouchi- bouguac Bay, New Brunswick: BRYANT WILLIAMS, A.T., see LEATHERMAN	1 2 1	15 49 15
<pre>sampling techniques for: LEATHERMAN Wave climate effects upon changing barrier island morphology, Kouchi- bouguac Bay, New Brunswick: BRYANT WILLIAMS, A.T., see LEATHERMAN Wisconsinan, Late, wood and other tree</pre>	1 2 1	15 49 15
<pre>sampling techniques for: LEATHERMAN Wave climate effects upon changing barrier island morphology, Kouchi- bouguac Bay, New Brunswick: BRYANT WILLIAMS, A.T., see LEATHERMAN Wisconsinan, Late, wood and other tree remains in the Presumpscot Formation,</pre>	1 2 1	15 49 15
<pre>sampling techniques for: LEATHERMAN Wave climate effects upon changing barrier island morphology, Kouchi- bouguac Bay, New Brunswick: BRYANT WILLIAMS, A.T., see LEATHERMAN Wisconsinan, Late, wood and other tree remains in the Presumpscot Formation, Portland, Maine: HYLAND</pre>	1 2 1 3	15 49 15 103
<pre>sampling techniques for: LEATHERMAN Wave climate effects upon changing barrier island morphology, Kouchi- bouguac Bay, New Brunswick: BRYANT WILLIAMS, A.T., see LEATHERMAN Wisconsinan, Late, wood and other tree remains in the Presumpscot Formation, Portland, Maine: HYLAND Wood, Wisconsinan, Late, and other tree maning in the Presumpscot Formation, Portland, Maine: HYLAND</pre>	1 2 1 3	15 49 15 103
<pre>sampling techniques for: LEATHERMAN Wave climate effects upon changing barrier island morphology, Kouchi- bouguac Bay, New Brunswick: BRYANT WILLIAMS, A.T., see LEATHERMAN Wisconsinan, Late, wood and other tree remains in the Presumpscot Formation, Portland, Maine: HYLAND Wood, Wisconsinan, Late, and other tree remains in the Presumpscot Formation, Portland, Maine: HYLAND</pre>	1 2 1 3	15 49 15 103

Number Page