

Index to Reports in Volume 3

Citation refers to Issue Number, and Page Numbers.

Age and origin, Perry Formation, Charlotte County, New Brunswick, Canada: McILWAINÉ	2-3	56-60
Analysis, basin and paleocurrent, Meguma Group, Nova Scotia: CAMPBELL & SCHENK	1	27-30
Arctic waters, sedimentation in, Western Queen Elizabeth Islands, District of Franklin, Canada: PELLETIER	4	90-99
Atlantic Oceanographic Group, marine geological investigations in Gulf of St. Lawrence, 1964-1966: LORING	1	24-26
Basin and paleocurrent analysis, Meguma Group, Nova Scotia: CAMPBELL & SCHENK	1	27-30
Bay of Fundy, Minas Basin, tidal delta and ebb-flood channel system in: SWIFT et al	1	12-16
Bay of Fundy, origin: SWIFT and LYALL	2-3	67-70
Bay of Fundy: reconnaissance by sub-bottom profiler: SWIFT & LYALL	2-3	67-70
Bay of Fundy, structure of Minas Passage: SWIFT et al	4	112-118
Beaver Harbour, N. B., plant fossils: HELMSTAEDT	4	119
Bibliography, geophysical surveys, continental shelf south of Nova Scotia: HOOD	1	6-11
BURNS, H. W., Jr., see SWIFT	1	17-23
Bottom sediment distribution, near-shore, Cape Sable Island, Nova Scotia, Canada: LYALL & GEES	2-3	65-66
Bottom sediment samples, factors influencing benthonic Foraminifera in, Gulf of St. Lawrence: SCHAFER	2-3	71
Bottom sediments, Arctic waters, Queen Elizabeth Islands, District of Franklin, Canada: PELLETIER	4	90-99
Bottom sediments, Grand Banks, distribution of Foraminifera in: SEN GUPTA	2-3	61-63
Bottom sediments, Gulf of St. Lawrence, investigations of: LORING	1	24-26
Bottom sediments, Port au Port Bay, Newfoundland: SHEARER	1	5
CAMPBELL, F. H. A. and SCHENK, P. E., Paleocurrent and basin analysis of the Meguma Group (lower Paleozoic), Nova Scotia	1	27-30
Cape Sable Island, Nova Scotia, Canada, near-shore sediment distribution: LYALL & GEES	2-3	65-66
Carboniferous, lower, plant fossils from, Beaver Harbour, N. S.: HELMSTAEDT	4	119
CARR, P. A., Salt-water intrusion studies in the Maritimes	4	109-111
Channel system, ebb-flood, Minas Basin, Bay of Fundy: SWIFT et al	1	12-16
Clay flake orientation in sensitive Pleistocene clay: O'BRIEN & HARRISON	1	1-4
Continental Shelf, south of Nova Scotia, geophysical surveys on: HOOD	1	6-11
Continuous seismic profiles, Bay of Fundy: SWIFT & LYALL	2-3	67-70
Continuous seismic profile, Halifax Harbour, Nova Scotia: GRANT	2-3	64
Dates, radiocarbon, Tyrrell Sea area: WAGNER	4	100-104
Delta, tidal, Minas Basin, Bay of Fundy: SWIFT et al	1	12-16
Devonian, Perry formation, age and origin, Charlotte County, New Brunswick, Canada: McILWAINÉ	2-3	56-60
Distribution, Foraminifera, Sediments of Grand Banks: SEN GUPTA	2-3	61-63
Distribution, living benthonic Foraminifera, Northumberland Strait: SCHAFER	4	105-108
Distribution, near-shore sediment, Cape Sable Island, Nova Scotia, Canada: LYALL & GEES	2-3	65-66
District of Franklin, Canada, sedimentation in Arctic waters, Queen Elizabeth Islands: PELLETIER	4	90-99
Ebb-flood channel system, Minas Basin, Bay of Fundy: SWIFT et al	1	12-16
Factors influencing numerical accuracy of benthonic Foraminifera population counts, bottom sediment samples, Gulf of St. Lawrence: SCHAFER	2-3	71

Fluviomarine outwash terrace, raised, genesis of, Minas Basin, Nova Scotia:			
SWIFT & BURNS	1	17-23	
Foraminifera, benthonic, distribution in sediments of Grand Banks: SEN GUPTA	2-3	61-63	
Foraminifera, benthonic, factors influencing numerical accuracy of population counts, bottom sediment samples, Gulf of St. Lawrence: SCHAFER	2-3	71	
Foraminifera, living benthonic, distribution, Northumberland Strait: SCHAFER	4	105-108	
Fossils, plant, Beaver Harbour, N. B.: HELMSTAEDT	4	119	
GEES, R. A., see LYALL	2-3	65-66	
Genesis of raised fluviomarine outwash terrace, Minas Basin, Nova Scotia:			
SWIFT & BURNS	1	17-23	
Geochemical investigations of bottom sediments from Gulf of St. Lawrence by Atlantic Oceanographic Group, 1964-1966: LORING	1	24-26	
Geochemical studies of bottom sediments, Arctic waters, Queen Elizabeth Islands, District of Franklin, Canada: PELLETIER	4	90-99	
Geological interpretation from geophysical surveys, continental shelf south of Nova Scotia: HOOD	1	6-11	
Geological, marine, investigations by Atlantic Oceanographic Group, Gulf of St. Lawrence, 1964-1966: LORING	1	24-26	
Geophysical surveys, continental shelf south of Nova Scotia: HOOD	1	6-11	
Grand Banks, distribution of Foraminifera in sediments of: SEN GUPTA	2-3	61-63	
GRANT, A. C., A continuous seismic profile from Halifax Harbour, Nova Scotia	2-3	109-111	
Gulf of St. Lawrence, factors influencing benthonic Foraminifera in bottom sediment samples: SCHAFER	2-3	71	
Gulf of St. Lawrence, marine geological investigations in, Atlantic Oceanographic Group, 1964-1966: LORING	1	24-26	
Halifax Harbour, Nova Scotia, continuous seismic profile from: GRANT	2-3	64	
HARRISON, W., see O'BRIEN	1	1-4	
HELMSTAEDT, H., Note on plant fossils from Beaver Harbour, N. B.	4	119	
HOOD, P., Geophysical surveys on the continental shelf south of Nova Scotia	1	6-11	
Hudson Bay, radiocarbon dates, Tyrrell Sea area: WAGNER	2-3	100-104	
Intrusion studies, salt-water, Maritimes: CARR	4	109-111	
Investigations, marine geological by Atlantic Oceanographic Group, Gulf of St. Lawrence, 1964-1966: LORING	1	24-26	
JAGODITS, F. L., see SWIFT	4	112-118	
Living benthonic Foraminifera, distribution, Northumberland Strait: SCHAFER	4	105-108	
LORING, D. H., Resumé of marine geological investigations carried out by the Atlantic Oceanographic Group in the Gulf of St. Lawrence, 1964-1966	1	24-26	
LYALL, A. K. and GEES, R. A., Near-shore sediment distribution around Cape Sable Island, Nova Scotia, Canada	2-3	65-66	
LYALL, A. K., see SWIFT	1	12-16	
LYALL, A. K., see SWIFT	2-3	67-70	
McILWAINE, W. H., Age and origin of the Perry formation (Devonian), Charlotte County, New Brunswick, Canada	2-3	56-60	
MCMULLEN, R. M., see SWIFT	1	12-16	
MANISTRE, B. L., see SWIFT	4	112-118	
Marine geological investigations by Atlantic Oceanographic Group, Gulf of St. Lawrence, 1964-1966: LORING	1	24-26	
Maritimes, salt-water intrusion studies: CARR	3	109-111	
Meguma Group, paleocurrent and basin analysis of, Nova Scotia: CAMPBELL & SCHENK	1	27-30	
Micrographs, electron, of clay flake orientation in Pleistocene clay: O'BRIEN and HARRISON	1	1-4	
Minas Basin, Nova Scotia, genesis of raised fluviomarine outwash terrace:			
SWIFT & BURNS	1	17-23	
Minas Basin, tidal delta and ebb-flood channel system in: SWIFT et al	1	12-16	
Minas Passage, Bay of Fundy, structure of: SWIFT et al	3	112-118	

Near-shore sediment distribution, Cape Sable Island, Nova Scotia, Canada:		
LYALL & GEES	2-3	65-66
New Brunswick, Bay of Fundy, reconnaissance by sub-bottom profiler: SWIFT and LYALL	2-3	67-70
New Brunswick, Canada, Charlotte County, Perry formation, age and origin: McILWAINE	2-3	56-60
New Brunswick, plant fossils from Beaver Harbour: HELMSTAEDT	4	119
New Brunswick, salt-water intrusion studies: CARR	4	109-111
Newfoundland, bottom sediments from Port au Port Bay: SHEARER	1	5
New York (state), clay flake orientation in Pleistocene clay from: O'BRIEN & HARRISON	1	1-4
Northumberland Strait, distribution of living benthonic Foraminifera in: SCHAFER	4	105-108
Nova Scotia, Bay of Fundy, reconnaissance by sub-bottom profiler: SWIFT and LYALL	2-3	67-70
Nova Scotia, Canada, near-shore sediment distribution, Cape Sable Island: LYALL & GEES	2-3	65-66
Nova Scotia, continental shelf south of, geophysical surveys on: HOOD	1	6-11
Nova Scotia, continuous seismic profile from Halifax Harbour: GRANT	2-3	64
Nova Scotia, Minas Basin, Bay of Fundy, tidal delta and ebb-flood channel system in: SWIFT et al	1	12-16
Nova Scotia, Minas Basin, genesis of raised fluviomarine outwash terrace: SWIFT and BORNS	1	17-23
Nova Scotia, paleocurrent and basin analysis of Meguma Group in: CAMPBELL & SCHENK	1	27-30
Nova Scotia, structure of Minas Passage, Bay of Fundy: SWIFT et al	3	112-118
O'BRIEN, N. R., and HARRISON, W., Clay flake orientation in a sensitive Pleistocene clay	1	1-4
Orientation, clay flake in Pleistocene clay: O'BRIEN & HARRISON	1	1-4
Origin and age, Perry formation, Charlotte County, New Brunswick, Canada: McILWAINE	2-3	56-60
Outwash terrace, raised fluviomarine, genesis of, Minas Basin, Nova Scotia: SWIFT & BORNS	1	17-23
Paleocurrent analysis, Perry formation, Charlotte County, New Brunswick, Canada: McILWAINE	2-3	56-60
Paleocurrent and basin analysis, Meguma Group, Nova Scotia: CAMPBELL & SCHENK	1	27-30
Paleozoic, lower, paleocurrent and basin analysis of Meguma Group, Nova Scotia: CAMPBELL & SCHENK	1	27-30
PATERSON, N. R., see SWIFT	4	112-118
PELLETIER, B. R., Sedimentation in Arctic waters of the western Queen Elizabeth Islands, District of Franklin, Canada	4	90-99
Perry formation, age and origin, Charlotte County, New Brunswick, Canada: McILWAINE	2-3	56-60
Plant fossils, Beaver Harbour, N. B.: HELMSTAEDT	4	119
Pleistocene clay, clay flake orientation in: O'BRIEN & HARRISON	1	1-4
Population counts, benthonic Foraminifera, factors influencing numerical accuracy of, bottom sediment samples, Gulf of St. Lawrence: SCHAFER	2-3	71
Port au Port Bay, Newfoundland, bottom sediments from: SHEARER	1	5
Prince Edward Island, salt-water intrusion studies: CARR	4	109-111
Queen Elizabeth Islands, District of Franklin, Canada, sedimentation in Arctic waters: PELLETIER	4	90-99
Radiocarbon dates, Tyrrell Sea area: WAGNER	4	100-104
Raised fluviomarine outwash terrace, genesis of, Minas Basin, Nova Scotia: SWIFT & BORNS	1	17-23
Salt-water intrusion studies, Maritimes: CARR	4	109-111
SCHAFFER, C. T., Factors influencing the numerical accuracy of benthonic Foraminifera population counts from bottom sediment samples, Gulf of St. Lawrence	2-3	71

SCHAFFER, C. T., Preliminary survey of the distribution of living benthonic Foraminifera in Northumberland Strait	4	105-108
SCHENK, P. E., see CAMPBELL	1	27-30
Sediment distribution, near-shore, Cape Sable Island, Nova Scotia, Canada: LYALL & GEES	2-3	65-66
Sediment samples, bottom, factors influencing benthonic Foraminifera in, Gulf of St. Lawrence: SCHAFFER	2-3	71
Sediments, bottom, Gulf of St. Lawrence, investigations of: LORING	1	24-26
Sediments, bottom, Port au Port Bay, Newfoundland: SHEARER	1	5
Sediments, Grand Banks, distribution of Foraminifera in: SEN GUPTA	2-3	61-63
Sedimentation in Arctic waters, western Queen Elizabeth Islands, District of Franklin, Canada: PELLETIER	4	90-99
Seismic profiles, continuous, Bay of Fundy: SWIFT and LYALL	2-3	67-70
Seismic profile, continuous, Halifax Harbour, Nova Scotia: GRANT	4	64
SEN GUPTA, B. K., Distribution of Foraminifera in the sediments of the Grand Banks: a preliminary report	2-3	61-63
SHEARER, J. M., Bottom sediments of Port au Port Bay, Newfoundland	1	5
Shelf, continental, south of Nova Scotia, geophysical surveys on: HOOD	1	6-11
Stratigraphy, Minas Passage, Bay of Fundy: SWIFT et al	4	112-118
Structure, Bay of Fundy: SWIFT and LYALL	2-3	67-70
Structure, Minas Passage, Bay of Fundy: SWIFT et al	4	112-118
Sub-bottom profiler, reconnaissance by, Bay of Fundy: SWIFT & LYALL	2-3	67-70
Surveys, geophysical, continental shelf south of Nova Scotia: HOOD	1	6-11
SWIFT, D. J. P., McMULLEN, R. M., LYALL, A. K., A tidal delta with an ebb-flood channel system in the Minas Basin, Bay of Fundy: preliminary report	1	12-16
SWIFT, D. J. P. and BURNS, H. W., Jr., Genesis of the raised fluviomarine outwash terrace, north south of the Minas Basin, Nova Scotia; a preliminary report	1	17-23
SWIFT, D. J. P. and LYALL, Anil K., Bay of Fundy: reconnaissance by sub-bottom profiler	2-3	67-70
SWIFT, D. J. P.; JAGODITS, F. L.; MANISTRE, B. L.; and PATERSON, N. R., Structure of the Minas Passage, Bay of Fundy: preliminary report	4	112-118
Terrace, raised fluviomarine outwash, genesis of, Minas Basin, Nova Scotia SWIFT & BURNS	1	17-23
Tidal currents in Minas Basin, Bay of Fundy: SWIFT et al	1	12-16
Tidal delta, Minas Basin, Bay of Fundy: SWIFT et al	1	12-16
Tyrrell Sea area, radiocarbon dates: WAGNER	4	100-104
Uplift rate, Tyrrell Sea area: WAGNER	4	100-104
WAGNER, F. J. E., Additional radiocarbon dates, Tyrrell Sea area	4	100-104
Water, salt, intrusion studies, Maritimes: CARR	4	109-111
Waters, sedimentation in Arctic, western Queen Elizabeth Islands, District of Franklin, Canada: PELLETIER	4	90-99