

Overview of age constraints for gold mineralization in central and western Newfoundland and new $^{40}\text{Ar}/^{39}\text{Ar}$ ages for muscovite from selected auriferous zones

HAMISH A.I. SANDEMAN¹, IAN W. HONSBERGER² AND ALFREDO CAMACHO³

1. Geological Survey Division, Department of Industry, Energy and Technology, Government of Newfoundland and Labrador, St. John's, Newfoundland and Labrador A1B 3J5, Canada
 2. Geological Survey of Canada, Ottawa, Ontario K1A 0E8, Canada
 3. University of Manitoba, Department of Geological Sciences, Winnipeg, Manitoba R3T 2N2, Canada
- *Corresponding author: <hamishsandeman@gov.nl.ca>

Date received: 19 May 2022 † *Date accepted 25 October 2022*

ABSTRACT

This contribution presents new $^{40}\text{Ar}/^{39}\text{Ar}$ laser step-heating data for muscovite associated with four significant orogenic gold-mineralized zones in central and western Newfoundland to build upon existing geochronological constraints for orogenic gold mineralization across much of the Newfoundland Appalachians. Additionally, we provide the first $^{40}\text{Ar}/^{39}\text{Ar}$ laser step-heating data for detrital muscovite from gold-mineralized sandstone of the Wigwam Formation (Botwood Group) at the Moosehead gold prospect. Most known gold zones on the island occur in proximity to reactivated crustal-scale faults and related structures, and are commonly localized within competent, rigid lithological units (e.g., granitoid rocks); although, some vein systems cut less competent, clastic sedimentary rock sequences. Host rocks range in age from Neoproterozoic to Devonian and may extend into the Carboniferous. Robust Pragian to Emsian (ca. 410–375 Ma) rutile, muscovite and zircon ages constrain the timing of gold mineralization in parts of the Exploits Subzone, whereas less precise age estimates for gold mineralization in the Notre Dame Subzone and Laurentian margin range from Wenlock to Emsian (ca. 433–375 Ma). Collectively, the geochronological data suggest that fluid-pressure cycling associated with gold mineralization in central and western Newfoundland occurred during progressive Siluro-Devonian tectonics associated with the waning stages of the Salinic orogenic cycle and spanning the Acadian and Neocadian orogenic cycles. Multiple, polyphase, overlapping orogenic events allowed for repeated, gold mineralizing fluid flow events, particularly in proximity to long-lived, reactivated crustal-scale fault zones.

RÉSUMÉ

La présente contribution fait état de nouvelles données obtenues par chauffage échelonné au laser $^{40}\text{Ar}/^{39}\text{Ar}$ de muscovite associée à quatre zones minéralisées aurifères orogéniques stratégiques dans le centre et l'ouest de Terre-Neuve permettant de mieux préciser les limites géochronologiques existantes de la minéralisation aurifère orogénique dans la majeure partie des Appalaches terre-neuviennes. Nous fournissons de plus les premières données par chauffage échelonné au laser $^{40}\text{Ar}/^{39}\text{Ar}$ de muscovite détritique provenant de grès minéralisé aurifère de la Formation de Wigwam (groupe de Botwood) dans la zone d'intérêt aurifère de Moosehead. La majorité des zones aurifères connues sur l'île se trouve à proximité de failles d'échelle crustale et de structures connexes, et elles se situent communément à l'intérieur d'unités lithologiques rigides parallèles (p. ex. des roches granitiques), bien que certains réseaux filoniens recoupent des séquences de roches sédimentaires clastiques moins parallèles. L'âge des roches hôtes varie du Néoprotérozoïque au Dévonien et peut s'étendre au Carbonifère. Des datations robustes du Praguien à l'Emsien (env. 410 à 375 Ma) de rutile, de muscovite et de zircon limitent le moment de la minéralisation de l'or dans des parties de la sous-zone Exploits, alors que des estimations moins précises de l'âge de la minéralisation aurifère dans la sous-zone Notre-Dame et de la marge laurentienne varient du Wenlock à l'Emsien (env. 433 à 375 Ma). Les données géochronologiques laissent collectivement supposer que les cycles de pression hydraulique associés à la minéralisation d'or dans le centre et l'ouest de Terre-Neuve se sont manifestés durant la progression tectonique siluro-dévonienne associée aux stades de ralentissement du cycle orogénique salinique et s'étendant aux cycles orogéniques acadien et néocadien. Plusieurs phénomènes orogéniques polyphasés se chevauchant ont permis des écoulements de fluides minéralisateurs aurifères répétés, en particulier à proximité de zones de failles d'échelle crustale réactivées persistantes.

[Traduit par la rédaction]

INTRODUCTION

Prior to the late 1980s, the vast majority of gold produced in the Newfoundland Appalachians was derived as a by-product from auriferous volcanic massive sulphide deposits such as the Rambler, Ming and in particular, Buchans deposits of the central Newfoundland Dunnage Zone (see Piercey 2007; Galley *et al* 2007). Positive market conditions and the introduction of flow through shares stimulated an Island-wide gold exploration boom in the late 1980s that led to the discovery of many of the gold deposits that are past producers (Fig. 1) (e.g., Hope Brook Deposit, Big Ridge Resources; Hammerdown and Orion, Maritime Resources) and are currently in production (e.g., Stog'èr Tight deposit, Anaconda Mining). In addition to these deposits, prospective gold deposits and occurrences seeing renewed exploration today include: (1) Valentine Lake Project, Marathon Gold Corporation; (2) Cape Ray Deposits, Matador Mining; (3) Appleton-JBP linears (now Queensway Project), New Found Gold; (4) Little River Prospects (now Golden Baie), Canstar Resources; and (5) Moosehead, Sokoman Minerals.

The geoscientific knowledge base related to gold mineralized zones in central and western Newfoundland is lean compared to other global gold producing districts. Early gold-related studies, subsequent to the explosion of exploration in the late 1980s, provided some of the first descriptive and comparative documentation of the gold occurrences on the Island (e.g., Tuach 1987; Tuach *et al* 1988; Evans 1991, 1993, 1999, 2004; Churchill *et al* 1993; Wilson and Evans 1994). Extant geochronological investigations on gold zones are limited to those from the Hope Brook (Dubé *et al* 1995) and Cape Ray deposits (Dubé *et al* 1996; Dubé and Lauzière 1997) and select geochronological studies at the Nugget Pond (Sangster *et al* 2008) and Hammerdown (Ritcey *et al* 1995) deposits. Subjects of more recent gold-related, paragenetic, litho-geochemical and geochronological studies include: the Rattling Brook (Kerr and van Breemen 2007) and Thor prospects (Minnett *et al* 2012) in western Newfoundland (White Bay); the Pine Cove (Ybarra 2020) and Ming (Au-rich VMS) deposits of Baie Verte Peninsula (Pilote *et al* 2020); and the central Newfoundland gold district (Valentine Lake deposit and Wilding Lake prospect; Honsberger *et al* 2022a).

Robust U–Pb geochronological constraints on gold mineralization are largely restricted to the Neoproterozoic (ca. 570–585 Ma) epithermal prospects of eastern Newfoundland (e.g., O'Brien *et al* 2001; Sparkes *et al* 2005; 2016; Sparkes and Dunning 2014), with fewer constraints on the timing of orogenic gold mineralization in central and western Newfoundland (e.g., ca. 374 Ma muscovite, Sandeman and Dunning 2016; ca. 410 Ma rutile, Honsberger *et al* 2022a). Kerr and Selby (2012) summarized many of the earlier geochronological data for gold mineralization, mostly on the Baie Verte Peninsula, and this contribution builds upon their compilation. We review existing age constraints on gold mineralization in the context of our new $^{40}\text{Ar}/^{39}\text{Ar}$ geochronological data for muscovite from four gold-mineralized

zones across western and central Newfoundland and discuss the data with respect to Paleozoic orogenesis.

CRUSTAL-SCALE ARCHITECTURE OF NEWFOUNDLAND

Williams *et al* (1988) subdivided the Newfoundland Appalachians into four major pre-Silurian tectonic-stratigraphic domains including, from west to east, the Humber, Dunnage, (including the Notre Dame and Exploits subzones), Gander and Avalon zones. The Gander Zone was further divided into Meelpeag, Mount Cormack and Gander Lake subzones. These zones broadly correspond to (also from west to east): the Laurentian margin (Humber); the Dashwoods terrane (extended Laurentian margin and Notre Dame Subzone); Ganderia (peri-Gondwana terranes including all Gander subzones and overthrust intraoceanic Exploits Subzone rocks); and Avalonia (e.g., van Staal *et al* 1996; van Staal and Barr 2012; Fig. 1). The Bras d'Or terrane of Cape Breton Island may extend to the Cinq-Cerf and Grey River areas in southern Newfoundland and its basement may underlie all the Exploits Subzone (Barr *et al* 2014).

The geographic/aerial extent of the Exploits Subzone and its boundaries in northeast-central Newfoundland is misrepresented on most extant compilation maps, particularly its interpreted southeastern termination at the former Gander River Ultrabasic Belt (GRUB Line), now known as the Gander River Complex (cf. O'Neill and Blackwood 1989). The Exploits Subzone includes Ordovician volcanic and sedimentary rocks of the Baie Du Nord Group that are structurally interleaved with Ganderian basement in the Meelpeag Subzone (e.g., Valverde-Vaquero and van Staal 2002). The correlated Ordovician Baie D'Espoir and Davidsville groups in central Newfoundland stratigraphically overlie and are imbricated with Cambrian ophiolites and Gander zone basement (e.g., Blackwood and Green 1983; Colman-Sadd *et al* 1992). Moreover, brachiopod and trilobite-bearing strata of Late to Middle Ordovician (Darriwilian) age, comparable to other Exploits Subzone fauna, are exposed ~30 km east of the Gander River Complex at Indian Bay Big Pond (Fig. 1) in the Gander Lake Subzone (Wonderly and Neumann 1984). The latter are spatially accompanied by pillow basalt, siltstone, conglomerate, and minor gabbro along northeast-trending curvilinear magnetic highs that likely represent klippe of Exploits Subzone assemblages structurally above and interleaved with Ganderian margin sedimentary strata (Miller and Weir 1982; Wonderly and Neumann 1984; O'Neill and Colman-Sadd 1993). Exploits Subzone rocks occur sporadically across parts of the poorly exposed Gander Lake subzone and are likely tectonically intercalated with Gander Zone rocks. Emplacement of these klippe is constrained to Middle Ordovician (Colman-Sadd *et al* 1992; Sandeman and Dickson 2019); thus, the Exploits Subzone and Gander Zone were amalgamated by that time and were no longer independent terranes after the Middle




Figure 1. Simplified geological map of the Island of Newfoundland showing the major geological terranes, tectonic boundaries and faults (modified after Colman-Sadd *et al* 1990) and the locations of the gold mineralized zones discussed in this report. AZ–Apsy zone; BB–Beaverbrook; BR–Browning; BV–Big Vein; CR–Cape Ray; DL–Duder Lake; DVFZ–Doucers Valley Fault zone; GBF–Green Bay Fault; GHF–Gunflap Hills Fault; GL Subzone–Gander Lake Subzone; H–Hammerdown; IBBP–Indian Bay Big Pond; JP–Jonathons Pond; JV–Jaclyn Vein; KP–Kettle Pond; KZ–Keats Zone; LJ–Little Joanna; LP–Leprechaun Pond; LR–Little River; M–Moosehead; MC Subzone–Mount Cormack Subzone; MH–Mosquito Hill; MP Subzone–Meelpaeg Subzone; NAF–Northern Arm Fault; NP–Nugget Pond; NPL–Noel Paul’s Line; PC–Pine Cove; QV–Quinlan Veins; RG–Road Gabbro; RZ–Reid Zone; S–Shrik; SL–The Slip; SR–Salmon River; ST–Stoger’tite; T1–Titan east; T2–Titan west; TG–True Grit; TH–Thor; TW–Twilite; VLSz–Victoria Lake shear zone; WH–Woods Lake and Hill Top; WL–Wilding Lake; YF–Yellow Fox.

Ordovician. Gander River Complex and other linear strongly magnetic anomalies in Gander Zone represent remnant overthrust Ordovician ophiolitic rocks, and Ordovician to Silurian cover rocks of Ganderia that were subsequently structurally modified during younger orogenic events.

The four major pre-Silurian terranes (Laurentian margin, Dashwoods terrane, composite Ganderian terranes, and Avalonia; van Staal and Barr 2014) are separated and transected by several major, long-lived polyphase fault zones. Progressing eastwards, the terrane-bounding fault zones include: the Baie Verte–Brompton Line separating the Laurentian margin from Dashwoods terrane; the Mekwe'jit Line (formerly Red Indian Line; RIL; see White and Waldron 2022) separating Dashwoods terrane from arcs and backarcs of the leading Ganderian margin; the Noel Paul's Line, Dog Bay Line, and Gander River Complex separating allochthonous blocks of composite Ganderia and; the Dover-Hermitage Bay Fault Zone separating Ganderia and Avalonia (e.g., Blackwood and Kennedy 1975; van Staal *et al* 2009; van Staal and Barr 2012; van Staal *et al* 2014, and references therein).

Despite the prevalence of terrane-bounding faults, orogenic gold mineralization in Newfoundland tends to be associated with crustal-scale fault zones that transect the pre-Silurian terranes. In the west, for example, the Silurian to Carboniferous Cabot–Doucers Valley fault system (Lock 1969; Tuach 1987) merges with the Baie Verte–Brompton Line (Fig. 1) to separate the Laurentian margin from Dashwoods terrane (Waldron and van Staal 2001). Furthermore, splays of the Baie Verte–Brompton Line must have been active post-Ordovician accretion to form Silurian vein-hosted gold mineralization (Dubé *et al* 1993; Poulsen *et al* 2000). In central Newfoundland, the approximately five-million-ounce gold deposit at Valentine Lake and numerous gold prospects along strike (e.g., Wilding Lake prospect) are hosted along an Early Devonian thrust-backthrust system (Honsberger *et al* 2022a) that imbricated rocks of the Exploits Subzone and Gander Zone and uplifted the Meelpaeg nappe along the Noel Paul's Line.

The new $^{40}\text{Ar}/^{39}\text{Ar}$ geochronology samples in the present study are from: (1) the White Bay area along the Doucer's Valley–Cabot Fault system of the Laurentian margin (Jacksons Arm, Shrik prospect); (2) the southwestern confluence of the Exploits and Meelpaeg subzones with Dashwoods terrane (Wood Lake South zone and Hill Top showing; Leprechaun Pond deposit, Valentine Lake); and (3) the northeastern Exploits Subzone of central Newfoundland (Moosehead prospect; Fig. 1). Existing geochronological constraints for additional auriferous zones are also illustrated in Figure 1.

AURIFEROUS ZONES INVESTIGATED WITH $^{40}\text{AR}/^{39}\text{AR}$ GEOCHRONOLOGY

Jacksons Arm trend (Shrik prospect)

The Jacksons Arm gold trend (Shrik, Boot n Hammer, and Stocker zones) occur ~ 4 km north of the community

of Jacksons Arm on the Coney Head Peninsula in western White Bay (Fig. 1; Reid and Myllyaho 2012; Myllyaho 2013; English *et al* 2017). The showings occur in a curvilinear zone along the margin of the Ordovician Coney Head Complex (ca. 478 Ma; Dunning 1987; Fig. 2), an ophiolite-related tonalite and marginal basaltic and sedimentary units. The Coney Head Complex is unconformably overlain by syntectonic, clastic sedimentary and bimodal volcanic rocks of the orogenic gold-mineralized Silurian Sops Arm Group (e.g., Heyl 1937; Betz 1948; Kerr 2006a and b; Sandeman and Dunning 2016). The unconformity and the volcanic-sedimentary rocks of the Sops Arm Group are thrust imbricated with tonalite of the Coney Head Complex along broadly north-striking, east-dipping fault zones (this study; Magna Terra Minerals 2022). A sample of altered tonalite and adhering quartz vein (HS12-200C), weakly anomalous in Bi (3.9 ppm), As (24 ppm), Ag (0.1 ppm), and Au (4650 ppb), was collected from the Shrik trench where fine- to medium-grained, variably foliated, sericite-altered, pyritic and quartz-veined tonalite of the Coney Head Complex is exposed over a ~10 m × ~30 m area (Figs. 3a and b). The tonalite is locally cut by irregular, pinch and swell quartz veins with rare pyrite, but the relationships between deformation, veining, and gold mineralization remain unresolved. The tonalite wall rock contains quartz, sericitized plagioclase, mats of intergrown fine-grained muscovite and sparse pyrite and goethite (Fig. 3c).

Wood Lake South (Main) zone - Hill Top showing

The Wood Lake South (Main) gold zone and Hill Top showing occur in central-western Newfoundland near the confluence of the Exploits, Meelpaeg, and Notre Dame subzones (Figs 1 and 4). The Hill Top gold showing consists of two narrow (1–10 cm wide), southeast-northwest-trending, steeply dipping, pinch and swell pyrite + arsenopyrite-bearing quartz veins that cut Ordovician (ca. 467 Ma) Peter Strides monzogranite (e.g., van Egmond 2004; van Staal *et al* 2005a; Valverde-Vaquero *et al* 2006; Sandeman *et al* 2014; Sandeman 2014a) of the Meelpaeg nappe and extend sporadically along strike for ~50 m. The quartz veins are hosted in an inferred northwest-trending mineralized fracture zone that cuts the monzogranite (Figs. 4 and 5). The host granite is strongly foliated, with a strong quartz mineral lineation, and is cut by translucent-white, foliated and lineated, barren quartz veins oriented sub-parallel to the foliation. The $^{40}\text{Ar}/^{39}\text{Ar}$ geochronology sample from Hill Top (HS10-59A) is a light grey to pink, fine- to medium-grained, quartz-veined, foliated and lineated monzogranite adjacent to the two quartz veins (Figs. 5a and 5b). In thin section, polyhedral and sutured quartz porphyroclasts (≤ 1.5 mm) are surrounded by fine-grained intergrown anhedral albite, quartz, and sericite (Fig. 5c). Rare euhedral monazite and subhedral rutile are intergrown with quartz and sericite in the groundmass (Fig. 5d).

Gold-mineralized quartz veins of the Wood Lake South (Main) zone are hosted by massive to brecciated and/or(?)




Figure 2. Simplified geology of part of the western White Bay area, northwestern Newfoundland (modified after Kerr 2006b; Sandeman and Dunning 2016), showing the approximate location of $^{40}\text{Ar}/^{39}\text{Ar}$ muscovite sample HS12-200C (yellow star), as well as the locations of precious- and base-metal showings of the region (Geoscience Atlas 2022)

mylonitic, orange-pink monzogranitic rocks of the Peter Strides Granite Suite. In the main trench (Fig. 6a), intensely deformed monzogranite is intruded by a less deformed, fine- to medium-grained, sericitic monzogranite. The less intensely deformed monzogranite is cut by an extensive array of steeply dipping irregular fractures and anastomosing, pinch and swell quartz veins (≤ 10 cm). Both the veins and fractures have minor pyrite + hematite \pm arsenopyrite, accompanied by adjacent wall rock sericitization, albitization and silicification. Muscovite locally forms randomly

oriented clumps and masses in the matrix of brecciated monzogranite, but occurs more typically as wispy platelets along fractures (Figs. 6b and 6c). Further details of the geology of the main trench and associated auriferous zones are provided by van Egmond and Cox (2005) and Sandeman *et al.* (2014).

The $^{40}\text{Ar}/^{39}\text{Ar}$ geochronology sample (HS13-063A) from the Wood Lake South (Main) zone trench consists of an orange-pink, medium-grained, brecciated, and foliated monzogranite exposed below a strand of mylonitic monzogran-




Figure 3. Photographs and photomicrographs of Shrik Showing sample HS12-200C. (a) Photograph of the Shrik discovery trench looking north with quartz-veined, altered tonalite in the centre, bounded to the north and south by sericite altered tonalite. The field of view is 10 m. (b) Close-up of the Shrik trench with quartz veined tonalite mantled by altered tonalite, and the location of sample HS12-200C. Field of view is 15 cm. (c) Photomicrograph in plane polarized light (PPL) showing the mineralogy of the altered tonalite and the random, mat-like nature of the muscovite in the sample. The brown tint of the muscovite is because of goethite staining. Mineral abbreviations after Whitney and Evans (2010).

ite (Figs. 4 and 6). In thin section, large (≤ 2 mm) polyhedral and sutured quartz grains are surrounded by fine-grained albite, sericite, and quartz. Subhedral pyrite and arsenopyrite are dispersed throughout and locally concentrated in vugs and fractures (Fig. 6b). Relatively large (≤ 1 mm), weakly to non-aligned muscovite grains are common (Figs. 6c and d).

Leprechaun Pond deposit (Valentine Lake)

The Leprechaun Pond deposit of the Valentine Lake gold property in central-western Newfoundland occurs within the Exploits Subzone near the boundary with the Meelpaeg Subzone (Figs. 1 and 7). The Neoproterozoic to Silurian rocks of the Valentine Lake area trend northeasterly and are bisected by the Victoria Lake shear zone (Valverde-Vaquero *et al* 2006; van Staal *et al* 2005b). This major northeast-trending, variably southeast-dipping, crustal-scale shear zone extends from the Gunflap Hills Fault in the southwest, through central Newfoundland along the northern margin of the Meelpaeg nappe (Figs. 1 and 7). The northeastern trace of the Victoria Lake shear zone is poorly constrained, and it may continue north-eastward to the Bay of Exploits, or alternatively, it may bifurcate and verge to the east along the northern margin of the Mount Cormack Subzone (Honsberger *et al* 2022a).

To the northwest, in the structural footwall of the Victoria Lake shear zone, Neoproterozoic (ca. 570 Ma) orogenic gold-mineralized basement granitoid rocks of the Valentine Lake Intrusive Suite (Evans and Kean 2002; Rogers and van Staal 2002; Rogers *et al* 2006), are uplifted and juxtaposed against the orogenic gold-mineralized Rogerson Lake Conglomerate (Kean and Jayasinge 1980) along the northwest-dipping Valentine Lake shear zone. The Valentine Lake shear zone hosts an approximately five-million-ounce orogenic gold resource, including the Leprechaun Pond deposit, where bleached and altered Neoproterozoic trondhjemite-tonalite and associated rocks contain a stacked array of fault-fill and extensional quartz-tourmaline-pyrite (QTP) vein sets (Lincoln *et al* 2018). The Rogerson Lake Conglomerate crops out southeast of the Valentine Lake shear zone, and is an aerially extensive latest Silurian, syntectonic, clastic sedimentary unit of central Newfoundland that delineates the southwest-northeast-trending fault system that is host to many of the gold occurrences in the central Newfoundland gold district (Honsberger *et al* 2022b; this volume).

At the Wilding Lake gold property, ~36 km northeast along strike of the Leprechaun Pond deposit, orogenic gold mineralization is hosted within both the Rogerson Lake Conglomerate and the associated ca. 422 Ma felsic subvolcanic and volcanic rocks in the footwall of the northeastern extension of the Valentine Lake shear zone (WL on Fig. 1; Honsberger *et al* 2019a; 2019b; 2020a; 2020b; 2022a). Overlapping ca. 410 Ma ID-TIMS U–Pb ages for hydrothermal rutile in quartz veins from both the Leprechaun Pond deposit and the Elm Prospect of the Wilding Lake property (Honsberger *et al* 2022a) indicate an Early Devonian (late




Figure 5. Photographs and photomicrographs of Hill Top Showing sample HS10-059A. (a) Photograph viewed southeast at the narrow Hill Top veins. Note person in background. (b) Photograph of a foliation surface on the rock adjacent to the Hill Top vein, illustrating the strong L-fabric in the Peter Strides monzogranite defined by aligned quartz and muscovite. (c) Photomicrograph in crossed polars of the monzogranite wall rock illustrating the disseminated subhedral pyrite, the altered sericitic groundmass and the large, secondary muscovite grains. (d) Backscattered electron image illustrating the minerals and their detailed relationships. Note the subhedral rutile and euhedral monazite, targets for future geochronology.

(Fig. 8c); a 250–450 μm fraction was extracted for $^{40}\text{Ar}/^{39}\text{Ar}$ geochronology.

Moosehead prospect (North Pond trench; TRMH15-3)

The Moosehead Prospect occurs southeast of the town of Bishops Falls in north-central Newfoundland (Figs. 1 and 9; Morgan 2016; Froude 2019, 2021). It lies within the latest Silurian–Early Devonian, intraorogenic Botwood Basin, ~4 km west of the ca. 425–418 Ma Mount Peyton Intrusive Suite (Sandeman *et al* 2017) in the north-central Exploits Subzone (Figs. 1 and 9). Mineralization consists of extensional, auriferous, sulphide-sulfosalt-bearing quartz veins and breccias that cut mainly muscovite-bearing sandstone and siltstone of the Wigwam Formation of the Botwood Group (i.e., Botwood basin; Williams 1969; Dickson *et al* 2000; O'Brien 2003; Morgan 2016). Mafic volcanic rocks of the lower Botwood Group (Laurenceton Formation) occur immediately south and southeast of the prospect. Gabbroic and fine-grained mafic dykes locally cut the sedimentary rocks and are themselves locally bleached, silicified and

carbonate + pyrite \pm sericite altered (e.g., Clark 1999; Morgan 2016). Trenching uncovered medium-bedded muscovite-bearing sandstone, southwest-dipping in the north, and south-dipping in the south, with weakly sulphidic, highly disrupted, quartz-veined, sandstone-quartz breccia in between (Figs. 9a, b and 10). The sandstone is variably mineralized, and quartz veins are more abundant in the northern part of the trench. Mineralization occurs in the hanging wall of a discrete east-west fault zone (Figs. 9b and 10). Bedding and crosscutting quartz veins are deflected into the fault zone, consistent with dextral rotation and reverse(?) slip. Further description and assay information for the Moosehead prospect are presented in Morgan (2016).

The $^{40}\text{Ar}/^{39}\text{Ar}$ geochronology sample from the Moosehead trench is a beige-grey, medium-grained, medium-bedded, quartz-veined muscovite-bearing sandstone of the Wigwam Formation from immediately north of the east-west fault near the centre of the trench (Figs. 9b and 10a). In thin section, large (≤ 0.4 mm) quartz and muscovite grains are surrounded by fine-grained matrix of albite, sericite, and quartz with subhedral pyrite dispersed throughout (Fig. 10b).




Figure 6. Photographs and photomicrographs of Wood Lake South Main zone sample HS13-063A. (a) Photograph from the main trench with massive and brecciated, quartz-veined and sericite + pyrite \pm arsenopyrite mineralized monzogranite cutting mylonitic monzogranite having fabric-parallel, sulphide-poor, internally deformed quartz veins. The location of sample HS13-063A is shown and the Geotul is 54 cm in length. Mylonitic fabric trends 136° and dips 66° to the southwest. (b) Cut slab photograph of sample HS13-063A. White is altered feldspar, grey is quartz and reflective silver grains are muscovite. (c) Photomicrograph in PPL of the brecciated altered monzogranite illustrating the disseminated subhedral pyrite, the altered sericite groundmass and the large, secondary muscovite grains.

Relatively large (≤ 1 mm), bedding-parallel muscovite grains are common, and appear to be detrital in origin (Fig. 10c).

$^{40}\text{Ar}/^{39}\text{Ar}$ GEOCHRONOLOGY

Methods

Geochronology samples were collected during 2010–2015 from selected gold-mineralized zones and drill core across central and western Newfoundland (e.g., Sandeman *et al* 2013; Sandeman 2014b; Sandeman and Dunning 2016). Kilogram-sized hand samples or ~ 40 cm long pieces of split drill core were cut and cleaned of all weathered surfaces and crushed to ~ 1 cm chips. About 100 grams of each sample was gently hand pulverised with pestle and mortar and sieved to two size fractions, 250–420 μm and 180–250 μm . The grain separates were washed repetitively in deionized water and the solute drained. Cleaned and dried samples were then gently cleaned in dilute ($\sim 0.5\text{N}$) HNO_3 to remove adhering sulphides and carbonates from the grain separates. The cleaned separates were re-sieved and muscovite was hand-picked under a binocular microscope for laser step-heating analyses. Muscovite from the Shrik Showing (HS12-200C) was of poor quality, fine-grained, and only a single muscovite-rich (180–250 μm) grain separate was extracted.

The $^{40}\text{Ar}/^{39}\text{Ar}$ age data were obtained in the $^{40}\text{Ar}/^{39}\text{Ar}$ ThermoChronology Laboratory at Queen's University. Mineral separates and flux-monitors (standards) are wrapped in Al-foil, stacked sequentially into an 8.5-cm-long and 2.0-cm-diameter Al irradiation capsule, and then irradiated with fast neutrons in position 8D of the McMaster Nuclear Reactor (Hamilton, Ontario) for a duration of 72 h (at 3 MWH). Packets of flux monitors are located at ~ 0.5 cm intervals along the irradiation container and the J-value for an individual sample is determined by least-squares, second-order polynomial interpolation using replicate analyses of splits for each monitor position in the capsule. The samples are loaded into flat-bottomed pits in a copper sample-holder and placed beneath the ZnS view-port of a small, bakeable, stainless-steel chamber connected to an ultra-high vacuum purification system. Following bake out at 100°C , a 30 watt New Wave Research MIR 10-30 CO_2 laser with a faceted lens is used to heat samples for ~ 3 minutes at increasing percent power settings (2 to 45%; beam diameter 3 mm). After purification using hot and cold SAESC50 getters (for ~ 5 minutes), the evolved gas is admitted to an MAP 216 mass spectrometer, with a Baur Signer source and an analog electron multiplier (set to a gain of 100 over the Faraday detector). Measured argon-isotope peak heights are extrapolated to zero-time and corrected for discrimination using a $^{40}\text{Ar}/^{36}\text{Ar}$ atmospheric ratio of 295.5 and measured ratios of atmospheric argon. Blanks, measured routinely, are subtracted from the subsequent sample gas-fractions. The extraction blanks are typically $<10 \times 10^{-13}$, $<0.5 \times 10^{-13}$, $<0.5 \times 10^{-13}$, and $<0.5 \times 10^{-13}$ cm $^{-3}$ STP for masses 40,




Figure 7. Simplified geology of the area around the Leprechaun Deposit (modified after van Staal *et al* 2005b) showing the approximate locations of the $^{40}\text{Ar}/^{39}\text{Ar}$ muscovite sample and the recently dated CA-TIMS U–Pb titanite sample (VL335-245; Honsberger *et al* 2022) at the Leprechaun Pond deposit. The locations of precious- and base-metal showings are from the Geoscience Atlas (2022).

39, 37, and 36, respectively. The ^{39}Ar and ^{37}Ar are corrected for radioactive decay during and after irradiation. Corrections are made for neutron-induced ^{40}Ar from potassium, ^{39}Ar and ^{36}Ar from calcium, and ^{36}Ar from chlorine (Roddick 1983; Onstott *et al* 1991). Dates and errors are calculated using the procedure of Dalrymple *et al* (1981) and the constants of Steiger and Jäger (1977). Plateau and inverse isotope correlation dates are calculated using ISOPLOT v. 3.60 (Ludwig 2012). A plateau is herein defined as 3 or more contiguous steps containing $>50\%$ of the ^{39}Ar released, with a probability of fit >0.01 and MSWD <2 . Errors shown in Supplementary data Table S1 and in Figure 11 represent the analytical precision at 2σ , assuming that the errors in the ages of the flux monitors are zero. This is suitable for comparing within-spectrum variation and determining which steps form a plateau (e.g., McDougall and Harrison 1988, page 89). The dates and J-values are referenced to GA-1550biotite (98.5 Ma; Spell and McDougall 2003) and Hb-3Grhornblende (PP-20; 1073.6 Ma; Jourdan *et al* 2006). The complete $^{40}\text{Ar}/^{39}\text{Ar}$ dataset is in Supplementary data Table

S1, whereas the laser step-heating results are illustrated in Figure 11. Sample coordinates are NAD27 datum, zone 21.

Results

Shrik Showing (HS12-200C: UTM Zone 21, 516037 E, 5527889 N: muscovite)

Altered tonalite wall rock from this sample was crushed, sieved, cleaned, and small (≤ 0.5 mm), weakly to non-aligned, goethite-dusted muscovite grains were concentrated into a 180–250 μm grain separate for $^{40}\text{Ar}/^{39}\text{Ar}$ dating. The muscovite separate yielded a complex age spectrum (Fig. 11a). The initial low-T (power) step (1.43% of ^{39}Ar released) yielded an anomalously young age (ca. 336 Ma) but the next 6 steps (30.3% of ^{39}Ar released) yielded anomalously old ages ranging from ca. 390 to 428 Ma and are attributed to the incorporation of excess argon. The higher temperature (power) steps, excluding the final young gas fraction, form a relatively flat segment representing 58.9 % of the $^{39}\text{Ar}\%$




Figure 8. Photographs and photomicrograph of Leprechaun Pond deposit. (a) Photograph from the main trench to the northwest of Leprechaun Pond illustrating the sericite-altered trondhjemite-tonalite of the Valentine Lake Intrusive Suite cut by quartz-tourmaline-pyrite veins. (b) Cut core photograph of quartz-tourmaline vein cutting bleached trondhjemite (DDH VL-24-62.1m). (c) Photomicrograph in PPL of a quartz-tourmaline vein containing pyrite, calcite and muscovite cutting altered trondhjemite host rock composed of saussuritized plagioclase and quartz. Muscovite forms coarse plates along fractures in plagioclase and in vein margins.

released and yield an age of 379.7 ± 1.4 Ma (2σ ; POF = 0.36; MSWD = 1.10). The volume of ^{39}Ar released is slightly less than the suggested convention of >60% volume for a plateau age (McDougall and Harrison 1988; Ludwig 2012; Schaen *et al* 2021), but the quasi-plateau appears to have geological significance.

**Hill Top Showing (HS10-59A: UTM Zone 21,
443526 E, 5334700 N: muscovite)**

Analyses of relatively large (250–425 μm) muscovite grains yielded a relatively simple $^{40}\text{Ar}/^{39}\text{Ar}$ age spectrum (Fig. 11b). The nine higher temperature (power) steps formed a relatively flat segment representing 91.4% of the ^{39}Ar released and gave a plateau age of 409.1 ± 1.3 Ma (2σ ; POF = 0.77; MSWD = 0.42).

**Wood Lake South (Main) zone (HS13-063A: UTM
Zone 21, 500450 E, 5504824 N: muscovite)**

Two fractions (250–425 μm and 180–250 μm) of muscovite were extracted for $^{40}\text{Ar}/^{39}\text{Ar}$ analysis. The nine highest (out of 15) temperature (power) steps of the 250–425 μm fraction, representing 91.4 % of the ^{39}Ar released gives a plateau age of 403.1 ± 1.2 Ma (2σ ; POF = 0.91; MSWD = 0.42; Fig. 11c). The second 180–250 μm aliquot (Fig. 11d) yielded a quasi-plateau age of 398.4 ± 1.3 Ma, representing 56.9% of the ^{39}Ar released (2σ ; MSWD = 0.74; POF = 0.65), due largely to an unfortunately large-volume last step. The plateau age of 403.1 ± 1.2 Ma is considered the best estimate of the age of the muscovite.

**Leprechaun Pond Deposit (Valentine Lake: VL-24_62.1m:
UTM Zone 21, 486400 E, 5355880 N: muscovite)**

The muscovite at the Leprechaun Pond deposit yielded a somewhat jagged age spectrum (Fig. 11e), but five of the steps formed a flat segment, comprising 59.9% of the ^{39}Ar released, and yielded a quasi-plateau age of 384.2 ± 1.6 Ma (2σ ; POF = 0.57; MSWD = 0.74) and is interpreted as the best estimate of the age of the muscovite.

**Moosehead Prospect (HS15-137; 613561E,
5428182N: muscovite)**

Two muscovite fractions (250–425 μm and 180–250 μm) were extracted for $^{40}\text{Ar}/^{39}\text{Ar}$ geochronology and yielded two similar age spectra (Figs. 11f, g). The coarse-grained (250–425 μm) aliquot yielded a well-defined $^{40}\text{Ar}/^{39}\text{Ar}$ plateau age of 457.3 ± 1.2 Ma (2σ ; POF = 0.94; MSWD = 0.44), representing 91.4% of the ^{39}Ar released (Fig. 11f). The 180–250 μm grain-size fraction (Fig. 11g), similarly yielded a well-defined $^{40}\text{Ar}/^{39}\text{Ar}$ plateau age of 453.5 ± 1.1 Ma (2σ ; POF = 0.48; MSWD = 0.97), representing 97.0% of the ^{39}Ar released. The older age of 457.3 ± 1.2 Ma for the coarser-grained 250–450 μm fraction is interpreted as the best estimate of the $^{40}\text{Ar}/^{39}\text{Ar}$ age of the muscovite.




Figure 9. (a) Simplified geology of the northeastern Botwood Basin and the location of the Moosehead Prospect and other local mineral occurrences (modified after the Geoscience Atlas (2022)). (b) Geological map of trench TRMH15-3 at the Moosehead Prospect with the locations of the $^{40}\text{Ar}/^{39}\text{Ar}$ muscovite sample, other lithogeochemical samples and the location of DDH MH-02-15 (see Morgan 2016).

DISCUSSION

Integration of these new $^{40}\text{Ar}/^{39}\text{Ar}$ geochronological data with existing regional geochronological constraints on gold mineralized zones and magmatic events elucidates deformational, metamorphic, and tectonic intervals that were favourable for gold-mineralization across Newfoundland (Fig. 12). The collective compiled data are diverse as they include age constraints determined via different radiometric methods on various phases/mineral species: some of which

are more robust than others. These include studies applying $^{40}\text{Ar}/^{39}\text{Ar}$ cooling ages on dynamo-metamorphism and mineralization-associated potassic alteration; Re–Os ages on sulphide minerals; implied maximum ages based on U–Pb ages of host rocks and direct U–Pb CA-TIMS dating of the minerals coeval with quartz vein emplacement. For example, the interpreted age of the Hammerdown gold deposit (Springdale Peninsula) comes from U–Pb (zircon) thermal ionization mass spectrometry (TIMS), which yielded an age of 437 ± 4 Ma for a felsic dyke cut by an auriferous vein




Figure 10. Photographs and photomicrographs of the Moosehead Prospect trench. (a) Photograph looking north from the east-west fault with decimetre-scale bedded muscovitic sandstone cut by an array of narrow, weakly sulphidic quartz veins. These veins largely trend northwest-southeast and are rotated into an east-west orientation near the main fault, indicating this is a dextral fault. (b) Close up of the warped bedding and crosscutting quartz veins at sample location HS15-137. Note the folding of bedding and the small yellow arrows pointing to quartz veins. (c) Photomicrograph in PPL of the sandstone showing disseminated subhedral pyrite, carbonate-sericite-altered groundmass and large, bedding-parallel muscovite grains.

(Ritcey *et al* 1995). This is therefore a maximum age constraint for gold mineralization. In contrast, the ca. 410 Ma chemical abrasion (CA) TIMS ages for rutile from auriferous quartz veins at the Wilding Lake gold prospect and Leprechaun Pond gold deposit (Honsberger *et al* 2022a) provide high-precision maximum constraints on the age of orogenic gold-mineralized quartz vein emplacement. Detailed investigations of vein and alteration mineral parageneses, their relationships to gold deposition and systematic geochronological study of different mineral species are required to better understand the timing of gold mineralization in Newfoundland.

The collective geochronological data outline two distinct, broad age-ranges for orogenic-style gold mineralization in the central and western Newfoundland Appalachians (Fig. 12). These include an older, Wenlock (middle Silurian) to Lochkovian (Lower Devonian) ca. 440 to ca. 405 Ma event, and a younger, Middle Devonian event at ca. 400–375 Ma. The former includes muscovite growth/resetting intervals for gold-bearing zones of the Laurentian margin, Dashwoods terrane, western Exploits subzone and along the Meelapaeg–Exploits subzones boundary; the latter includes muscovite from gold-bearing zones across the Laurentian Margin, Dashwoods terrane, Exploits Subzone, and Gander Zone (Fig. 12).

The muscovite from the Shrik showing occurs in an Ordovician, intraoceanic arc-related tonalite that has a strongly metaluminous composition (Sandeman, unpublished data 2022). The muscovite is spatially associated with quartz veining and bleaching of the tonalite resulting from hydrothermal fluid flow; therefore, it formed as a result of metasomatism, opposed to by igneous or dynamothermal metamorphic processes. Accordingly, the $^{40}\text{Ar}/^{39}\text{Ar}$ geochronological result of 379.7 ± 1.4 Ma is interpreted as a maximum age constraint for gold mineralization.

The host rocks of the Hill Top and Woods Lake South zones are muscovite-biotite-bearing, strongly peraluminous granitoid rocks of the Ordovician Peter Strides suite (part of the Meelapaeg Subzone) that comprise the hanging wall of the Victoria Lake-Valentine Lake fault system. The northern portion of the Meelapaeg Subzone records metamorphic U–Pb monazite and titanite ages and $^{40}\text{Ar}/^{39}\text{Ar}$ hornblende, muscovite and biotite cooling ages that range from ca. 418 to 400 Ma (Valverde-Vaquero *et al* 2003). Muscovite from the Hill Top (409.1 ± 1.3 Ma) and Woods Lake zones (403.1 ± 1.2 Ma) therefore likely represent Acadian metamorphic cooling temperatures in the immediate hanging wall of the Victoria Lake Shear Zone, which represent best approximations of maximum ages of gold mineralization.

Metaluminous tonalite and trondhjemite of the Valentine Lake Intrusive Suite host the Leprechaun Pond gold deposit. These rocks do not contain primary muscovite and the white mica is hydrothermal in origin. However, the relationship between the muscovite and deformation of the auriferous quartz-tourmaline-pyrite veins is ambiguous. The hydrothermal rutile age of ca. 410 Ma (see Honsberger *et al* 2022a) is significantly older than the muscovite plateau

age of 384.2 ± 1.6 Ma (this study), suggesting either a single protracted alteration and mineralization event, or multiple discrete events.

In contrast to the other dated samples, muscovite from the sandstone of the Moosehead prospect (ca. 457.3 ± 1.2 Ma and 453.5 ± 1.1 Ma) is detrital in origin and oriented parallel to bedding. Therefore, these ages represent metamorphic cooling ages for the metasedimentary or granitoid sources for the sandstone. One proximal potential Ordovician (Darriwillian; Cohen *et al* 2013) source terrane for the detrital muscovite is the Mount Cormack Complex (Gander Zone) of central Newfoundland (Fig. 1; Williams *et al* 1988; Colman-Sadd *et al* 1992). The Mount Cormack Complex preserves a central core of foliated, upper amphibolite-grade migmatitic psammite and pelite that were metamorphosed and intruded by garnet-muscovite syenogranite (Through Hills granite) at ca. 465–464 Ma (Colman-Sadd *et al* 1992; Valverde-Vaquero *et al* 2006). Metamorphic ages from the complex include: a monazite age of 462 ± 1 Ma; a titanite age of 460 ± 3 Ma; a $^{40}\text{Ar}/^{39}\text{Ar}$ hornblende plateau age of 465.7 ± 6.4 Ma; and a $^{40}\text{Ar}/^{39}\text{Ar}$ biotite plateau age of 439.4 ± 2.4 Ma (Valverde-Vaquero *et al* 2003, 2006). The hornblende and biotite ages bracket the interval defined by the Moosehead trench muscovite ages and support the suggestion of the Mount Cormack Complex as a possible detrital muscovite source. Muscovite from the Meelpaeg nappe yield younger, Devonian regional metamorphic cooling ages ranging from ca. 418 to 394 Ma (Valverde-Vaquero *et al* 2003), similar to those obtained in the Gander Lake Subzone (ca. 404 to 388 Ma; O'Neill and Lux 1989; O'Neill and Colman-Sadd 1993). The apparent absence of older, Ordovician muscovite in these areas does not entirely preclude them as possible sources, however, this appears to be the case. To the west of the Botwood Basin, an alternative Ordovician, muscovite-bearing source might be represented by the muscovite-biotite-psammite of the reworked Laurentian margin (e.g., Fleur de Lys Supergroup, Baie Verte Peninsula).

The Early Devonian muscovite ages from central Newfoundland (Hill Top showing and Wood Lake South zone) are similar to, but slightly younger than, the ca. 410 Ma ages determined along strike to the northeast for hydrothermal rutile in gold-mineralized veins from the Leprechaun Pond deposit (Valentine Lake) and Wilding Lake prospect (Honsberger *et al* 2022a). This may reflect a lower $^{40}\text{Ar}/^{39}\text{Ar}$ closure temperature for muscovite relative to the U–Pb closure temperature for rutile (ca. 600°C; Vry and Baker 2006). The ca. 410 Ma rutile ages are interpreted to provide the best minimum estimates for quartz vein emplacement and initial orogenic gold mineralization along the Victoria Lake–Valentine Lake fault corridor (Fig. 1), whereas the muscovite


ages provide the best estimates for potassic alteration associated with quartz vein emplacement. Furthermore, the ca. 403 and 409 Ma muscovite ages are also in broad agreement with two identical 406 ± 2 Ma $^{40}\text{Ar}/^{39}\text{Ar}$ plateau ages for alteration-related muscovite from the Mosquito Hill gold prospect along the southeastern margin of the Mount Cormack Complex (Fig. 1; Sandeman *et al* 2013).


Early Devonian metamorphism, metasomatism and orogenic gold mineralization along the Victoria Lake–Valentine Lake fault corridor is compatible with the timing of initial Acadian thrusting after ca. 418 Ma in southcentral Newfoundland (Dunning *et al* 1990; Valverde-Vaquero and van Staal 2001; van der Velden *et al* 2004; Valverde-Vaquero *et al* 2006). Furthermore, Early Devonian mineralization coincides with ca. 415 to 410 Ma Acadian ductile deformation in north-central and southwestern Newfoundland (Dunning *et al* 1990; Dubé *et al* 1996; McNicoll *et al* 2006). In southwestern–central Newfoundland, $^{40}\text{Ar}/^{39}\text{Ar}$ geochronology of white mica provides evidence for low-*P* – low-*T* Acadian metamorphism between ca. 408 and 390 Ma (Willner *et al* 2018). On the Baie Verte Peninsula, orogenic gold mineralization may have been initiated earlier than in central Newfoundland as Re–Os pyrite geochronology yielded ages of 420 ± 7 Ma and 411 ± 7 Ma for the Stog'er Tight and Pine Cove deposits (Kerr and Selby 2012), respectively. Similarly, hydrothermal zircon from the Stog'er Tight deposit was dated at 420 ± 5 Ma (Fig. 1; Ramezani *et al* 2002).

The new Middle–Upper Devonian muscovite ages for the Leprechaun deposit (ca. 384 Ma) at Valentine Lake and Shrik showing at White Bay (ca. 380 Ma), contrast with Early Devonian orogenic gold-bearing vein formation documented in both locations (ca. 410 Ma rutile; Honsberger *et al* 2022a; ca. 419 to ca. 408 Ma; Kerr and van Breemen 2007; Minnett *et al* 2012). However, the new data are compatible with a previous age determination for secondary muscovite at White Bay (ca. 374 Ma, Sandeman and Dunning 2016), as well as with ca. 374 ± 8 Ma hydrothermal xenotime from a gold-mineralized vein at the Nugget Pond deposit on the Baie Verte Peninsula (Sangster *et al* 2008). Moreover, a gold-mineralized mafic sill/dyke that cuts the Late Silurian Indian Islands Group near Gander Bay at the Titan prospect (Fig. 1; McNicoll *et al* 2006) yielded a SHRIMP U–Pb zircon age of 381 ± 5 Ma, providing a maximum, late Devonian age for the gold mineralization. All these data support the existence of a Middle–Upper Devonian gold mineralization event that is superimposed on Early Devonian quartz vein systems in western, west-central, and central Newfoundland.

All geochronologically constrained orogenic gold occurrences in central and western Newfoundland are Late

Figure 11. (next page) $^{40}\text{Ar}/^{39}\text{Ar}$ age spectra for muscovite grain separates from the samples under investigation. (a) Shrik Showing muscovite HS12-200C (250–425 μm). (b) Hill Top Showing muscovite HS10-059A (250–425 μm). (c) Wood Lake South (Main) zone muscovite HS13-063A-1 (250–425 μm). (d) Wood Lake South (Main) zone muscovite HS13-063A-2 (180–250 μm). (e) Leprechaun Pond Deposit muscovite VL-24-62.1m (250–425 μm). (f) Moosehead Prospect detrital muscovite HS15-137-1 (250–425 μm). (g) Moosehead Prospect detrital muscovite HS15-137-2 (180–250 μm).





SANDEMAN ET AL. – Overview of age constraints for gold mineralization in central and western Newfoundland and new ⁴⁰Ar/³⁹Ar ages for muscovite from selected auriferous zones

Figure 12. Diagram summarizing geochronological constraints on gold mineralized zones of central and western Newfoundland. Timescale from the International Commission on Stratigraphy stratigraphic chart (Cohen *et al* 2013; updated 2021). Also outlined are the approximate estimates for the orogenic events of the Newfoundland Appalachians (e.g., van Staal and Barr 2012). The orogenic-style mineralization occurs in two broad intervals, one in the latest Silurian to Early Devonian and a second in the Middle to Late Devonian. The latter may be synchronous with, and immediately postdate accretion of the Meguma terrane. Data sources: 1- Minnett *et al* (2012); 2- Kerr and van Breemen (2007); 3- Dubé *et al* (1996); 4- Sandeman and Dunning (2016); 5- Kerr and Selby (2012); 6- Ramezani *et al* (2002); 7- Sangster *et al* (2008); 8- Ritcey *et al* (1995); 9- Sandeman (2014d); 10- Honsberger *et al* (2022); 11- Sandeman *et al* (2013); 12- Colman-Sadd *et al* (1992); 13- Valverde-Vaquero *et al* (2003); 14- Sandeman and Dickson (2019); 15- Sandeman *et al* (2017); 16- Sandeman and Spurrell (2020); 17- McNicoll *et al* (2006).

Silurian to Middle–Upper Devonian and define two gold-forming intervals broadly between ca. 433 and 405 Ma and ca. 390 and 372 Ma. Spatiotemporal coincidence of orogenesis, metamorphism, magmatism and precious metal mineralization occurs across much of the Laurentian margin, Dashwoods terrane, the western Exploits Subzone and selected parts of the eastern Exploits Subzone. The oldest orogenic gold deposits occur in the west along the Laurentian margin and eastern Dashwoods terrane and are associated with the terminal stages of the Middle to Late Silurian Salinic orogenic cycle (ca. 435–427 Ma: e.g., van Staal and Barr 2012). Moreover, regional magmatic, sedimentary, metamorphic, and structural events indicate that although terrane accretion progressed oceanward (present-day east) throughout the Paleozoic (e.g., van Staal *et al* 2014), earlier accreted terranes were affected by mineralizing events related to subsequent accretion of outboard terranes (e.g., Honsberger *et al* 2022a). Thus, mineralized zones in central and western Newfoundland yield Middle–Late Silurian to Middle Devonian ages that post-date accretion of the encompassing terrane. For example, accretion of Dashwoods to the Laurentian margin took place in the Ordovician (Waldron and van Staal 2001), but gold mineralization at White Bay (Humber Zone, Fig. 1) is Late Silurian to Early or Middle Devonian (Fig. 12). Furthermore, ca. 410 Ma orogenic gold mineralization along the Victoria Lake–Valentine Lake fault corridor post-dates ca. 435 Ma accretion of the composite Gander Zone/Exploits Subzone (Ganderia) to the Dashwoods terrane and composite Laurentia (e.g., Honsberger *et al* 2022a). These relationships indicate that other tectonic processes in addition to accretion, such as oroclinal bends and orogen-parallel along strike temporal variations in the kinematics of orogenic accommodation, were critical to generating the orogenic gold-mineralized faults in central and western Newfoundland, as is presumed elsewhere along the Iapetan suture zone of the Appalachian mountain belt (Romer and Kroner 2018).

CONCLUSIONS

1. New $^{40}\text{Ar}/^{39}\text{Ar}$ geochronological constraints across central and western Newfoundland are consistent with two distinct intervals of orogenic gold mineralization between ca. 433 and 405 Ma and ca. 390 and 372 Ma.

2. The late Silurian–Early Devonian event includes muscovite growth/resetting intervals for gold-bearing zones of the Laurentian Margin, Dashwoods terrane, western Exploits subzone and along the Meelapaeg–Exploits subzones boundary, whereas the younger Devonian interval includes muscovite from gold-bearing zones across the Laurentian Margin, Dashwoods terrane, Exploits Subzone, and Gander Zone.

3. $^{40}\text{Ar}/^{39}\text{Ar}$ plateau ages for detrital muscovite from Wigwam Formation sandstone at the Moosehead prospect (ca. 457.3 ± 1.2 to 453.5 ± 1.1 Ma) suggest the Mount Cormack Complex is an erosional source for the sedimentary units of

the Botwood Basin.

4. Detailed investigations of quartz vein and alteration mineral parageneses, their relationships to gold deposition and systematic geochronological investigations of a number of different mineral species at select auriferous zones is necessary to better constrain the timing of gold mineralization in central and western Newfoundland.

ACKNOWLEDGEMENTS

H. Sandeman wishes to acknowledge the capable field assistance of Matthew Minnett, Jonathon Hull and Kendra Power along with the staff of the Geological Survey of Newfoundland and Labrador for making the research happen smoothly. Dr. Doug Archibald (formerly of Queen's University) provided much early technical assistance with the $^{40}\text{Ar}/^{39}\text{Ar}$ geochronology. Jeremy Powell and two anonymous reviewers provided constructive comments. This is Natural Resources Canada contribution number 20220066.

REFERENCES

- Barr, S. M., Dehler, S. A., and Zsámboki, L. 2014. Connecting Cape Breton Island and Newfoundland, Canada: geophysical modeling of pre-Carboniferous 'basement' rocks in the Cabot Strait area. *Geoscience Canada*, 41, pp. 186–206. <https://doi.org/10.12789/geocanj.2014.41.041>
- Betz, F.J. 1948. Geology and mineral deposits of southern White Bay. Newfoundland Geological Survey Bulletin, 24, 26 p.
- Blackwood, R.F. and Green, L. 1983. Great Gull Lake, Fortune Bay District, Newfoundland. Government of Newfoundland and Labrador, Department of Mines and Energy, Mineral Development Division, Map 82-071, scale 1:50 000.
- Blackwood, R.F. and Kennedy, M.J. 1975. The Dover Fault: western boundary of the Avalon Zone in northeastern Newfoundland. *Canadian Journal of Earth Sciences*, 12, pp. 320–325. <https://doi.org/10.1139/e75-027>
- Churchill, R.A., Wilton, D.H.C., and Evans, D.T.W. 1993. Geology, alteration assemblages and geochemistry of the Duder Lake Gold showings, northeastern Newfoundland. *In* Current Research, Government of Newfoundland and Labrador, Department of Mines and Energy, Geological Survey Branch, Report 93-01, pp. 317–333.
- Clark, D. 1999. First, third and fifth year assessment report on geophysical and diamond drilling exploration for licence 4821 on claim block 8377 and claims 17325 and 17327 in the Bishops Falls area, 3 reports. Newfoundland and Labrador Geological Survey, Assessment File NFLD/2928, 170 p.
- Cohen, K.M., Finney, S.C., Gibbard, P.L., and Fan, J.X. 2013. Updated 2021: The ICS International Chronostratigraphic Chart. *Episodes*, 36, pp. 199–204. <https://doi.org/10.18814/epiugs/2013/v36i3/002>
- Colman-Sadd, S., Hayes, J.P., and Knight, I. 1990. Geology of the Island of Newfoundland. Government of Newfoundland and Labrador, Department of Mines and Energy, Geological Survey Branch, Map 90-01, scale: 1:1 000 000.
- Colman-Sadd, S.P., Dunning, G.R., and Dec, T. 1992. Dunning–Gander relationships and Ordovician orogeny in central Newfoundland: a sediment provenance and U/Pb age study. *American Journal of Science*, 292, pp. 317–355. <https://doi.org/10.2475/ajs.292.5.317>
- Dalrymple, G.B., Alexander, Jr., E.C., Lanphere, M.A., and Kraker, G.P. 1981. Irradiation of samples for $^{40}\text{Ar}/^{39}\text{Ar}$ dating using the Geological Survey TRIGA Reactor. U.S. Geological Survey, Professional Paper 1176, 55 p. <https://doi.org/10.3133/pp1176>
- Dickson, W.L., O'Brien, B.H., and Colman-Sadd, S.P. 2000. Geology of the Botwood map area [NTS 2E/3], central Newfoundland. Government of Newfoundland and Labrador, Department of Mines and Energy, Geological Survey, Open File 2E/03/1067 Version 2.0, map 2000–11, scale 1:50 000.
- Dubé, B. and Lauzière, K. 1997. Gold metallogeny of the Cape Ray fault zone, southwest Newfoundland. Geological Survey of Canada, Bulletin 508, 90 p. <https://doi.org/10.4095/209256>
- Dubé, B., Lauzière, K., and Poulsen, H.K. 1993. The Deer Cove deposit: an example of 'thrust' related breccia-vein type gold mineralization in the Baie Verte Peninsula, Newfoundland: *In* Current Research, Part D, Geological Survey of Canada, Paper 93-1D, p. 1–10. <https://doi.org/10.4095/134265>
- Dubé, B., Dunning, G., and Lauzière, K. 1995. Geology of the Hope Brook Mine, Newfoundland, Canada: A preserved Late Proterozoic high-sulfidation epithermal gold deposit and its implications for exploration. *Economic Geology*, 93, pp. 405–436. <https://doi.org/10.2113/gsec-geo.93.4.405>
- Dube, B., Dunning, G.R., Lauziere, K., and Roddick, J.C. 1996. New insights into the Appalachian Orogen from geology and geochronology along the Cape Ray fault zone, southwest Newfoundland. *Geological Society of America Bulletin*, 108, pp. 101–116. [https://doi.org/10.1130/0016-7606\(1996\)108<0101:NIITAO>2.3.CO;2](https://doi.org/10.1130/0016-7606(1996)108<0101:NIITAO>2.3.CO;2)
- Dunning, G.R. 1987. U/Pb geochronology of the Coney Head Complex, Newfoundland. *Canadian Journal of Earth Sciences*, 24, pp. 1072–1075. <https://doi.org/10.1139/e87-104>
- Dunning, G.R., O'Brien, S.J., Colman-Sadd, S.P., Blackwood, R.F., Dickson, W.L., O'Neill, P.P., and Krogh, T.E. 1990. Silurian Orogeny in the Newfoundland Appalachians. *Journal of Geology*, 98, pp. 895–913. <https://doi.org/10.1086/629460>
- English, M., Sparrow, B., Walsh, M., and Copeland, D.A. 2017. First and seventh year assessment report on data compilation and interpretation for licences 22503M, 23772M, 23774M, 24522M, 24523M and 24524M on claims in the White Bay area, Newfoundland and Labrador. Newfoundland and Labrador Geological Survey, As-

- assessment File 12H/15/2320, 31 p.
- Evans, D.T.W. 1991. Gold metallogeny, eastern Dunnage Zone, central Newfoundland. *In* Current Research, Government of Newfoundland and Labrador, Department of Mines and Energy, Geological Survey Branch, Report 91-01, pp. 301–318.
- Evans, D.T.W. 1993. Gold mineralization in the eastern Dunnage Zone, central Newfoundland. *In* Current Research, Government of Newfoundland and Labrador, Department of Mines and Energy, Geological Survey Branch, Report 93-01, pp. 339–349.
- Evans, D.T.W. 1999. Epigenetic gold mineralization, Baie Verte Peninsula, Newfoundland. *In* Current Research, Government of Newfoundland and Labrador, Department of Mines and Energy, Geological Survey, Report 99-01, pp. 163–182.
- Evans, D.T.W. 2004. Epigenetic gold occurrences, Baie Verte Peninsula, [NTS 12H/09, 16 and 12I/01], Newfoundland. Government of Newfoundland and Labrador, Department of Natural Resources, Geological Survey, Mineral Resource Report 11, 176 p.
- Evans, D.T.W. and Kean, B.F. 2002. The Victoria Lake Supergroup, central Newfoundland – its definition, setting and volcanogenic massive sulphide mineralization. Newfoundland and Labrador Department of Natural Resources, Geological Survey Open File NFLD/2790, 80 p.
- Froude, T. 2019. Sokoman Iron Acquires a 100% Interest in Moosehead Gold Project. Sokoman Iron URL: <<https://sokomanmineralscorp.com/2019/02/01/sokoman-iron-acquires-a-100-interest-in-moosehead-gold-project/>> Press release, February 1, 2019.
- Froude, T. 2021. Sokoman reports first barge-based drill results Moosehead Gold Project, Central Newfoundland; Phase 6 drill Program doubled to 100,000 m. Sokoman Minerals Corp. URL: <<https://sokomanmineralscorp.com/2021/11/10/sokoman-reports-first-barge-based-drill-results-moosehead-gold-project-central-newfoundland/>> Press release November 10, 2021.
- Galley, A.G., Hannington, M.D., and Jonasson, I.R. 2007. Volcanogenic massive sulphide deposits. *In* Mineral Deposits of Canada: A Synthesis of Major Deposit-Types, District Metallogeny, the Evolution of Geological Provinces, and Exploration Methods. *Edited by* W.D. Goodfellow. Geological Association of Canada, Mineral Deposits Division, Special Publication, 5, pp. 141–161.
- Geoscience Atlas. 2022. Newfoundland and Labrador Department of Industry, Energy and Technology, Geological Survey. URL <<https://geoatlas.gov.nl.ca/Default.htm>>, 9 May 2022.
- Heyl, G.R. 1937. The geology of the Sops Arm area, White Bay, Newfoundland. Newfoundland Department of Natural Resources, Geology Section, Bulletin 8, 42 p.
- Honsberger, I.W., Bleeker, W., Kamo, S.L., Evans, D.T.W., and Sandeman, H.A.I. 2019a. A Neoproterozoic age for granodiorite underlying Rogerson Lake Conglomerate; Confirmed Ganderian basement in the Wilding Lake area, central Newfoundland gold district. Government of Newfoundland and Labrador, Department of Natural Resources, Geological Survey, St. John's, Open File 12A/07/1774, 12 p.
- Honsberger, I.W., Bleeker, W., Sandeman, H.A.I., and Evans, D.T.W. 2019b. Structural geology of a gold-bearing quartz vein system, Wilding Lake region, central Newfoundland. *In* Current Research, Newfoundland and Labrador Department of Natural Resources Geological Survey, Report 19-1, pp. 23–38.
- Honsberger, I.W., Bleeker, W., Sandeman, H.A.I., Evans, D.T.W., and Kamo, S.L. 2020a. The Wilding Lake gold prospect, central Newfoundland: a lithological and structural synthesis. Geological Survey of Canada, Open File 8658, 2 sheets. <https://doi.org/10.4095/321490>
- Honsberger, I.W., Bleeker, W., Sandeman, H.A.I., Evans, D.T.W., and Kamo, S.L. 2020b. Vein-hosted gold mineralization in the Wilding Lake area, central Newfoundland: Structural geology and vein evolution. *In* Targeted Geoscience Initiative 5: Contributions to the Understanding of Canadian Gold Systems. *Edited by* P. Mercier-Langevin, C.J.M. Lawley, and S. Castonguay. Geological Survey of Canada, Open File 8712, pp. 179–192. <https://doi.org/10.4095/326020>
- Honsberger, I.W., Bleeker, W., Kamo, S.L., Sandeman, H.A.I., Evans, D.T.W., Rogers, N., van Staal, C.R., and Dunning, G.R. 2022a. Latest Silurian syntectonic sedimentation and magmatism and Early Devonian orogenic gold mineralization, central Newfoundland Appalachians, Canada: Setting, structure, litho-geochemistry, and high-precision U–Pb geochronology. Geological Society of America Bulletin. <https://doi.org/10.1130/GSAB.S.18858221.v1>
- Honsberger, I.W., Wouter, B., Kamo, S.L., Sutcliffe, C.N., and Sandeman, H.A.I. 2022b. U–Pb geochronology of Late Silurian (Wenlock to Pridoli) volcanic and sedimentary rocks, central Newfoundland Appalachians: targeting the timing of transient extension as a prelude to Devonian orogenic gold mineralization. *Atlantic Geoscience*, 58, pp. 215–237. <https://doi.org/10.4138/atlgeo.2022.009>
- Jourdan, F., Verati, C., and Féraud, G. 2006. Intercalibration of the Hb3gr $^{40}\text{Ar}/^{39}\text{Ar}$ dating standard. *Chemical Geology*, 231, pp. 77–189. <https://doi.org/10.1016/j.chemgeo.2006.01.027>
- Kean, B.F. and Jayasinghe, N.R. 1980. Geology of the Lake Ambrose [12A/10]-Noel Pauls Brook [12A/9] map areas, central Newfoundland. Government of Newfoundland and Labrador, Department of Mines and Energy, Mineral Development Division, Report 80-02, 1980, 34 p, 2 maps, scale 1: 50 000.
- Kerr, A. 2006a. Silurian rocks of the Sops Arm group, western Newfoundland: some new food for future digestion. *In* Current Research. Government of Newfoundland and Labrador, Department of Natural Resources, Geological Survey, Report 06-1, pages 91–117.
- Kerr, A. 2006b. Mesothermal gold mineralization in the Silurian Sops Arm group, western Newfoundland: a descriptive and historical overview. *In* Current Research. Government of Newfoundland and Labrador, Department

- ment of Natural Resources, Geological Survey, Report 06-1, pages 61–90.
- Kerr, A. and Selby, D. 2012. The timing of epigenetic gold mineralization on the Baie Verte Peninsula, Newfoundland, Canada: new evidence from Re–Os pyrite geochronology. *Mineralium Deposita*, 47, pp. 325–337. <https://doi.org/10.1007/s00126-011-0375-2>
- Kerr, A. and van Breemen, O. 2007. The timing of gold mineralization in White Bay, western Newfoundland: Evidence from $^{40}\text{Ar}/^{39}\text{Ar}$ studies of mafic dykes that predate and postdate mineralization. *Atlantic Geology*, 43, pp. 148–162. <https://doi.org/10.4138/5623>
- Lincoln, N., Farmer, R., Eccles, R., and Deering, P. 2018. NI 43-101 Technical Report preliminary economic assessment of the Valentine Lake gold project Newfoundland, NL, Canada. Prepared by Lycopodium Minerals Canada Ltd in accordance with the requirements of National Instrument 43-101, “Standards of Disclosure for Mineral Project”, of the Canadian Securities Administrators Qualified Persons. Lycopodium Minerals Canada, 5060 Spectrum Way, Suite 400, Mississauga, Ontario L4W 5N5.
- Lock, B.E. 1969. Paleozoic Wrench Faults in Canadian Appalachians: Chapter 55: Late Orogenic Stratigraphy and Structure: Discussion. *In* North Atlantic: Geology and Continental Drift, AAPG Memoir M 12, 789–790.
- Ludwig, K.R. 2012. Isoplot 4.15: A Geochronological Toolkit for Microsoft Excel: Berkeley Geochronological Center. URL <http://www.bgc.org/isoplot_etc/isoplot/Isoplot4_15files.zip> 19 May 2022.
- Magna Terra Minerals 2022. Maps and Figures -Great Northern. URL <https://www.magnaterraminerals.com/MTT/great_northern_and_viking_projects/4932>, 19 May 2022.
- McDougall, I. and Harrison, T.M. 1988. Geochronology and thermochronology by the $^{40}\text{Ar}/^{39}\text{Ar}$ method. Oxford Monographs on Geology and Geophysics #9, Oxford, United Kingdom, Oxford University Press, 212 p.
- McNicol, V., Squires, G.C., Wardle, R.J., Dunning, G.R., and O'Brien, B.H. 2006. U–Pb geochronological evidence for Devonian deformation and gold mineralization in the eastern Dunnage Zone, Newfoundland. *In* Current Research. Newfoundland and Labrador Department of Natural Resources, Geological Survey, Report 06-1, pp. 43–60.
- Miller, H.G. and Weir, C. 1982. The northwest portion of the Gander Zone — a geophysical interpretation. *Canadian Journal of Earth Sciences*, 19, pp. 1371–1381. <https://doi.org/10.1139/e82-119>
- Minnett, M., Sandeman, H., and Wilton, D. 2012. Geochemistry of the host rocks and timing of gold-electrum mineralization at the Viking Property, Newfoundland. *In* Current Research, Newfoundland Department of Mines and Energy Geological Survey, Report 12-1, pp. 61–84.
- Morgan, J. 2016. First year assessment report on geological, geochemical and trenching exploration for licences 22789M–22800M and 23582M–23587M on claims in the Bishops Falls area, central Newfoundland. Newfoundland and Labrador Geological Survey, Assessment File NFLD/3360, 317 p.
- Myllyaho, J. 2013. First and second year assessment report on prospecting and geochemical and trenching exploration for licences 19543M, 19546M, 19902M, 20193M, 20197M, 20382M and 20417M on claims in the Jacksons Arm area, White Bay, Newfoundland. Newfoundland and Labrador Geological Survey, Assessment File 12H/15/2091, 192 s.
- O'Brien, B.H. 2003. Geology of the central Notre Dame Bay region, northeastern Newfoundland. Government of Newfoundland and Labrador, Department of Mines and Energy, Geological Survey, Report 3-03, 170 p.
- O'Brien, S.J., Dunning, G.R., Dubé, B., O'Driscoll, C.F., Sparkes, B., Israel, S., and Ketchum, J. 2001. New insights into the Neoproterozoic geology of the central Avalon Peninsula (parts of NTS map areas 1N/6, 1N/7 and 1N/3), eastern Newfoundland. *In* Current Research, Government of Newfoundland and Labrador, Department of Mines and Energy, Geological Survey, Report 1-1, pp. 169–189.
- O'Neill, P. and Blackwood, F. 1989. A proposal for revised stratigraphic nomenclature of the Gander and Davidsville groups and the Gander River Ultrabasic Belt, of northeastern Newfoundland. *In* Current Research. Government of Newfoundland and Labrador, Department of Mines, Geological Survey, Report 89-01, pp. 127–130.
- O'Neill, P.P. and Colman-Sadd, S.P. 1993. Geology of the eastern Gander (NTS 2D/15) and western Gambo (NTS 2D/16) map areas, Newfoundland. Government of Newfoundland and Labrador, Department of Mines and Energy, Geological Survey Branch, Report 93-2, 53 p.
- O'Neill, P. and Lux, D. 1989. Tectonothermal history and ^{40}Ar – ^{39}Ar geochronology of the northeastern Gander Zone, Weirs Pond area (2E/1). *In* Current Research. Government of Newfoundland and Labrador, Department of Mines, Geological Survey, Report 89-1, pp. 131–139.
- Onstott, T.C., Phillips, D., and Pringle-Goodell, L. 1991. Laser microprobe measurement of chlorine and argon zonation in biotite. *Chemical Geology*, 90, pp. 145–168. [https://doi.org/10.1016/0009-2541\(91\)90040-X](https://doi.org/10.1016/0009-2541(91)90040-X)
- Piercy, S.J. 2007. Volcanogenic massive sulphide (VMS) deposits of the Newfoundland Appalachians: An overview of their setting, classification, grade-tonnage data, and unresolved questions. *In* Current Research, Newfoundland and Labrador Department of Natural Resources. Geological Survey, Report 07-1, pp. 169–178.
- Pilote, J.-L., Piercy, S., and Mercier-Langevin, P. 2020. Evolution of the seafloor hydrothermal system associated with the Ming VMS deposit, Newfoundland Appalachians, and its controls on base and precious metal distribution. *Mineralium Deposita*, 55. <https://doi.org/10.1007/s00126-019-00899-z>
- Poulsen, K.H., Robert, F., and Dubé, B. 2000. Geological classification of Canadian gold deposits: Geological Survey of Canada, Bulletin 540, 106 p. <https://doi.org/10.4095/211094>

- Ramezani, J., Dunning, G.R., and Wilson, M.R. 2002. Geologic setting, geochemistry of alteration, and U-Pb age of hydrothermal zircon from the Silurian Stog'er Tight gold prospect, Newfoundland Appalachians, Canada. *Exploration and Mining Geology*, 9, pp.171–188. <https://doi.org/10.2113/0090171>
- Reid, W. and Myllyaho, J. 2012. First year assessment report on prospecting, geochemical and trenching exploration for licences 18403M, 19525M, 19530M and 19542M on claims in the Jacksons Arm area, White Bay, Newfoundland. Newfoundland and Labrador Geological Survey, Assessment File 12H/15/2066, 84 p.
- Ritcey, D.H., Wilson, M.R., and Dunning, G.R. 1995. Gold mineralization in the Paleozoic Appalachian Orogen: Constraints from geologic, U-Pb and stable isotope studies of the Hammerdown prospect, Newfoundland. *Economic Geology*, 90, pp. 1955–1965. <https://doi.org/10.2113/gsecongeo.90.7.1955>
- Roddick, J.C. 1983. High precision intercalibration of $^{40}\text{Ar}/^{39}\text{Ar}$ standards. *Geochimica et Cosmochimica Acta*, 47, pp. 887–898. [https://doi.org/10.1016/0016-7037\(83\)90154-0](https://doi.org/10.1016/0016-7037(83)90154-0)
- Rogers, N. and van Staal, C. 2002. Toward a Victoria Lake Supergroup: a provisional stratigraphic revision of the Red Indian to Victoria lakes area, central Newfoundland. *In Current Research, Government of Newfoundland and Labrador, Department of Mines and Energy, Geological Survey, Report 2-1, pages 185–195.*
- Rogers, N., van Staal, C.R., McNicoll, V.J., Pollock, J., Zagorevski, A., and Whalen, J. 2006. Neoproterozoic and Cambrian arc magmatism along the eastern margin of the Victoria Lake Supergroup: A remnant of Ganderian basement in central Newfoundland? *Precambrian Research*, 147, pp. 320–341. <https://doi.org/10.1016/j.precamres.2006.01.025>
- Romer, R.L. and Kroner, U. 2018, Paleozoic gold in the Appalachians and Variscides. *Ore Geology Reviews*, 92, pp. 475–505. <https://doi.org/10.1016/j.oregeorev.2017.11.021>
- Sandeman, H.A.I. 2014a. South Wood Lake (Staghorn) gold prospect lithogeochemical database (map area NTS 12A/04). Government of Newfoundland and Labrador, Department of Natural Resources, Geological Survey, Open File 012A/04/1563, 12 p.
- Sandeman, H.A.I. 2014b. The timing of mineralization at the Jaclyn zone gold deposit, central Newfoundland: constraints from $^{40}\text{Ar}/^{39}\text{Ar}$ studies of white mica alteration adjacent to auriferous quartz veins. *In Current Research, Newfoundland and Labrador Department of Natural Resources Geological Survey, Report 14-1, pp. 63–77.*
- Sandeman, H.A.I. and Dickson, W.L. 2019. An Ordovician, $^{40}\text{Ar}/^{39}\text{Ar}$ step-heating age for fabric-forming hornblende in amphibolite, the Great Bend Complex, central Newfoundland (NTS 2D/5). *In Current Research, Newfoundland and Labrador Department of Natural Resources Geological Survey, Report 19-1, pp. 85–96.*
- Sandeman, H.A.I. and Dunning, G.R. 2016. Preliminary U–Pb geochronology and petrochemistry of volcanic rocks and felsic dykes of the Silurian Sops Arm Group, White Bay, western Newfoundland (NTS 2H/10 and 15). *In Current Research, Newfoundland and Labrador Department of Natural Resources Geological Survey, Report 16-1, pp. 39–69.*
- Sandeman H.A.I. and Spurrell, C. 2020. The Yellow Fox showing: monzogranite-hosted, fracture related antimony–silver–gold mineralization near the eastern margin of the Mount Peyton Intrusive Suite, central Newfoundland (NTS map area 2d/14). *In Current Research, Newfoundland and Labrador Department of Natural Resources Geological Survey, Report 20-1, pp. 179–205.*
- Sandeman, H.A.I., Wilton, D.H.C., Conliffe, J., Froude, T. and O'Driscoll, J.M. 2013. Geological setting, geochronological constraints and the nature of mineralization at the Mosquito Hill (Huxter Lane) gold deposit, central Newfoundland. *In Current Research, Newfoundland Department of Mines and Energy Geological Survey, Report 13-1, pp. 167–188.*
- Sandeman, H.A., Hull, J.R., and Wilton, D.H.C. 2014. Geology, lithogeochemistry and mineralization at the South Wood Lake gold prospect (Staghorn property), Exploits–Meelpaeg subzones boundary, western-central Newfoundland. *In Current Research, Newfoundland and Labrador Department of Natural Resources Geological Survey, Report 14-1, pp. 79–98.*
- Sandeman, H.A.I., Dunning, G.R., McCullough, C.K., and Peddle, C. 2017. U–Pb geochronology, petrogenetic relationships and intrusion-related precious-metal mineralization in the northern Mount Peyton intrusive suite: implications for the origin of the Mount Peyton Trend, central Newfoundland (NTS 2D/04). *In Current Research, Newfoundland and Labrador Department of Natural Resources Geological Survey, Report 17-1, pages 189–217.*
- Sangster, A.L., Douma, S.L., and Lavigne, L. 2008. Base metal and gold deposits of the Betts Cove Complex, Baie Verte Peninsula, Newfoundland. *In Mineral Deposits of Canada: A Synthesis of Major Deposit Types, District Metallogeny, the Evolution of Geological Provinces and Exploration Methods. Edited by W.D. Goodfellow. Geological Survey of Canada, Mineral Deposits Division, Special Volume 5, pp. 703–723.*
- Schaen, A.J., and 34 others. 2021. Interpreting and reporting $^{40}\text{Ar}/^{39}\text{Ar}$ geochronologic data. *Geological Society of America Bulletin*, 133, pp. 461–487. <https://doi.org/10.1130/B35560.1>
- Sparkes, G.W. and Dunning, G.R. 2014. Late Neoproterozoic epithermal alteration and mineralization in the western Avalon Zone: A summary of mineralogical investigations
- Sparkes, G.W., O'Brien, S.J., Dunning, G.R., and Dubé, B. 2005. U-Pb geochronological constraints on the timing of magmatism, epithermal alteration and low-sulphidation gold mineralization, eastern Avalon Zone, Newfoundland. *In Current Research, Government of Newfoundland and Labrador, Department of Natural Resources, Geological Survey, Report 5-1, pp.115–130.*

- and new U/Pb geochronological results. *In* Current Research, Government of Newfoundland and Labrador, Department of Natural Resources, Geological Survey, Report 14-1, pp. 99–128.
- Sparkes, G.W., Ferguson, S.A., Layne, G.D., Dunning, G.R., O'Brien, S.J., and Langille, A. 2016. The nature and timing of Neoproterozoic high-sulphidation gold mineralization from the Newfoundland Avalon Zone: Insights from new U–Pb ages, ore petrography and spectral data from the Hickey's Pond prospect. *In* Current Research, Government of Newfoundland and Labrador, Department of Natural Resources, Geological Survey, Report 16-1, pp. 91–116.
- Spell T.L. and McDougall, I. 2003. Characterization and calibration of $^{40}\text{Ar}/^{39}\text{Ar}$ dating standards. *Chemical Geology*, 198, pp. 189–211. [https://doi.org/10.1016/S0009-2541\(03\)00005-6](https://doi.org/10.1016/S0009-2541(03)00005-6)
- Steiger, R.H. and Jäger, E. 1977. Subcommittee on geochronology: Convention on the use of decay constants in geo- and cosmochronology. *Earth and Planetary Science Letters*, 36, pp. 359–362. [https://doi.org/10.1016/0012-821X\(77\)90060-7](https://doi.org/10.1016/0012-821X(77)90060-7)
- Tuach, J. 1987. Mineralized environments, metallogenesis, and the Doucers Valley Fault Complex, western White Bay: a philosophy for gold exploration in Newfoundland. *In* Current Research, Government of Newfoundland and Labrador, Department of Mines and Energy, Mineral Development Division, Report 87-01, pp. 129–144.
- Tuach, J., Dean, P.L., Swinden, H.S., O'Driscoll, C.F., Kean, B.F., and Evans, D.T.W. 1988. Gold mineralization in Newfoundland: a 1988 review. *In* Current Research, Government of Newfoundland and Labrador, Department of Mines, Mineral Development Division, Report 88-01, pp. 279–306.
- Valverde-Vaquero, P. and van Staal, C. 2001. Relationships between the Dunnage-Gander zones in the Victoria Lake-Peter Strides Pond area. *In* Current Research, Government of Newfoundland and Labrador, Department of Mines and Energy, Geological Survey, Report 1-1, pp. 159–167.
- Valverde-Vaquero, P. and van Staal, C. 2002. Geology and magnetic anomalies of the Exploits-Meelpaeg boundary zone in the Victoria Lake area (central Newfoundland): regional implications. *In* Current Research, Government of Newfoundland and Labrador, Department of Mines and Energy, Geological Survey, Report 2-1, pp. 197–209.
- Valverde-Vaquero, P., van Staal, C.R., van der Velden, A., and Dunning, G.R. 2003. Acadian orogenesis and high-grade metamorphism in the Central Mobile Belt of central Newfoundland. Geological Society of America, Northeastern Section Annual Meeting, Abstracts with Programs, 35(2), 23 p.
- Valverde-Vaquero, P., van Staal, C.R., McNicoll, V., and Dunning, G.R. 2006. Mid–Late Ordovician magmatism and metamorphism along the Gander margin in central Newfoundland. *Journal of the Geological Society, London*, 163, pp. 347–362. <https://doi.org/10.1144/0016-764904-130>
- van der Velden, A.J., van Staal, C.R., and Cook, F.A. 2004. Crustal structure, fossil subduction, and the tectonic evolution of the Newfoundland Appalachians: Evidence from a reprocessed seismic reflection survey. *Geological Society of America Bulletin*, 116, pp. 1485–1498. <https://doi.org/10.1130/B25518.1>
- van Egmond, R. 2004. Second, third, fifth and sixth year assessment report on geological, geochemical and geophysical exploration for licences 6333M, 8378M, 8490M, 8516M and 9162M on claims in the Wood Lake area, southwestern Newfoundland, 2 reports. Newfoundland and Labrador Geological Survey, Assessment File 12A/04/1135, 133 p.
- van Egmond, R. and Cox, E. 2005. Third year supplementary and seventh year assessment report on diamond drilling exploration for licences 6333M and 10608M on claims in the Wood Lake area, central Newfoundland. Newfoundland and Labrador Geological Survey, Assessment File 12A/04/1274, 226 p.
- van Staal, C. and Barr, S.M. 2012. Lithospheric architecture and tectonic evolution of the Canadian Appalachians and associated Atlantic margin. *In* Tectonic Styles in Canada: the Lithoprobe Perspective. Edited by J.A. Percival, F.A. Cook, and R.M. Clowes. Geological Association of Canada, Special Paper, 49, pp. 41–95.
- van Staal, C.R., Sullivan, R.W., and Whalen, J.B. 1996. Provenance and tectonic history of the Gander Margin in the Caledonian/Appalachian Orogen: Implications for the origin and assembly of Avalonia. *In* Avalonian and related Peri-Gondwanan terranes of the Circum-North Atlantic. Edited by R.D. Nance and M.D. Thompson. Geological Society of America Special Paper, 304, pp. 347–367. <https://doi.org/10.1130/0-8137-2304-3.347>
- van Staal, C.R., Valverde-Vaquero, P., Zagorevski, A., Pehrson, S., Boutsma, S., and van Noorden, M.J. 2005a. Geology, King George IV Lake, Newfoundland and Labrador. Geological Survey of Canada, Open File 1665, scale 1:50 000. <https://doi.org/10.4095/221286>
- van Staal, C.R., Valverde-Vaquero, P., Zagorevski, A., Rogers, N., Lissenberg, C.J., and McNicoll, V.J. 2005b. Geology, Victoria Lake, Newfoundland and Labrador; Geological Survey of Canada, Open File 1667, scale 1:50 000. <https://doi.org/10.4095/221287>
- van Staal, C.R., Whalen, J.B., Valverde-Vaquero, P., Zagorevski, A., and Rogers, N. 2009. Pre-Carboniferous, episodic accretion-related, orogenesis along the Laurentian margin of the northern Appalachians. Geological Society, London, Special Publications, 327, pp. 271–316. <https://doi.org/10.1144/SP327.13>
- van Staal, C. R., Zagorevski, A., McNicoll, V. J., and Rogers, N. 2014. Time-transgressive Salinic and Acadian orogenesis, magmatism and Old Red Sandstone sedimentation in Newfoundland. *Geoscience Canada*, 41(2), pp. 138–164. <https://doi.org/10.12789/geocanj.2014.41.031>
- Vry, J.K. and Baker, J.A. 2006. LA-MC-ICPMS Pb–Pb dating of rutile from slowly cooled granulites: confirmation of

- the high closure temperature for Pb diffusion in rutile. *Geochimica et Cosmochimica Acta*, 70, pp. 1807–1820. <https://doi.org/10.1016/j.gca.2005.12.006>
- Waldron, J. W.F. and van Staal, C. R. 2001. Taconian orogeny and the accretion of the Dashwoods block: A peri-Laurentian microcontinent in the Iapetus Ocean. *Geology* 29, 811–814. [https://doi.org/10.1130/0091-7613\(2001\)029<0811:TOATAO>2.0.CO;2](https://doi.org/10.1130/0091-7613(2001)029<0811:TOATAO>2.0.CO;2)
- White, S.E. and Waldron, J.W.F. 2022, Along-strike variations in the deformed Laurentian margin in the Northern Appalachians: Role of inherited margin geometry and colliding arcs. *Earth Science Reviews*, 226. <https://doi.org/10.1016/j.earscirev.2022.103931>
- Whitney, D.L. and Evans, B.W. 2010. Abbreviations for names of rock-forming minerals. *American Mineralogist*, 95, 185–187. <https://doi.org/10.2138/am.2010.3371>
- Williams, H. 1969. Stratigraphy of Botwood Map - Area, Northeastern Newfoundland. Geological Survey of Canada, Open File 113, 104 p. <https://doi.org/10.4095/130357>
- Williams, H., Colman-Sadd, S.P., and Swinden, H.S. 1988. Tectonicstratigraphic subdivisions of central Newfoundland. *In* Current Research, Part B. Eastern and Atlantic Canada, Geological Survey of Canada, Paper 881B, pp. 91–98. <https://doi.org/10.4095/122425>
- Willner, A.P., van Staal, C.R., Zagorevski, A., Glodnye, J., Romere, R.L., and Sudof, M. 2018, Tectonometamorphic evolution along the Iapetus suture zone in Newfoundland: Evidence for polyphase Salinic, Acadian and Neoacadian very low- to medium-grade metamorphism and deformation: *Tectonophysics*, 742–743, p. 137–167. <https://doi.org/10.1016/j.tecto.2018.05.023>
- Wilson, M. and Evans, D.T.W. 1994. Epigenetic gold occurrences in the eastern Dunnage Zone, Newfoundland: preliminary stable-isotope results. *In* Current Research, Government of Newfoundland and Labrador, Department of Mines and Energy, Geological Survey Branch, Report 94-01, pp. 211–223.
- Wonderly, P.F., and Neuman, R.B. 1984. The Indian Bay Formation: fossiliferous Early Ordovician volcanogenic rocks in the northern Gander Terrane, Newfoundland, and their regional significance. *Canadian Journal of Earth Sciences*, 21, pp. 525–532. <https://doi.org/10.1139/e84-057>
- Ybarra, S. 2020. Litho geochemistry and hydrothermal alteration of the Pine Cove orogenic gold deposit, Baie Verte Peninsula, Newfoundland, Canada. Unpublished *M.Sc.* thesis Memorial University of Newfoundland and Labrador, St. John's, Newfoundland and Labrador, 112 p.

Editorial responsibility: Mitchell Kerr