

---

# Atlantic Geoscience Society

## ABSTRACTS

*52<sup>nd</sup> Colloquium & Annual General Meeting 2026*

---

TRURO, NOVA SCOTIA

---

The Atlantic Geoscience Society held its the 52<sup>nd</sup> annual meeting in Truro, Nova Scotia, at the Inn on Prince on February 6–7, 2026. Over 100 presentations were made (83 oral and 28 poster), of which 49 were student presentations. AGS continues to be a at the leading edge of geoscience in Atlantic Canada as well as a great venue for student presentations. On behalf of the society, we thank Colloquium organizers Jared Butler, Lynn Dafoe, Maureen Matthew, Michael Parsons, Rob Raeside, Carla Skinner, and Deanne van Rooyen, as well the numerous session chairs and judges, for facilitating an excellent meeting with about 210 registrants. AGS acknowledges support from the corporate sponsors and partners for the meeting: BGC Engineering, Geoscientists Nova Scotia, Engineers and Geoscientists New Brunswick, New Brunswick Department of Natural Resources, Nova Scotia Department of Natural Resources, Dillon Consulting, and Natural Resources Canada.

In the following pages, we are pleased to publish the abstracts of oral and poster presentations from the meeting representing a wide variety of topics. Best undergraduate and graduate student presentations are indicated by an asterisk in the authorship and a footnote. The meeting included six special sessions: (1) Ongoing developments in coastal and marine geoscience in Atlantic Canada; (2) Sedimentological and paleontological applications in advancing a sustainable energy future; (3) Advances in Appalachian tectonics: insights from field and laboratory studies; (4) Educational outreach; (5) Terrestrial and marine surface processes in Atlantic and Arctic Canada; (6) Advancing critical mineral development in Atlantic Canada. The remaining talks were organized into one general session: Geoscience research developments. Also included were three pre-conference workshops: (1) QAQC methods; (2) Mineral exploration in glaciated terrain; and (3) Building community within the geosciences.

The traditional Saturday evening Awards Banquet and ‘get together’ was highlighted by a talk by Tracy Webb “Geoscience education through the decades-a look back to look forward” and the announcement of the society awards, as well as student prizes for best student poster and oral presentations. Although the abstracts have been edited as necessary for clarity and to conform to Atlantic Geoscience format and standards, the journal editors do not take responsibility for their content or quality.

THE EDITORS

---

## Apatite for knowledge: what can we learn about the Archean Eon from inclusions found in zircon?

---

CHARBEL ABOU-KHALIL AND JILLIAN KENDRICK

*Department of Earth Science, Saint Mary's University,  
Halifax, Nova Scotia B3H 3C3, Canada  
<charbel.abou-khalil@smu.ca>*

One of the most widely debated topics in geology is crustal evolution in the Archean Eon, a time that is poorly understood due to limited preserved crust. There are, however, examples of well-preserved Archean crust, which are largely composed of tonalite-trondhjemite-granodiorite suites (TTGs). TTGs represent some of the earliest continental crust that can be dated back to the Archean and their chemical compositions are typically linked to the depth and tectonic setting of their source. However, the complexity of their magmatic evolution is increasingly being recognized, and unravelling this could provide new insights into their origin. Magmatic zircon in felsic igneous rocks has been shown to contain a record of melt evolution in the form of mineral and melt inclusions, and this approach has not yet been widely applied to TTGs. In this study, we investigate the magmatic evolution of the original melt that formed Mesoproterozoic to Neoproterozoic TTGs and related felsic rocks within the Kapuskasing Uplift in the Superior Province, Canada, by examining inclusions in zircon separates. We found that the mineral inclusions are consistent with the observed mineral assemblages in the rocks and confirm that epidote in the TTGs is magmatic in origin, providing constraints on P-T conditions of crystallization. The F and Cl content of apatite inclusions varies systematically with the age of the sample, and may indicate a shift toward F-bearing, Cl-poor sources over time, which has implications for halogen cycling in the region near the end of the Archean Eon.

---

## Ichnology and sedimentology of the middle Pennsylvanian Minto Formation at Robertson Point, central New Brunswick, Canada: implications for paleobiodiversity, provenance, and paleoenvironmental settings

---

LUKE ALLEN<sup>1</sup>, ROWAN NORRAD<sup>2</sup>, MATTHEW STIMSON<sup>3</sup>,  
OLIVIA KING<sup>4</sup>, DAVE KEIGHLEY<sup>1</sup>, STEVEN HINDS<sup>3,5</sup>,  
ADRIAN PARK<sup>5</sup>, AND BAILEY MALAY<sup>3</sup>

1. *Department of Earth Sciences, University of New Brunswick, Fredericton, New Brunswick B3H 0A4, Canada  
<luke.allen@unb.ca>*
2. *Department of Earth and Environmental Science, Acadia University, Wolfville, Nova Scotia B4P 2R6, Canada*
3. *Natural History Department, New Brunswick Museum,*

*SaintJohn, New Brunswick E2K 1E5, Canada  
4. Department of Earth Sciences, Carleton University,  
Ottawa, Ontario K1S 5B6, Canada*

*5. New Brunswick Department of Natural Resources, Fredericton,  
New Brunswick E3B 5H1, Canada*

Robertson Point, situated along the southern shoreline of Grand Lake, New Brunswick, is a prolific fossil locality that continues to yield a diverse array of tetrapod and invertebrate trace fossils from the middle Pennsylvanian Minto Formation. The Minto Formation was deposited within the Marysville Sub-basin of the Maritimes Basin and is historically thought to represent a tidally influenced peat-forming wetland within back-barrier or delta front depositional settings. New stratigraphic data and interpretations suggest that the stratigraphy at Robertson Point represents fluvial, low sinuosity channel and overbank deposits of large tributary rivers that drained to the northeast, flowing from the crystalline highlands that bordered the southeastern sub-basin margin. These tributaries likely fed into a central flowing axial river system that existed in the center of the basin, resembling a paleoenvironmental setting analogous to the modern-day St. Lawrence River Valley. The tetrapod ichnofaunal assemblage is dominated by the ichnogenera, *Batrachichnus salamandriodes*, *Characichnos* sp., *Limnopus heterodactylus*, *Matthewichnus* sp., *Notalacerta missouriensis*, cf. *Psuedobradypus* sp, and an unidentified, potentially new morphology that resembles both *Notalacerta* and *Dromopus*. Tetrapod trackways provide ichnological evidence as a proxy for biodiversity of terrestrial amphibians (temnospondyls, microsaur, stem tetrapods, anthracosaurs) and early amniotes (reptiles and/or synapsids). Equally as diverse is the invertebrate ichnofaunal assemblage of surface trails (*Gordia*, *Helminthoidichnites*, *Helminthopsis*), infaunal burrows (*Taenidium*, *Planolites*) and walking traces (*Stiaria*, *Diplichnites*, *Kouphichnium*). *Kouphichnium* is one of the few invertebrate traces that is confidently assigned to a limulid or eurypterid trace maker. The presences of this trace could be interpreted as evidence of a distal connection to open water and is thought to represent an upriver excursion to reproduce inland. The ichnofaunal assemblage at Robertson Point is interpreted to have been preserved in a moderately to well drained floodplain paleoenvironment.

---

## Hydraulic controls on pCO<sub>2</sub> damping across river steps and cascades under baseline and alkalinity-dosed conditions

---

ISAAC BAHLER AND SHANNON STERLING

*Hydrology and Climate Change Laboratory, Department of Earth and Environmental Sciences, Dalhousie University,  
Halifax, Nova Scotia B3H 4R2, Canada  
<isaac.bahler@dal.ca>*

Anthropogenic climate change is amplifying carbon-cycle perturbations across aquatic and terrestrial systems, increasing the need for accurate greenhouse-gas accounting. Rivers, though limited in global surface area, exert outsized influence on the carbon and alkalinity balance through coupled weathering and gas-exchange processes and are estimated to emit  $\sim 1.8 \text{ Pg C yr}^{-1}$  as  $\text{CO}_2$ . A key unresolved challenge is the absence of a systematic, scalable approach to quantify  $\text{CO}_2$  evasion from short, high-energy lotic segments - hydraulic “hotspots” where dissolved  $\text{CO}_2$  and exchange rates change sharply over space and time. Discrete features such as steps, cascades, and waterfalls can dominate reach-scale  $\text{CO}_2$  evasion despite occupying negligible surface area, yet prevailing monitoring approaches rarely resolve step-specific contributions, often miss local dynamics occurring over seconds to minutes, and do not yield low-cost proxies suitable for widespread deployment. This knowledge gap is especially consequential for River Alkalinity Enhancement (RAE), a carbon dioxide removal (CDR) strategy in which alkaline minerals, such as calcite, are added to raise alkalinity and dissolved inorganic carbon, lower aqueous  $\text{pCO}_2$ , and promote conversion of atmospheric  $\text{CO}_2$  to bicarbonate for long-term ocean storage. If dosed waters traverse high-energy steps during equilibration, turbulence-driven gas exchange may provide a mechanism for improving removal efficiency and CDR credibility by reversing the gradient of  $\text{CO}_2$  invasion. This research presents the on-going development of an approach to quantify step-resolved  $\text{CO}_2$  evasion by measuring  $\text{pCO}_2$  “damping” across discrete hydraulic steps, with avenues to examine other factors influencing simulated reach evasion. Controlled mesocosm experiments systematically vary hydraulic conditions while collecting high-frequency dissolved  $\text{CO}_2$  observations under baseline and calcite-dosed scenarios, enabling empirical constraints that support scalable hot-spot accounting, improved RAE citing and design.

---

### Toward understanding geological relations in southern New Brunswick, Canada, one U–Pb zircon date at a time

---

SANDRA M. BARR<sup>1</sup>, DEANNE VAN ROOYEN<sup>1</sup>, CHRIS E. WHITE<sup>1</sup>, MICHAEL A. HAMILTON<sup>2</sup>, SUSAN C. JOHNSON<sup>3</sup>, ADRIAN F. PARK<sup>4</sup>, AND JAMES L. CROWLEY<sup>5</sup>

1. *Department of Earth and Environmental Science, Acadia University, Wolfville, Nova Scotia B4P 2R6, Canada*  
<Sandra.barr@acadiau.ca>
2. *Jack Satterly Geochronology Laboratory, University of Toronto, Toronto, Ontario M5S 3B1, Canada*
3. *New Brunswick Department of Natural Resources, Sussex, New Brunswick E4E 5L2, Canada*
4. *New Brunswick Department of Natural Resources, Fredericton, New Brunswick E3B 5H1, Canada*

5. *Department of Geosciences, Boise State University, Boise, Idaho 83725, USA*

Southern New Brunswick is a collage of fault-bounded Neoproterozoic and lower Paleozoic belts/terrane with few original stratigraphic relationships or fossils preserved. Hence ages provided by U–Pb dating of zircon via chemical abrasion-isotope dilution-thermal ionization mass spectrometry (CA-ID-TIMS) and laser ablation-inductively coupled plasma-mass spectrometry are essential for constraining geological relationships. The New River belt contains felsic volcanic rocks and porphyry with ages of ca. 551 Ma, the same as similar rocks in the Avalonian Caledonia belt. Both belts contain Cambrian sedimentary rocks that have been assigned to the Saint John Group, but the New River belt also includes different Cambrian rocks that include rhyolite dated at  $515.96 \pm 0.50 \text{ Ma}$  (CA-ID-TIMS). The New River belt also includes both younger (ca. 550–530 Ma) and older (ca. 630–625 Ma) granitoid plutonic rocks that differ from plutons in the Caledonia terrane. Pre-Mesozoic rocks of Grand Manan Island in the Bay of Fundy have been correlated with the New River belt based on age and rock types, including quartzite units in which the principal zircon age modes are ca. 2.0, 2.5, and 2.75 Ga with no grains younger than Mesoproterozoic. These rocks indicate a large old cratonic source for Ganderia that differs from that for quartzite sources in Avalonia. The Avalonian Caledonia terrane includes continental margin volcanic-plutonic arcs with ages of ca. 690–680 Ma and ca. 620–615 Ma, and voluminous extensional or plume-related magmatism at 551.5–549.5 Ma (Coldbrook Group and related plutons). Overlying Cambrian sedimentary rocks (Saint John Group) have a maximum depositional age of ca. 534 Ma. Southwest of Saint John, the Burchill Wind Project has revealed rock units that may be part of the Carboniferous Partridge Island block, a tectonically emplaced assemblage of Devonian and Carboniferous metasedimentary, sedimentary, and igneous rocks with associated iron oxide-copper-gold (IOCG) mineralization.

---

### Lithochemical discrimination of tungsten-associated intrusive rocks using pXRF, southwestern and central New Brunswick, Canada

---

MARCELA BARRERA-CORTES<sup>1</sup>, DAVID R. LENTZ<sup>1</sup>, AND KATHLEEN G. THORNE<sup>2</sup>

1. *Department of Earth Sciences, University of New Brunswick, Fredericton, New Brunswick E3B 5A3, Canada*  
<marcela.barreraco@unb.ca>
2. *New Brunswick Department of Natural Resources, Fredericton, New Brunswick E2A 7B8, Canada*

In situ portable X-ray fluorescence (pXRF) analyses of immobile elements provide a robust geochemical framework

to minimize the effects of hydrothermal alteration (greisenization) and to assess litho-geochemical differences among intrusive rocks associated with tungsten (W) occurrences at lake George (IG), Flume Ridge (FR), Foster lake (FL), Wildcat (WC), Kedron (KD), and Burnt Hill (BH) in southern and central New Brunswick. Measurements were conducted directly on rock slabs, using the average of three analyses per sample to reduce small-scale heterogeneity, and data were calibrated using certified reference materials. Scatterplots of Nb/Ti and Y/Ti versus Zr/Ti reveal systematic compositional contrasts among the intrusions. KD and WC samples are characterized by elevated Nb, Y, and Zr concentrations relative to IG and FL, indicating highly differentiated felsic magmas for the former, and less evolved, intermediate to felsic compositions for the latter. BH samples are comparatively sparse and scattered, suggesting a heterogeneous and overall intermediate compositional character. The Ti versus Zr diagram highlights distinct fractionation trends, suggesting the involvement of at least two magmatic precursors: a dacitic affinity with moderately high Ti values for FL, IG, and some BH samples, and a rhyolitic affinity with lower Ti contents, encompassing WC, KD, and most BH samples. Additionally, Nb versus Ti relationships indicate contrasting magma source affinities. Granites from WC, FL, IG, and part of BH plot within the syn-collisional granite (syn-COIG) field, whereas KD and most BH samples exhibit higher Nb and Y contents, plotting predominantly within the within-plate granite (WPG) field and the upper ocean-ridge granite (ORG) field, suggesting partial hotspot-influenced magmatic contributions. Overall, these immobile-element relationships indicate that W-associated intrusions along the Central and Southern Plutonic Belt of New Brunswick record contributions from distinct magmatic sources and variable degrees of fractionation, rather than a single homogeneous magma.

---

**A blast furnace from the past: using Renaissance-period smelting techniques and ore sources to assess stable copper isotope fractionation for contact period L'nu artifact provenance studies**

---

ERIC BECHARD<sup>1</sup>, JACOB HANLEY<sup>1</sup>, DELANEY CARTER<sup>1</sup>,  
AND KATIE COTTREAU-ROBINS<sup>2</sup>

1. Department of Earth Science, Saint Mary's University,  
Halifax, Nova Scotia B3H 3C3, Canada  
<eric.bechard@smu.ca>
2. Nova Scotia Museum, 1747 Summer St, Halifax,  
Nova Scotia B3H 3A6, Canada

Heavy metal stable isotope systems (e.g., Pb, Sn, Cu, Fe) are applied in fields like archaeometallurgy to evaluate geologically distinct sources of ore as it relates to the proven-

ance of metallic artefacts and stratigraphic source of ore in historically mined ore systems. However, compared to Pb, Cu isotope systematics have received much less attention. The objective of this study is to elucidate the natural variations in stable Cu isotopes in relevant historical Cu ores (16<sup>th</sup>–17<sup>th</sup> C) and suspected compositionally linked artifacts from sites of L'nu trade, habitation and burial in the Maritimes Basin that date to the same period. Fourteen samples of ore were measured for  $\delta^{65}\text{Cu}$  across five Renaissance mining districts in Sweden, Norway and Germany with a focus on Falun, Sweden, considered to be the most important copper source during this period. Fourteen samples of artifact copper metal previously characterized for stable Pb isotopes were assessed for  $\delta^{65}\text{Cu}$ , including seven L'nu artifacts and seven Renaissance Swedish coins. Five samples of matte were also produced from experiments designed from historical accounts of Renaissance smelting techniques at Falun during its peak production period to quantify Cu isotope fractionation during ore roasting, pulverization and smelting. All artifacts yielded  $\delta^{65}\text{Cu}$  values ranging from 0.16 to 0.60‰ (mean = 0.49‰). This is consistent with the ore samples at Falun (mean = 0.39‰). Excluding two outliers at -0.13‰ and 1.3‰ these fit within expected error ( $\pm 0.08\%$ , 2s). Other historically important mining districts show consistent isotopic signatures given historical context and geology, except for Garpenberg which ranges from 0.02‰ and 3.83‰. Significant outliers in Falun and Garpenberg potentially indicate natural fractionation due to oxidation and weathering of the ore minerals along alteration zones or different ore mineralization. Future work includes characterization of stratigraphic  $\delta^{65}\text{Cu}$  fractionation in the natural Falun system to resolve this issue.

---

**Lithology, geochronology, and tectonic setting of the Rockville Notch Group in the western Meguma terrane, Nova Scotia, Canada\***

---

LESLIE BOATENG-DWOMOH, DEANNE VAN ROOYEN,  
AND CHRIS E. WHITE

*Department of Earth and Environmental Science,  
Acadia University, Wolfville, Nova Scotia B4P 2R6, Canada  
<0309230b@acadiau.ca>*

The Silurian to early Devonian Rockville Notch Group (RNG) consists of sedimentary and volcanic rocks that unconformably overlie the Halifax Group of the Meguma terrane. The RNG occurs north of the Chebogue Point Shear Zone and in the Annapolis Valley area. It consists of, from oldest to youngest, the White Rock, Kentville, and Torbrook/New Canaan formations. In the Torbrook and Wolfville areas, the lowermost White Rock Formation overlies either the Elderkin Brook, Hellsgate Falls, or Lumsden Dam

formations of the Halifax Group. The metamorphic grade of the Halifax Group rocks ranges from sub-greenschist facies in the Wolfville area, to greenschist facies in the Torbrook area. Detailed field mapping and petrographic studies reveal lithological variations in stratigraphy of the RNG. In Fales River, the White Rock Formation consists of a basal rhyolite flows and crystal to crystal lithic tuff overlain by sandstone beds. In the Wolfville area, the base of the White Rock Formation is marked by interbedded sandstone and siltstone. Ironstone occurs in lenses throughout the White Rock Formation. The Kentville Formation in Fales River consists of black slate with abundant graptolite fossils overlain by laminated siltstone with limestone nodules. A complete section of the Torbrook Formation in the Spinney Brook area comprises sandstone and siltstone at both the base and the top of the formation. The middle of the Torbrook Formation includes minor mafic tuff, ironstone, and interlayers of black slate and limestone. These units are contact metamorphosed by the South Mountain Batholith and related plutons which produced a narrow contact aureole with cordierite porphyroblasts preserves at the highest grade. New U–Pb dating of rocks throughout the Rockville Notch Group will yield the maximum depositional ages of the formations, determine how the provenance of the Rockville Notch Group changes throughout its history, and constrain its tectonic affinity.

\*Winner: AGS Graham Williams Award for best graduate poster presentation

---

### Timing and kinematics of deformation relative to contact and regional metamorphism in the Tor Bay area, Nova Scotia, Canada

---

KAREN BROOKMAN, DEANNE VAN ROOYEN,  
AND CHRIS E. WHITE

*Department of Earth and Environmental Science,  
Acadia University, Wolfville, Nova Scotia B4P 2R6, Canada  
<0302569b@acadiau.ca>*

The eastern Meguma terrane consists of Cambrian to Ordovician metasedimentary rocks of the Goldenville and Halifax groups. These units are intruded by Devonian granitoid plutons which have well-developed contact aureoles that overprint regional metamorphic assemblages. In the Tor Bay area, the Goldenville Group comprises the Taylors Head and overlying Beaver Bank formations. The younger Halifax Group consists of the Cunard Formation and the laterally equivalent Cole Harbour Formation. All rocks have a strong near-vertical, E–W striking, composite foliation, axial planar to regional upright, gently east and west plunging, folds. In the Mn-rich Beaver Bank Formation greenschist-facies rocks are characterized by muscovite + garnet (spessartine) + quartz + ilmenite + chlorite ± plagioclase and grade

into lower-amphibolite facies characterized by cordierite + ilmenite ± biotite. In the pelitic Cole Harbour Formation greenschist-facies rocks are characterized by muscovite + quartz + ilmenite + chlorite ± garnet and grade into lower-amphibolite-facies rocks characterized by muscovite + biotite + andalusite ± garnet. In the sulphide-rich Cunard Formation greenschist-facies rocks are characterized by muscovite + chlorite + graphite + quartz + pyrite and lower-amphibolite-facies rocks are characterized by andalusite + cordierite ± biotite. All assemblages show a retrograde metamorphic overprint (sericite + chlorite). Porphyroblasts and overprinting relationships indicate that spessartine garnet was the first major porphyroblast to grow during  $M_1$  regional metamorphism closely associated with  $D_1/S_1$  regional deformation. Cordierite and andalusite porphyroblasts are interpreted as  $M_2$  associated with high heat flow in contact aureoles because they overgrow  $D_1/S_1/M_1$  mineral assemblages and related textures. The  $S_2/D_2$  structures define the dominant E–W-striking regional foliation in which kinematic indicators show both flattening and shear deformation. This  $S_2/D_2$  deformation happened during cooling, because the high-grade assemblages are overprinted by chlorite and sericite (interpreted as  $M_3$  retrograde assemblages) occurring in pressure shadows, boudin necks, and as mantles around porphyroblasts.

---

### Mapping of glacial erratic boulders for mineral exploration in southwestern Nova Scotia, Canada, using ultra-high-resolution LiDAR

---

SYED BUKHARI<sup>1</sup>, NICK EYLES<sup>1</sup>, ROGER PAULEN<sup>2</sup>,  
LAVANECHA CHANDRAN<sup>1</sup>, AND DENISE BRUSHETT<sup>3</sup>

- 1. Department of Physical and Environmental Sciences, University of Toronto-Scarborough, Toronto, Ontario M1C 1A4, Canada  
<sajidb.bukhari@mail.utoronto.ca>*
- 2. Geological Survey of Canada, Natural Resources Canada, Ottawa, Ontario K1A 0G1, Canada*
- 3. Nova Scotia Department of Municipal Affairs, Halifax, Nova Scotia B3J 2M4, Canada*

Mineral exploration in glaciated terrains is dependent on a thorough understanding of the history and mode of flow of Pleistocene mid-latitude ice sheets to successfully identify potential mineralized targets in bedrock partially or completely covered by glacial sediments. Mapping of glaciated terrains and exploration using boulder trains and mineralized dispersal trains on the glaciated crystalline shields of Canada and Fennoscandia is challenged by vegetative covers and remote sensing data of inadequate resolution. New generation LiDAR data and mapping techniques provide an excellent solution by allowing the removal of vegetation to generate bare earth digital elevation models and identify glacial erratic boulders. Targeted drone-

based LiDAR mapping was carried out at an area of known lithium and tin mineralisation obscured by thick forest cover in Nova Scotia, which has been affected by several Pleistocene glaciations, most recently by the Appalachian Glacier Complex some 20 000 years ago. The methodology developed herein permitted successful mapping of erratic boulders and their spatial density. This approach can now be scaled up for regional mineral exploration projects and for geomorphic mapping of modern and ancient glaciated terrains to determine former ice flow trajectories and landform evolution in Atlantic Canada.

---

**Turning up the heat: garnet chemistry and phase equilibria constraints on high-temperature, low-pressure metamorphism in the southwestern Meguma terrane, Nova Scotia, Canada**

---

JARED P. BUTLER<sup>1</sup>, ANTOINE GODET<sup>2</sup>, JAMIE A. CUTTS<sup>3</sup>,  
AND DAWN A. KELLETT<sup>1</sup>

1. Geological Survey of Canada, Natural Resources Canada, Dartmouth, Nova Scotia B3B 1A6, Canada  
<jared.butler@nrcan-rncan.gc.ca>
2. Geological Survey of Canada, Natural Resources Canada, Québec, Québec G1K 9A9, Canada
3. Geological Survey of Canada, Natural Resources Canada, Ottawa, Ontario K1A 0E8, Canada

Staurolite-grade metapelites in southwestern Nova Scotia record high-temperature, low-pressure metamorphism associated with Devonian orogenesis. However, quantitative constraints on the timing and conditions of this metamorphism are lacking, limiting our ability to relate it to the broader tectonic and metallogenic evolution of the Meguma terrane. We present preliminary results from a petrological and geochronological study designed to address this gap, focusing on the pressure-temperature evolution of metapelites from the early Silurian White Rock Formation, north of the Chebogue Point shear zone in the Yarmouth area. Petrographic observations show that M<sub>1</sub> metamorphism produced syn- to post-tectonic garnet ± staurolite ± andalusite assemblages. Overprinting fabrics and chlorite strain fringes on M<sub>1</sub> porphyroblasts represent later D<sub>2</sub>/M<sub>2</sub> deformation and metamorphism. M<sub>1</sub> garnet preserves prograde major element zoning with rimward decreasing spessartine and increasing almandine, relatively constant pyrope and grossular contents, and narrow, higher-grossular outer rims. Concentric trace element zoning in garnet is superimposed on patchy, matrix-inherited patterns, suggesting limited post-growth element mobility. Elevated REE concentrations associated with higher-grossular outer rims imply growth during prograde break-

down of allanite or other Ca- and REE-bearing phases. Phase equilibria modeling in the MnNCKFMASHTO system using Theriak Domino indicates garnet core growth at ~3.0–3.5 kbar/525–540°C followed by an increase in P-T conditions to ~4.5–5.5 kbar/550–615°C, with outer rims recording somewhat higher conditions. Efforts are underway to constrain the timing of metamorphism by garnet Lu-Hf and U-Pb monazite geochronology. Comparable high-temperature, low-pressure metamorphism has been documented elsewhere in the Meguma terrane. Although some occurrences are spatially associated with late Devonian plutons, textural evidence and field relations suggest that these are not contact metamorphic rocks but instead record a regionally elevated geothermal gradient (~40°C/km) that preceded pluton emplacement. Such conditions are difficult to reconcile with crustal thickening alone, implying that other processes, such as lithospheric thinning or delamination, played an important role in Devonian orogenesis in the Meguma terrane.

---

**Lithology and thickness variations in Lower Pennsylvanian rocks of the southern Maritimes Basin, Canada**

---

JOHN CAREY<sup>1</sup>, PAUL DURLING<sup>1</sup>, PETER GILES<sup>1</sup>, ZELIA PEREIRA<sup>2</sup>, MARCIA MENDES<sup>2</sup>, VANIA CORREIA<sup>1</sup>, AND  
ROB FENSOME<sup>1</sup>

1. Geological Survey of Canada, Natural Resources Canada, Dartmouth, Nova Scotia B3B 1A6, Canada  
<john.carey@nrcan-rncan.gc.ca>
2. Laboratório Nacional de Energia e Geologia, Zambujal Neighborhood, Alfragide 2610-999 Amadora, Portugal

The Hastings Uplift is an uplifted basement block in northern Nova Scotia and southeastern New Brunswick, southeast of the Sackville Basin, north of the Athol syncline and across the Northumberland Strait from Prince Edward Island. Understanding of the stratigraphy of the Hastings Uplift and its relationship to other parts of the Maritimes Basin has been hampered by limited outcrop and biostratigraphic data. New palynological data from the Hastings No. 1 well and a nearby borehole provide insights to the thickness and facies patterns in the lower Pennsylvanian rocks. In the Hastings well, the Mississippian Mabou Group is overlain by a sand-dominated succession with interbedded grey siltstone approximately 1650 m thick most of which belongs to the *Raistrickia fulva* zone (late Bashkirian). These rocks resemble but are largely younger than the Boss Point Formation in the well-studied stratigraphic succession at Joggins in the Athol syncline; there, the *R. fulva* zone comprises a red and grey clastic succession with intercalated coal-bearing strata. The Hastings succession more closely resembles that in the Sackville Basin where the

Boss Point Formation is estimated to be up to 2400 m thick, spans the *R. saetosa* (early Bashkirian) and *R. fulva* zones, and coal measures are absent. The Bashkirian succession in western Cape Breton is also thick (~2200 m) and broadly resembles that of the Athol syncline. In contrast, Bashkirian rocks in Prince Edward Island are comparatively thin (<1000 m thick), which may also be the case in the Tatamagouche syncline southeast of the Hastings Uplift. These results suggest that Bashkirian strata are thicker proximal to ancestral basement than nearer the basin center, possibly due to greater erosion or a longer hiatus in late Bashkirian time. The observed lithology and thickness variations have important implications for potential carbon sequestration in rocks of this age.

---

**Analysis of reef island building, sediment production, and island elevation, Utila, Honduras**

---

KASIA CHMIEL

*Department of Geological Sciences, University of Manitoba,  
Winnipeg, Manitoba R3T 2N2, Canada  
<kasiachmiel27@outlook.com>*

Coral reef islands are amongst the most diverse yet vulnerable ecosystems on Earth. They yield important ecosystem services, providing habitat for both animals, including endemic and threatened species, and humans. In some cases, they house entire nations, supporting the livelihoods of billions of people globally. As these landforms are typically below 5 m above mean sea level (MSL), they are under threat from inundation due to sea level rise (SLR) and increasing storm events. Reef island dynamics are highly understudied, particularly within the Caribbean region, highlighting the need for a deeper understanding of potential responses to global warming induced oceanographic changes. Reef island longevity is a function of its 3D profile; hence, this thesis assesses the relationship between sediment production rates and elevation of three reef islands: Morgans Cay (MC), Water Cay (WC), and South West Cay (SWC), in the Caribbean Sea off the coast of Utila, Honduras. Sediment transport dynamics are identified as driving process behind island geomorphology therefore transport processes including currents and waves are investigated. To calculate sediment production rates, sediment budget surveys in three zones around each island, seagrass, patch reef and reef crest, are the main data collection method. Current and wave data are retrieved, in addition to elevation data by topographic surveys of each island. Results revealed no significant relationship between sediment production rates and island elevation. MC had the highest production yet

was the smallest island whereas WC exhibited the lowest production rates yet was the largest island. Oceanic data suggest transportation of island grade material away from MC north-westerly towards SWC. WC revealed extensive patches of dead coral, possibly causing rapid influxes of sediment grade material to the island. Additionally, significant anthropogenic disturbances and manmade sediment trapping features were observed at WC. The energy transition will reduce the pressures of global warming and SLR on future carbonate production and sedimentological regimes, allowing reef island accretion to keep pace with SLR.

---

**Understanding of geohazards and constraints for development of Nova Scotia, Canada's designated offshore Wind Energy Areas**

---

JADE CHUNG<sup>1</sup>, VICTOR WEBER<sup>2</sup>, MAEVE STORY<sup>2</sup>,  
AND DIANE BAXTER<sup>2</sup>

1. BGC Engineering Inc., Vancouver British Columbia V6Z 0C8, Canada <jchung@bgcengineering.ca>
2. Cathie Group, Newcastle upon Tyne NE1 3NP, UK

The Cathie Group have collaborated with BGC Engineering to complete a regional desktop study of Nova Scotia's designated offshore Wind Energy Areas (WEAs). located on the continental shelf, three WEAs form part of land Tenure Process NS25-1R launched in October 2025 by the Canada-Nova Scotia Offshore Energy Regulator: Sydney Bight, Middle Bank, and French Bank. This study uses public datasets to assess geological, geospatial, and anthropogenic conditions, then provides preliminary development recommendations from global offshore wind experience. The study includes: a regional geological data review to understand depositional history; a compilation of specific site conditions including bathymetry, seabed morphology, and seabed conditions; an identification of potential geohazards; and a preliminary assessment of anthropogenic features and metocean parameters that may influence site suitability. These data are then used to develop geoscience recommendations for additional geotechnical and geophysical surveys to reduce uncertainty in site understanding, to support the development of ground models for effective risk mitigation, and to identify applicable complementary studies (e.g., sediment mobility, cable burial risk). The study also identifies preliminary development considerations such as the suitability of the seabed and subsurface for industry-specific foundations, anchors and cables, and anticipated challenges for engineering design and construction. Screening of bathymetry data across the WEAs indicates that water depths would be consistent with the development of either bottom-fixed (e.g., Middle Bank) or

anchored floating technologies (e.g., French Bank). Geoscience data review suggests that area-wide survey strategies should aim to collect additional primarily coarse seabed sediments at the Sydney Bight to reduce risk for foundation design. Soil investigations within the French Bank will be required to reduce uncertainty as highly variable subsurface conditions are interpreted with potential for bedrock outcropping. last, the Middle Bank has potential for scour depressions and large ripples, of which a high-resolution geophysical dataset would assist understanding.

---

### Evaluating subsurface carbonate reservoirs for CO<sub>2</sub> storage

---

HILARY CORLETT

*Department of Earth Sciences, Memorial University of Newfoundland, St. John's, Newfoundland and Labrador A1C 5S7, Canada <hilary.corlett@mun.ca>*

Meeting our global targets for reduction of greenhouse gases requires us to cut back on emissions through shifting to different energy sources, but also to store CO<sub>2</sub> that is produced through our current production. Canada runs four of the largest seven subsurface geological CO<sub>2</sub> storage operations in the world in Alberta and Saskatchewan. Three of the four involve injection of liquid CO<sub>2</sub> into carbonate reservoirs. There are many things that can make carbonate reservoirs ideal candidates for CO<sub>2</sub> storage, but there are considerations to take into account when evaluating permeability, porosity, and reactivity in carbonates. In this talk, I review the projects that currently operate in western Canada and discuss an integrated approach to the evaluation of carbonate reservoirs for CO<sub>2</sub> storage using various analytical methods. Examples of carbonate reservoirs and analogues will be included from the Western Canada Sedimentary Basin and closer to home in the offshore Newfoundland Jeanne D'Arc Basin. Evaluation of the petrophysical properties of carbonates requires knowledge of the changes in biota on the carbonate platform through geological time that result in variable carbonate environments and therefore, complex textures. Assessment of the complex diagenetic history of highly reactive carbonate sediments from the time they are deposited to present day is also critical in understanding the physical properties of the rocks, their potential reactivity, and possible fluid flow and plume migration pathways linked to prior tectonism and diagenesis.

---

### Lived experience navigating the geosciences and field-based research with a physical disability: the good, the bad, and the things that drive me mad

---

HILARY CORLETT

*Department of Earth Sciences, Memorial University of Newfoundland, St. John's, Newfoundland and Labrador A1C 5S7, Canada <hilary.corlett@mun.ca>*

The first reaction I had to telling my family that I was going to go into geology was “oh, big surprise”, but the first reaction I had to telling my university advisors was “you cannot be in geology if you cannot participate in field work”. The reaction from my family is because I come from a family that is oddly filled with geologists and they knew how much I loved being outdoors and trying to understand the natural environment. The reaction from my university is because I told them I had a physical disability that can limit some of my mobility. I had not told them I could not do fieldwork, but rather that I would need to figure out my own way to do it. What they told me was a difficult thing to hear, but my aunt, who was one of the first female geology professors in the country, told me to persevere, and that she was sure I would be able to succeed. In this talk, I would like to share some of my experiences from over 20 years in the geosciences. I will also present some of the amazing adaptations that are being developed by various groups in the geosciences to promote inclusion in the field and the lab for people with differing abilities.

---

### Early Cretaceous palynoevent scheme for the Scotian Margin, Canada: application to offshore geological carbon storage and other energy transition studies

---

VÂNIA CORREIA<sup>1</sup>, ROBERT A. FENSOME<sup>1</sup>, LYNN T. DAFOE<sup>1</sup>, R. ANDREW MACRAE<sup>2</sup>, GRAHAM L. WILLIAMS<sup>1</sup>, CARLA SKINNER<sup>1</sup>, JOHN CAREY<sup>1</sup>, and RICARDO L. SILVA<sup>3</sup>

1. *Geological Survey of Canada, Natural Resources Canada, Dartmouth, Nova Scotia B3B 1A6, Canada <vania.correia@nrcan-rncan.gc.ca>*

2. *Department of Earth Science, Saint Mary's University, Halifax, Nova Scotia B3H 3C3, Canada*

3. *Department of Earth Sciences and BETY Lab, University of Manitoba, Winnipeg, Manitoba R3T 2N2, Canada*

Organic-walled microfossils (palynomorphs), including dinoflagellate, spores, and pollen, are extensively recovered from subsurface wells and boreholes, often when other fossil groups are scarce or difficult to extract from limited sample material; they are a standard biostratigraphic tool for relative age control. The Scotian Margin, located off Atlantic

Canada, has been studied biostratigraphically over the past 50 years, and palynoevents (e.g., inceptions, extinctions, and acmes of species) have been critical to understanding the geological history of this area and associated petroleum systems. However, the Late Triassic–Early Cretaceous interval lacks a detailed event-stratigraphic framework. With the urgent need to globally shift to more environmentally sustainable energy and Canada's current aim to achieve net-zero emissions by 2050, geological studies to support the development of renewable energy, such as offshore geological carbon storage (GCS) and offshore wind, are being actively advanced on the Scotian Margin. Assessing the potential of GCS requires a comprehensive knowledge of the subsurface geology, including age constraints and assessment of depositional paleoenvironments. We carried out palynological analyses of 115 samples from the Lower Cretaceous conventional cores of Panuke B-90, comprising the Upper Member of the Missisauga Formation and the Logan Canyon Formation, a key interval for ongoing GCS investigations. We have identified several palynoevents from four lithological intervals reflecting changes in the paleoenvironmental setting. Additionally, we have tied our results to selected palynoevents from previous biostratigraphic studies, resulting in a high-resolution Mesozoic palynoevent scheme for the Scotian Margin. These palynoevents are further integrated with other analyses in an ongoing event-stratigraphic framework. Palynology, combined with other disciplines, is effective in providing age-control refinements for the Scotian Margin, which is essential for establishing regional correlations, and for paleoenvironmental interpretations that help understand the nature of reservoir bodies in GCS studies.

---

**Bedrock occurrences tied to wind energy areas offshore  
Nova Scotia, Canada: linking seabed samples to  
subsurface and onshore geology**

---

LYNN T. DAFOE<sup>1</sup>, MARK E. DEPTUCK<sup>2</sup>, ROBERT A. FENSOME<sup>1</sup>, D. CALVIN CAMPBELL<sup>1</sup>, CHRIS E. WHITE<sup>3</sup>, JORDAN B.R. EAMER<sup>1</sup>, STÉPHANIE LARMAGNAT<sup>4</sup>, VÂNIA CORREIA<sup>1</sup>, QINGMOU LI<sup>5</sup>, KEVIN MACKILLOP<sup>1</sup>, ZÉLIA PEREIRA<sup>6</sup>, AND SEAN KELLY<sup>1</sup>

1. Geological Survey of Canada, Natural Resources Canada, Dartmouth, Nova Scotia B2Y 4A2 Canada  
<lynn.dafoe@nrcan-rncan.gc.ca>
2. Canada-Nova Scotia Offshore Energy Regulator, Dartmouth, Nova Scotia B3B 1W2 Canada
3. Department of Earth and Environmental Science, Acadia University, Wolfville, Nova Scotia B4P 2R6, Canada
4. Geological Survey of Canada, Natural Resources Canada, Québec City, Quebec G1K 9A9 Canada
5. Geological Survey of Canada, Natural Resources Canada, Ottawa, Ontario K1A 0E8 Canada

6. Laboratório Nacional de Energia e Geologia,  
Rua da Amieira, S. Mamede de Infesta, Porto, Portugal

Wind energy development in offshore Nova Scotia requires an understanding of seabed conditions and seabed geology to guide infrastructure design, with exposed and shallowly buried bedrock posing a challenge for conventional foundations. Three recently designated wind energy areas overlie varied bedrock: a Mesozoic–Cenozoic Scotian Basin passive margin succession below Middle Bank, Carboniferous Sydney Basin strata below Sydney Bight, and Meguma terrane rocks beneath French Bank. Our study focusses on archival and recent dredge and grab samples of bedrock and subcrop geology within these areas, with assessment of lithology, lithostratigraphy, age, paleoenvironments, geochemistry, and key petrophysical and mechanical properties. Where possible, samples are integrated with multibeam bathymetry, shallow seismic, potential fields, borehole data, and onshore bedrock geology. Adjacent to Middle Bank, two sample stations predominantly record bedded and bioturbated sandstone representing possible fluvial, estuarine, shoreface, and inner shelf settings, with rare shelfal limestone. Subcrop mapping from industry seismic and borehole information shows that the sandstone samples are likely Logan Canyon Formation. Similar lithologies in the nearby Dover A-43 well confirms this, with rare limestone likely correlating to the overlying Petrel Member of the Dawson Canyon Formation. In the Sydney Bight area, Carboniferous–Permian samples include red, grey and greenish sandstone and siltstone with current ripples, planar lamination, cross bedding and/or mottling representing fluvial channel, crevasse splay and floodplain paleoenvironments. Extrapolation of onshore structure and geology combined with borehole data indicates a possible shallow subcrop transition between the Morien and Pictou groups in northeast plunging folds. In the French Bank area, bedrock is imaged in multibeam data and samples confirm the presence of laminated metasandstone consistent with Goldenville Group in tight east–west oriented folds. Bedrock subcrop mapping and seabed samples thus provide key insights into occurrences impacting future offshore wind development, both within designated wind energy areas and on the broader continental shelf.

---

**Distinguishing Tonian and Ediacaran rocks in the  
northern Cobequid Highlands, Nova Scotia, Canada**

---

TIERNAN DAVIES, SANDRA M. BARR, CHRIS E. WHITE,  
AND DEANNE VAN ROOYEN

Department of Earth and Environmental Science,  
Acadia University, Wolfville, Nova Scotia B4P 2R6, Canada  
<0304939d@acadiau.ca>

The Cobequid Highlands in northern mainland Nova Scotia remains one of the most enigmatic components of Avalonia in the northern Appalachian orogen. The highlands are divided by faults into the Bass River, Mount Ephraim, and Jeffers blocks, all of which contain small areas of Neoproterozoic rocks with uncertain relations both to one another and to other parts of Avalonia. Although initially interpreted to be mainly Ediacaran, recent U–Pb zircon dating has shown that both Tonian and Ediacaran volcanic and plutonic rocks are present. However, because they formed in similar volcanic arc settings, establishing criteria to distinguish rocks of the two ages without U–Pb zircon dating at every outcrop is challenging. Tonian ages of 755 Ma to 725 Ma are well established for volcanic rocks of the Dalhousie Mountain Formation and Mount Ephraim Plutonic Suite in the Mount Ephraim block in the eastern highlands. Recent studies of the central Jeffers block reveal that it consists mainly of tuffaceous volcanic rocks intruded by two granodioritic plutons, one Tonian (Bulmer Brook pluton, ca. 740 Ma) and the other Ediacaran (Lily Pond pluton, ca. 626 Ma). Although trace element characteristics enable these two dated plutons to be distinguished, subdividing the host volcanic rocks, all currently assigned to the Ediacaran Jeffers Group, is difficult. U–Pb dating and chemical characteristics of three tuffaceous rocks in the New Annan area support the interpretation that the northern part of the central block of the Jeffers Group is Tonian and related to the Dalhousie Mountain Formation in the Mount Ephraim block, whereas the southern part may be Ediacaran and related to dated volcanic and plutonic rocks of that age in the western part of the Jeffers block.

---

**Separating advective from conductive heat transport signatures in Distributed Temperature Sensing (DTS) data for dam seepage monitoring**

---

EMANUEL DE GANTE CARRILLO, KARL BUTLER,  
AND KERRY MACQUARRIE

*Department of Earth Sciences, University of New Brunswick,  
Fredericton, New Brunswick E3B 5A3, Canada  
<manuel.degante@unb.ca>8.12*

Fibre Optic Distributed Temperature Sensing (DTS) systems have been installed in water-retaining dams throughout the world to measure temperature and detect leaks or concentrated seepage. Where concentrated water seepage occurs temperature anomalies will be transported into the dam structure by advection, distorting the normal temperature pattern. This study developed a suite of data processing and visualization techniques designed to identify temperature anomalies suggestive of anomalous seepage within more than a decade of Distributed Temperature Sen-

ing (DTS) data collected at the Mactaquac Dam. The dataset is dominated by strong seasonal signals arising from conductive heat exchange with air- and reservoir-exposed surfaces, which obscure subtler advective signatures. Two complementary approaches are demonstrated to enhance anomaly detection. The first computes the temperature gradient along the DTS cable and applies median filtering to reduce systematic noise, effectively highlighting short-wavelength thermal variations. The second, more advanced method introduces the DBM (deGante-Butler-MacQuarrie) algorithm, which employs a modified Karhunen–Loève transform adapted from seismic data processing. Conceptually similar to Principal Component Analysis, this approach estimates and subtracts depth-dependent conductively transported temperature variations, enabling the extraction of more subtle thermal patterns associated with potential seepage. In addition, novel visualization strategies are presented, including offset temperature-depth and gradient-depth displays inspired by seismic trace plotting, as well as a new method for visualizing time series as a function of depth. Together, these techniques significantly improve the interpretation of seasonal behavior and the identification of subtle anomalies within long-term DTS data sets which are becoming increasingly common in dam seepage/leakage monitoring and other hydrological applications.

---

**Chlorite compositions, prograde metamorphism, and gold-related trace elements**

---

LAUREN DIPETRO AND RICHARD COX

*Department of Earth and Environmental Sciences, Dalhousie  
University, Halifax, Nova Scotia B3H 4R2, Canada  
<kdipietro2000@icloud.com>*

This project focuses on volatile-metalloid (VM) element (S, Se, Te, Sb, Bi, Hg, As, etc) behavior during progressive metamorphism of pelitic rocks from the West Highlands of Scotland. The metamorphism is represented by chlorite-mica schists to garnet-muscovite schists, with at least two stages of deformation and variable retrograde reactions present. VM-elements are commonly related to orogenic gold mobilization, and the compositions of sheet silicates have been used as indicators of gold deposition. In polymetamorphic terranes, different deformation events may be responsible for the reactions releasing volatile and metalloid elements. Microstructures preserved by sheet silicates are widely used to examine P–T conditions and deformation. This project investigates P–T conditions and trace-element variations during multistage deformation in samples from the West Highlands which were mapped and collected in May of 2025. The research will involve, (i) SEM

and petrographic analysis of chlorite schists and chlorite-altered retrograde garnets with identification of chlorite growth zoning during varying metamorphic conditions (ii) X-ray mapping and analyses of major elements in the sheet silicates to produce P-T-d maps, and in the future, (iii) trace-element analysis using LA-ICP-MS to determine which deformation event(s) released volatile, gold-related elements, and finally, (iv) determine the grades of metamorphism that release the majority of volatile elements.

---

**Cryptic microbial carbonates of the basal Windsor Group: a sedimentologic and petrographic reappraisal of the Macbeth Brook Formation, Cape Breton Island, Nova Scotia, Canada**

---

ZAILYNNE DURANT<sup>1,2</sup> AND EDWARD J. MATHESON<sup>2</sup>

*1. Department of Earth and Environmental Science, Acadia University, Wolfville, Nova Scotia B4P 2R6, Canada  
<zailynne@yahoo.ca>*

*2. Department of Math, Physics and Geology, Cape Breton University, Sydney, Nova Scotia B1M 1A2, Canada*

The Macbeth Brook Formation of the Sydney Basin is a laterally extensive and understudied component of the basal Windsor Group. Existing interpretations are based primarily on core observations, with limited detailed outcrop analysis and petrography. As a result, the stratigraphic and paleo-environmental relationships between the Macbeth Brook Formation and coeval microbial reef complexes of the geographically proximal and laterally equivalent Gays River Formation, as well as the more widely known Macumber Formation laminated limestone, remain incompletely understood. The Macbeth Brook Formation has been previously described as variable siliciclastics in bedded limestones that are stromatolitic or thickly bedded to massive with relatively sparse fossil content. Here we re-evaluate key outcrops of the Macbeth Brook Formation in the Coxheath Hills, along East Bay, and at the northern end of the Boisdale Hills, together with Gays River Formation outcrops near the transition between units. Petrographic analysis reveals that the Macbeth Brook Formation is dominated by microscopic microbial fabrics, including laminated, oncolitic, cryptic, and irregular textures that are difficult to recognize at outcrop scale. Fossil material is more common than previously reported, though largely present as thin-section-scale components intergrown with microbial fabrics, dominated by molluscs and poorly preserved enigmatic and problematica forms. These textures indicate in situ carbonate accumulation, rather than extensive allochthonous redeposition, with growth of significant microbial material coating individual fossil grains providing novel insight in fluctuating conditions

between those supportive of metazoan vs. microbial communities. Comparison with nearby Gays River reefal successions indicates a closer paleoenvironmental association than previously recognized in a spectrum of shallow-marine, microbially influenced carbonate environments. This work further highlights the need for re-evaluation of paleogeographic and relative water depth interpretations of the onshore Sydney Basin Windsor Sea.

---

**Organic geochemical analysis of potential intra-salt Early Jurassic oil shows of the Scotian Basin, Canada**

---

STEPHANIE ERHUNMWUNSEE, TODD VENTURA, AND  
ANDREW MACRAE

*Department of Earth Science, Saint Mary's University,  
Halifax, Nova Scotia B3H 3C3, Canada  
<stephanie.omoeghie.erhunmwunsee@smu.ca>*

Well-completion reports document numerous putative oil shows in salt-bearing units and associated carbonates within the Early Jurassic, Argo Formation of the Scotian Margin in the Orpheus Graben. These oil shows were, however, inferred from odour, fluorescence, and visual staining during drilling operations and require definitive confirmation by geochemical analysis of core plugs, sidewall cores, and cuttings. Unfortunately, these materials are often compromised by contamination from hydrocarbon in drilling fluids. This study evaluates whether oil-stained lithologies and salt-hosted fluid inclusions preserved in contaminated cuttings can be extracted and geochemically fingerprinted to distinguish indigenous in situ or migrated hydrocarbons from drilling-fluid contaminants. Eleven Early Jurassic carbonate-, siliciclastic-, and halite-dominated cuttings samples, along with two sidewall cores from the Argo F-38 well, were analyzed. Due to extensive drill-mud contamination, an experimental series of sequential solvent washes was applied before sample powdering and extraction. Hydrocarbon recoveries decreased exponentially with successive washes, indicating progressive removal of contaminant hydrocarbons. Apolar fractions were analyzed using comprehensive two-dimensional gas chromatography with flame ionization detection (GC×GC-FiD). Subtracted chromatograms were used to characterize drilling-fluid hydrocarbon signatures and isolate potential native hydrocarbon fingerprints. The results will assess whether indigenous hydrocarbons are present and, where confirmed, constrain their origin and maturity, providing new geochemical constraints on possible hydrocarbon generation within the Scotian Margin.

---

**Results of the Geoheritage Summer Institute  
and Geodiversity Day in the Cliffs of Fundy  
UNESCO Global Geopark**

---

TIM FEDAK

*Nova Scotia Museum of Natural History,  
Halifax, Nova Scotia B3H 3A6, Canada  
<tim.fedak@novascotia.ca>*

Three one-hour online seminars were delivered for the 2025 Geoheritage Summer institute (GSI) at noon on Fridays, June 6, July 4, and August 1. Fifty multidisciplinary participants were invited, representing geoscience, humanities, tourism, and economics, from universities, government, industry, and community members within the Cliffs of Fundy UNESCO Global Geopark. Thirty individuals (60%) registered through Zoom. The 2025 GSI was designed to strengthen engagement with geoheritage and geodiversity by fostering dialogue between academic and community stakeholders in the Cliffs of Fundy UNESCO Global Geopark (<http://fundygeopark.ca>). Each seminar provided a concise overview of a focused topic, followed by structured discussion. Participants were invited to complete a pre-session survey to assess interests and experience. Attendance varied by session, ranging from 7 to 11 participants. Across the three seminars, participants considered definitions of geoheritage and geoconservation, the UNESCO Sustainable Development Goals, and shared perspectives on interdisciplinary and community-based collaboration. The final seminar explored the development of a potential geoheritage course syllabus intended for both university students and community members, reflecting sustained interest in formal educational offerings. Building on the GSI discussions, organizers developed a plan to celebrate international Geodiversity Day (October 6) within the Geopark. A full-day itinerary, spanning sunrise to sunset was implemented, beginning at Spencers Island, moving to the Fundy Discover Site, and concluding at Five Islands Provincial Park. Outcomes from this event were documented in the Geologists' Association Magazine and featured on the international Geodiversity Day website (<http://geodiversityday.org>). The 2025 GSI pilot was successful. The small cohort size supported meaningful discussion, relationship building and collaboration. Continued interest in developing a geoheritage course, together with the international visibility generated by the Geodiversity Day celebration, highlights the program's value to the Geopark and the region. Funding was provided by the Canadian Geological Foundation and the Atlantic Geoscience Society Products and Grants Committee.

---

**Resolving the effects of circulation structures on  
microbial diversity in the North Atlantic**

---

ALEXIS GAUVIN<sup>1</sup>, UNYIME UMOH<sup>2</sup>,  
AND G. TODD VENTURA<sup>2</sup>

*1. Department of Earth and Environmental Sciences,  
Dalhousie University, Halifax, Nova Scotia B3H 4R2, Canada  
<emigau5492@gmail.com>*

*2. Department of Earth Science, Saint Mary's University,  
Halifax, Nova Scotia B3H 3C3, Canada*

Compared to other areas of the ocean, the North Atlantic is rich in photosynthetic activity, responsible for roughly one-third of the world's net carbon dioxide uptake. The primary driver of this activity is thought to be a disruption of ocean stratification by turbulent interactions of strong warm- and cold-water currents, which are projected to weaken due to anthropogenic warming of seawater. How the microbial communities that sustain this productivity are responding to such a shift is thus far poorly understood. Last summer, the One Ocean Expedition's transatlantic cruise brought together scientists from around the world to measure ocean productivity markers and compare them to satellite data. From this cruise, our group collected a total of 26 surface water and water column filtrate samples from 10 stations along a transect between Iceland and Spain. We are currently using lipids—extracted from the filtrate samples using a modified Bligh-Dyer technique and measured using ultra high-pressure liquid chromatography coupled to a quadrupole time-of-flight mass spectrometer—as cellular proxies to resolve the identity and relative abundance of various microorganism types throughout the transect. Particular focus is placed on recording the occurrence of chlorophyll and its derivative pheophytin, indicators of photosynthetic communities, as well as the occurrence of archaeol and GDGT-class lipids, indicative of heterotrophic archaeal communities. This dataset, combined with other geochemical (nitrate, ammonia, TOC) and geophysical (remote sensing) datasets taken from the cruise, should provide a clear snapshot of how microbial communities are stratified along our transect, which can then be related to known current structures in the North Atlantic to resolve the impact of changing circulation patterns.

---

**Geochronology and petrogenesis of  
granitoid rocks and related critical mineral  
systems in New Brunswick, Canada**

---

AARON L. BUSTARD<sup>1</sup>, AMIRABBAS KARBALAEIRAMEZANALI<sup>1</sup>,  
DUSTIN R.L DAHN<sup>1</sup>, AND AYALEW GEBRU<sup>2</sup>

1. *New Brunswick Department of Natural Resources, South Tetagouche, New Brunswick E2A 7B8, Canada*
2. *New Brunswick Department of Natural Resources, Fredericton, New Brunswick E3B 5H1, Canada*  
<ayalew.gebru@gnb.ca>

New Brunswick is host to diverse critical mineral systems associated with felsic magmatism. Although the timing of several of these systems is well constrained, many of the critical mineral systems in New Brunswick need additional geochronological work. This two-year project, supported in part by a federal grant through the Critical Minerals Geoscience and Data (CMDG) Initiative, aims to address knowledge gaps in the timing of critical mineral systems associated with granitoid intrusions in New Brunswick that are favorable for hosting critical minerals, and to investigate their petrological origins. Four types of systems will be investigated: (1) Greisen/quartz stockwork quartz vein-related systems, including Burnthill (W-Mo-Sn), Flume Ridge (W), and Foster Lake (W); (2) Intrusion-related polymetallic vein systems including Pokiok (Ag-Cu-Zn-Pb-Au), Zealand Station (Be-Mo-W), Lake George (W-Mo), Popelogan-South Zone (Cu-Zn), Malachite (Cu-Co), Keymet (Cu-Pb-Ag-Zn), and Nigadoo (Cu-Pb-Ag-Zn); (3) LCT pegmatite systems, including in the Splan Pond (Li), Tower Hill (Li), and Kedron (Li) areas; and (4) W-Cu-Sn-Mo systems including Basswood Ridge (Sn-W-Li), Tower Hill (Sn-Mo-Cu-Zn-Au), Sorrel Ridge (Sn-Mo), Pleasant Ridge (Cu-Zn-Li-Mo-Pb), Kedron (W-Sn-Li-Be-Pb-Y-Zn-In-Cu-Au), Beech Hill (Mo-Sn-Zn), True Hill (Bi-Mo-Sn-W), Whittier Ridge; Wildcat Brook (Mo-W-Pb-Zn), Benjamin River South (Cu-Mo), and Nicholas Denys (Cu-Mo). The geochronological approach will begin with in situ dating of muscovite using the Rb–Sr method and will be followed up with the  $^{40}\text{Ar}/^{39}\text{Ar}$  method on selected samples. Zircon mineral separates will be analysed using laser ablation U–Pb geochronology. Zircon crystals yielding the youngest ages from laser ablation will subsequently be dated using high-precision Thermal ionization Mass Spectrometry techniques. Lu–Hf isotopic analysis will be conducted on the same zircon crystals used for geochronology. Re–Os dating of arsenopyrite and molybdenite will be completed for systems where other isotopic systems are not feasible. Additional work will include whole-rock litho-geochemistry and complementary isotopic work.

---

**Antifouling paint particles as a source of metal and microplastic pollution in harbour sediments: Chester Back Harbour, Nova Scotia, Canada**

---

ETHAN GERACI<sup>1</sup>, IAN SPOONER<sup>1</sup>, AND MARK MALLORY<sup>2</sup>

1. *Department of Earth and Environmental Science, Acadia University, Wolfville, Nova Scotia B4P2R6, Canada*  
<ethangeraci@acadiau.ca>
2. *Department of Biology, Acadia University, Wolfville, Nova Scotia B4P2R6, Canada*

In recent years, antifouling paints have been identified as a significant source of microplastic and heavy metal pollution in marine environments. Most modern antifouling paints contain copper (I) oxide as a primary biocide in addition to booster biocides such as zinc oxide, zinc pyrithione, and copper pyrithione. When released from a vessel, antifouling paint particles (APPs) will continue to leach biocides per their original design, posing an environmental risk to sediment-dwelling organisms. Our research focussed on determining the concentration of heavy metals associated with modern and historic marine paints (Cu, Zn, and Pb) and the concentration of APPs in the sediments of Nova Scotia's small harbours. Sediment gravity cores were collected in Autumn 2025 from Chester Back Harbour and two reference sites in Mahone Bay with limited boating traffic. Metal analysis was performed using an Epsilon 1 X-Ray Fluorescence Spectrometer. Results from the Chester Harbour sample show concentrations of Cu and Pb (mg kg<sup>-1</sup>) in the core's surface sediments that exceed the interim sediment quality guidelines (ISQGs) for Cu (18.7 mg kg<sup>-1</sup>) and Pb (30.2 mg kg<sup>-1</sup>) in marine sediments. In the case of Cu, concentrations in the top 2 cm of sediment exceeded the probable effect levels (PELs) for copper in marine sediments (108 mg kg<sup>-1</sup>). Results from both reference sites showed concentrations of Cu and Pb that did not exceed the ISQGs. Results for Zn from all three sites also did not exceed the ISQGs (<124 mg kg<sup>-1</sup>). Future work will include the isolation of suspected APPs along with the application of FTIR and Raman spectroscopy for polymer analysis.

---

**On the discovery of a large *Taenidium* burrow from the Clifton Formation in New Brunswick, Canada**

---

KAYLA GERARD<sup>1</sup>, MARARET SCOTT<sup>2</sup>, MATT STIMSON<sup>3</sup>,  
BAILEY MALAY<sup>3</sup>, STEVEN HINDS<sup>4</sup>, ADRIAN PARK<sup>4</sup>,  
AND LYNN DAFOE<sup>5</sup>

1. *Department of Earth Sciences, University of New Brunswick, Fredericton, New Brunswick E3B 3A3, Canada*  
<kayla.gerard@unb.ca>
2. *Department of Earth Sciences, Memorial University of Newfoundland, St. John's, Newfoundland and Labrador A1C 5S7, Canada*
3. *Natural History Department, New Brunswick Museum, Saint John, New Brunswick E2K 1E5, Canada*
4. *New Brunswick Department of Natural Resources, Fredericton, New Brunswick E3B 5H1, Canada*

5. Geological Survey of Canada, Natural Resources Canada,  
Dartmouth, Nova Scotia B2Y 4A2 Canada

The Moscovian Clifton Formation crops out in northern New Brunswick (NB) along the shores of the Bay du Chaleur. The formation is well known for its well-preserved plant fossils and lycopod trees preserved in life position. Trace fossils from the Clifton Formation have sparsely been reported in literature with only a cameo-appearance of an amphibian track and an invertebrate trackway reported amidst a detailed description of the sequence stratigraphy of the formation and sedimentary lithofacies. We report here the first systematic description of a large *Taenidium* burrow discovered near the township of Stonehaven NB. The burrow was preserved within a grey-green fine-grained sandstone from floodplain deposits. The fossil is relatively large (5 cm diameter and 50 cm in length) compared to other examples that have been reported from the Paleozoic of Atlantic Canada. However, large examples of *Taenidium* are known. *Taenidium* is characterized as an unbranched, unlined, subhorizontal burrow with evenly spaced meniscate backfill. A unique morphological feature includes the presence of 3–5 mm scale grey mud intraclasts within the meniscate backfill. This unique feature has not been recorded for *Taenidium* burrows previously, and may represent a new ichnospecies, and thus would expand the ichnogenus concept to include a heterogenous infill including intraformational mudclasts with a similar burrow infill as the parent rocks matrix. The *Taenidium* burrow presented here is interpreted to having been produced by a tetrapod, making it one of the rare examples of Carboniferous tetrapod burrow in the fossil record.

---

### Evidence for widespread barrier overstepping in coarse-clastic paraglacial coastlines: insights from Chedabucto Bay, Nova Scotia, Canada\*

---

CAMERON GREAVES<sup>1,2</sup>, JORDAN B.R. EAMER<sup>2</sup>, AND  
VITTORIO MASELLI<sup>1,3</sup>

1. Department of Earth and Environmental Sciences,  
Dalhousie University, Halifax, Nova Scotia B3H 4R2, Canada  
<camdgreaves@gmail.com>

2. Geological Survey of Canada, Natural Resources Canada,  
Dartmouth, Nova Scotia B2Y 4A2 Canada

3. Department of Chemical and Geological Sciences,  
University of Modena and Reggio Emilia,  
via Giuseppe Campi 103-41125, Modena, Italy

Key insights on how coastlines will evolve with future sea-level rise can be obtained by examining past coastal systems that formed during post-glacial relative sea-level (RSL) rise. In Atlantic Canada, coastlines are dominantly paraglacial,

meaning that their geomorphology and morphodynamics are largely governed by the presence of glacial deposits from successive glaciations. Recent detailed mapping of uniquely preserved drowned (overstepped) barrier beach deposits in Chedabucto Bay, Nova Scotia has revealed not only the ubiquity of these features but also that barrier overstep may play a more dominant role in shoreline retreat than once thought. Many studies investigating barrier retreat through overstepping primarily focus on sandy systems while coarse-clastic systems remain understudied. Here, we investigate coarse-clastic paraglacial barrier systems by comparing the geophysical characteristics of modern barrier features surveyed through ground penetrating radar and LiDAR, with those now drowned offshore, imaged by multibeam bathymetry and seismic reflection data. We aim to investigate the environmental conditions that have led to their formation and preservation and to understand the differences observed in modern systems. Our mapping in Chedabucto Bay has revealed 62 individual relict paraglacial barriers, with glacio-isostatic RSL models suggesting systems formed between ca. 9 500 and ca. 5 500 yrs BP in a similar coastal configuration to that observed today, with recent radiocarbon ages supporting this regional model. From this catalogue of barrier features there is a greater prevalence of drift-aligned relict barriers, which we attribute to a sedimentological preservation bias from longshore transport dynamics along northern coast of the bay. From these findings we suggest that barrier overstepping plays an extremely important role in coastal dynamics for coarse-clastic paraglacial systems. This mechanism of coastal retreat must be considered by researchers who attempt to model long-term sediment budgets and shoreline retreat for these systems, especially during periods of high rates of RSL rise.

\*Winner: AGS Sandra Barr Award for best graduate student oral presentation

---

### Surficial geology and permafrost conditions within the involuted terrain near Tuktoyaktuk, Northwest Territories, Canada

---

BAILEY R. GRONDIN<sup>1</sup>, ALEXANDRE CHIASSON<sup>1,2</sup>,  
ALEJANDRO ALVAREZ<sup>1</sup>, STEVEN V. KOKELJ<sup>2</sup>,  
AND DUANE G. FROESE<sup>1</sup>

1. Department of Earth and Atmospheric Sciences,  
University of Alberta, Edmonton, Alberta T6G 2E3, Canada  
<bgrondin@ualberta.ca>

2. Northwest Territories Geological Survey,  
Government of the Northwest Territories, Yellowknife,  
Northwest Territories X1A 2L9, Canada

The Arctic coastal community of Tuktoyaktuk, Northwest Territories, faces severe coastal erosion and permafrost thaw hazards driven by its location on ice-rich terrain. Near Tukto-

yaktuk, Northwest Territories, faces severe coastal erosion and permafrost thaw hazards driven by its location on ice-rich terrain. Near Tuktoyaktuk, the involuted terrain is characterized by clusters of ice-cored hills (involute hills) with ridge forms along their tops separated by thermokarst basins and lakes. The origin of the involuted terrain is not well-understood, but a general model suggests that these ice-rich deposits formed in front of the retreating Laurentide ice Sheet along the Tuktoyaktuk coastlands through a combination of segregation and intrusive ice growth during the late Pleistocene. The last regional surficial geology mapping (1:500 000) was completed more than 30 years ago, prior to the availability of LiDAR, high resolution satellite imagery, modern glaciogenic concepts, and before the current rapid climate-driven landscape change began. Since ground ice content is the primary driver of permafrost thaw sensitivity, mapping the distribution, and understanding the development and degradation of involuted hills is crucial. In 2025, we undertook initial fieldwork, and developed the first high-resolution surficial geology and geohazard assessment of the involuted terrain, including: (i) producing 1:10 000 and 1:50 000 scale surficial geology and permafrost feature maps (1:5000); (ii) establishing relations between geology and ground ice by correlating electrical resistivity tomography surveys, headwall mapping of thaw slumps, and shallow drilling with ground ice observations; (iii) examining areas of recent landscape change by analyzing multi-temporal LiDAR (2011–2024) to quantify recent thermokarst features, and classify geohazards. The ultimate goal of this work is to establish surficial geology-ground ice relations to improve our understanding of ongoing climate-driven landscape change and provide baseline information for infrastructure planning in the region.

---

**Mafic-ultramafic melt inclusions in early magmatic apatite record low degrees of partial melting of a critical metal-enriched, suspected Neoproterozoic basement, Mersey Point, Nova Scotia, Canada**

---

JACOB HANLEY<sup>1</sup>, GEENA MORSE<sup>1</sup>, ANNA TEREKHOVA<sup>1</sup>, AND BRENDAN MURPHY<sup>2</sup>

1. *Department of Earth Science, Saint Mary's University, Halifax, Nova Scotia B3H 3C3, Canada*  
<jacob.hanley@smu.ca>

2. *Department of Earth and Environmental Sciences, St. Francis Xavier University, Antigonish, Nova Scotia B2G 2W5, Canada*

The Mersey Point gabbronorite intrusion is one of several very small Late Devonian mafic intrusions coeval with granitoids that host intrusion-related Sn-W-Mo deposits

in Nova Scotia. The gabbronorite is an olivine (>Fo<sub>80</sub>)-plagioclase mesocumulate with poikilitic orthopyroxene, clinopyroxene, and phlogopite. Apatite, spinel, and sulfides (Fe-Ni-Co-Cu-S) are present as accessory phases. Early apatite phenocrysts contain primary, high alkaline silicate melt inclusions (SMI) of picrobasaltic/basanitic (in primocrysts) to trachytic composition. Apatite trace element compositions (by LA-ICP-MS) indicate growth from plagioclase-saturated mafic-ultramafic magma and show strong crustal contamination. Compatible major/trace elements in the SMI show behaviors consistent with differentiation. However, incompatible LILE/HFSE show no enrichment with differentiation and, instead, reach extremely high concentrations in the most primitive melt inclusions. Notably, concentrations of critical metals W and Mo are 1–3 orders higher than in MORB and arc basalts; Sn concentrations are unremarkable. The compositional characteristics of the apatite and associated SMI provide the first direct evidence for the onset of partial melting of protolith lower crustal rocks by mafic end-member magmas within the Meguma lithotectonic Zone. Strong enrichment in W relative to Sn indicate very low degrees of partial melting of a phyllosilicate-rich crustal protolith, suggesting that in areas where larger volumes of such mafic magmas interacted with lower crust, higher degrees of partial melting may have produced Sn- and W-rich melts, a likely prerequisite for intrusion-related deposits in coeval granitoid rocks.

---

**Spatial distributions of surface textures, sedimentary structures, and neoichnological traces in the Bay of Fundy, Canada**

---

BRETTE HARRIS<sup>1</sup>, KELLY ROZANITIS<sup>2</sup>, JEFFREY KAVANAUGH<sup>2</sup>, AND MURRAY GINGRAS<sup>2</sup>

1. *Department of Earth Sciences, Memorial University of Newfoundland, St. John's, Newfoundland and Labrador A1B 3X5, Canada* <bretteh@mun.ca>

2. *Department of Earth and Atmospheric Sciences, University of Alberta, Edmonton, Alberta T6G 2E3, Canada*

Intertidal environments are shaped by complex interactions between physicochemical processes and biological activity. These interactions influence sedimentary structures and surface textures crucial for interpreting modern and ancient siliciclastic tidal systems. This study integrates high-resolution unmanned aerial vehicle (UAV) imagery, sedimentological analyses, and neoichnological observations to investigate the spatial distribution of surface features and modern organism traces in macrotidal intertidal flats. This project focuses on the mud-dominated inner estuary and fluvial-tidal transition zones within the Salmon and

Petitcodiac River estuaries (Bay of Fundy, Canada). Surface features were mapped and classified, quantifying the affinity of trace types to specific sedimentary surfaces using spatial statistics, specifically affinity factor analysis and Markov chain analyses with Monte Carlo simulation. Grain size distributions from sediment cores complement the spatial dataset. Results reveal distinct spatial relationships between sedimentary facies and biogenic structures, establishing quantitative baselines for sediment-biota relationships in macrotidal estuaries. We demonstrate that certain organism trace-making activity exhibits strong environmental control, clustering predictably on specific depositional surfaces. This integrated UAV-based neoichnological approach provides a novel framework for interpreting trace fossil assemblages in analogous ancient siliciclastic deposits and advances methodologies for high-resolution sedimentary investigations in coastal environments. The quantitative baselines established here offer a valuable reference for paleoenvironmental reconstruction and prediction of organism-sediment interactions across tidal systems.

---

**Preliminary petrographic and micro XRF insights into the origin and diagenetic evolution of siderite in the Ben Nevis Formation, Jeanne d'Arc Basin, offshore Newfoundland, Canada**

---

BRETTE HARRIS<sup>1</sup>, ERIC HUTTON<sup>1</sup>, NOAH SLANEY<sup>1</sup>, JAIMEE HODGSON<sup>1</sup>, SIMONE BOOKER<sup>2</sup>, ERIN ADLAKHA<sup>2</sup>, DAVID LOWE<sup>1</sup>, AND HILARY CORLETT<sup>1</sup>

1. *Department of Earth Sciences, Memorial University of Newfoundland, St. John's, Newfoundland and Labrador A1B 3X5, Canada <bretteh@mun.ca>*

2. *Department of Earth Science, Saint Mary's University, Halifax, Nova Scotia B3H 3C3, Canada*

Early diagenetic and burial siderite can provide new constraints on the paragenesis of cementation in marginal to shallow-marine reservoir successions. Building on previous research that uses siderite chemistry and morphology to track basin fluid evolution in Mesozoic clastic systems of the Scotian Basin, this study focuses on the Aptian–Albian Ben Nevis Formation of the Jeanne d'Arc Basin, offshore Newfoundland. Here we use hyperspectral and petrographic analysis, scanning electron microscopy (SEM), and micro-X-ray fluorescence ( $\mu$ XRF) elemental mapping to characterize siderite paragenesis and textural relationships to host sediments. Preliminary observations document multiple siderite habits, including nodules and pore-filling to grain-binding cements, which vary with diagenetic facies and burial depth. Variations in Fe, Mg, Mn, and Ca, suggest progressive compositional evolution linked to changes in diagenetic fluid composition and redox conditions. Textural criteria, including laminae deflection around siderite-cemented

domains, contrasts between sutured and open grain contacts, and cross-cutting relationships between diagenetic phases provisionally distinguish eodiagenetic concretions and cements from mesodiagenetic replacement and pore-filling phases. These preliminary results support a diagenetic model in which multiple generations of siderite cements and nodules occur with distinct chemical signatures and textures. Ongoing work integrating systematic petrography, SEM-EDS, quantitative geochemistry, and high-resolution  $\mu$ XRF guided by hyperspectral analysis of core will refine the paragenetic framework and assess how siderite morphology in the Ben Nevis Formation can be used as a proxy for fluid pathways, redox evolution, and reservoir quality in this key Atlantic Canadian petroleum system.

---

**Characterizing the geomorphology and dynamics of potential rock glaciers in parc national de la Gaspésie using LiDAR, thermal imaging, and GPR**

---

SHANNON HEGGER, KYNAN HUGHSON,  
AND GILLES NGORAN

*Department of Earth Sciences, University of New Brunswick, Fredericton, New Brunswick E3B 3A3, Canada  
<shannon.hegger@unb.ca>*

Three potential rock glaciers were visited in Parc National de la Gaspésie in summer of 2025, with the goal of contributing to the understanding of permafrost and periglacial processes in Eastern Canada. The presence of rock glaciers, along side their structures and activity, in eastern Canada has been fairly under documented and under characterized in recent decades. With the recent rise of global temperatures, it is more important than ever to study these features and document their climate sensitivity. Remote sensing surveys were conducted with a small remotely piloted aircraft system to collect photogrammetry, LiDAR, and thermal infrared data to produce high-resolution terrain models and multispectral orthophotos. These data were used in the following ways: high resolution imaging (orthophotos), mapping surface temperature anomalies (thermal imaging), identifying and mapping geomorphic facies (LiDAR), and examining internal reflectors, such as ice- and water-rich zones (GPR). Analysis conducted thus far includes mapping of geomorphological features based on orthophotos and LiDAR data and inferring relative chronostratigraphy. Additional analysis using thermal images have identified zones of surface temperature variation associated with geomorphological structures. Radargrams are currently undergoing examination for internal reflectors in order to map locations of potential ice- and water-rich zones to later cross-reference with temperature anomalies and geomorphological features.

---

**Revised structural and stratigraphic correlation of the Boyd Creek tuff of the eastern Moncton Sub-basin, southeastern New Brunswick**

---

STEVEN HINDS<sup>1</sup>, ADRIAN PARK<sup>1</sup>, MATT STIMSON<sup>2</sup>,  
ANDREW MACRAE<sup>3</sup>, AND OLIVIA KING<sup>4</sup>

1. *New Brunswick Department of Natural Resources, Fredericton, New Brunswick E3B 5H1, Canada*  
<Steven.Hinds@gnb.ca>
2. *Natural History Department, New Brunswick Museum, SaintJohn, New Brunswick E2K 1E5, Canada*
3. *Department of Earth Science, Saint Mary's University, Halifax, Nova Scotia B3H 3C3, Canada*
4. *Department of Earth Sciences, Carleton University, Ottawa, Ontario K1S 5B6, Canada*

The Boyd Creek tuff was recently radiometrically dated by U–Pb Laser Ablation method to be  $337.4 \pm 1.5$  Ma (Visean–Holkerian). This date is significant as there are very few units which provide a radiometric age to constrain the timing of stratigraphic deposition and tectonic events throughout the Devonian–Carboniferous sedimentary sequences of New Brunswick. Historically, age determinations for the Devonian–Carboniferous sediments are reliant on the study of micro and macro paleontology of spores and plants. The Boyd Creek tuff was logged in some early 1900s boreholes and very sporadic outcrops within the Stoney Creek to Gautreau area. During the 1940s exploration phase by Shell Oil, this tuff was interpreted to have been deposited within the Hillsborough Formation before being re-assigned to the Late Tournaisian during the early 2000s. Hence, the scarcity of outcrops of the Boyd Creek tuff and the uncertain age of the stratigraphy has complicated surface to subsurface geological correlations within the Berryton to Gautreau Village area of southeastern New Brunswick. The new age is consistent with a recent biostratigraphic study in the Gautreau–Weldon area and the Boyd Creek tuff is correlative with other Visean volcanic episodes across the southern portion of the onshore Maritimes Basin in New Brunswick. Very recently, a new outcrop of the Boyd Creek tuff was discovered along the Hiram Brook within the Stoney Creek Station area. This outcrop, along with new fieldwork, borehole, and seismic interpretations in adjacent areas reveals the Boyd Creek tuff is part of a doubly plunging syncline extending from the Berryton to Gautreau Village areas. The southern portion of this syncline near Baltimore is truncated by the Clover Hill Fault, which explains the absence of the Boyd Creek tuff in nearby boreholes.

---

**The East Kemptville Sn(-Ag-Cu-Zn) domain, southwestern Nova Scotia, Canada: a potential world class Sn-In-Li-Rb district**

---

RICK HORNE AND DANIEL KONTAK

1. *Everest Metals, 72 Johnstone Avenue, Dartmouth Nova Scotia B2Y 2K5, Canada* <rickhorne57@gmail.com>
2. *Harquail School of Earth Sciences, Laurentian University, Sudbury, Ontario P3E 2C6, Canada*

The ongoing quest for critical metals to support the green transition is based on a long and expanding list of metals that includes Sn. The demand for domestic sources of critical metals is important and exemplified by Sn for which there is no current primary production in North America. The former producing East Kemptville Sn mine and surrounding area define a northeast-trending “Sn domain” of 100 km that extends from the Dominique prospect south of Yarmouth eastwards to the Caledonia occurrence and thus represents Nova Scotia’s premiere critical metal target. This domain hosts numerous occurrences of Sn and related metals (W, Zn, Cu, In, Ag) within both the parental granite and host Meguma Supergroup metasedimentary rocks. Tin occurrences include all the classic Sn deposit types, including quartz vein-greisen zones within host granite and quartz veins and skarn in host metasediments. Although the East Kemptville Sn mine is the only currently defined economic resource, several significant occurrences in the area suggest further resources are likely to be defined through systematic exploration because all well-defined Sn domains globally host several or more deposits (e.g., Cornwall, Bolivia, Australia). In addition to Sn, mineralization includes potential Cu-Zn-in-Rb-Li resources as byproducts, with Cu (chalcopyrite) and Zn (sphalerite) both byproducts of historical mining, in addition to Ag and W, and with In concentrated in sphalerite having the potential for recovery at the smelting stage. Li and Rb are enriched in micas, including zinnwaldite, trillithionite, muscovite and biotite, but are decoupled from Sn on a local scale and thus may require selective mining and processing streams; however, as they occur in the same general area they thus benefit from common infrastructure.

---

**The diverse morphology and internal structure of North American pingos reveals complex ice growth and loss**

---

KYNAN HUGHSON

*Department of Earth Sciences, University of New Brunswick, Fredericton, New Brunswick E3B 3A3, Canada*  
<kynan.hughson@unb.ca>

Pingos are distinct ice-rich hills found in permafrost regions that serve as critical indicators of ground ice conditions and possess morphological analogs on planetary bodies like Mars and Ceres. Despite their cultural and scientific significance, the relationship between pingo sur-

face morphology and internal cryostratigraphy remains poorly constrained, often relying on classical models of monolithic ice cores. This study provides novel geophysical constraints on pingo internal structure by analyzing four pingos near Prudhoe Bay, Alaska, and nine within Pingo Canadian Landmark, near the Hamlet of Tuktoyaktuk in the Northwest Territories. We integrated high-resolution digital elevation models with ground penetrating radar and capacitively-coupled electrical resistivity surveys to resolve subsurface architecture and ground ice distribution. Our results challenge the prevalence of the idealized hydrostatic pingo model. Of the thirteen investigated sites only two, including the massive ibyuk pingo, displayed geophysical characteristics supporting a large, monolithic massive ice core. Instead, we observed significant structural diversity; many morphologically “mature” pingos lacked discrete central cores, exhibiting instead scattered lenticular ice bodies and complex layering. Observations of “young” pingos indicate that growth likely initiates through ice segregation, transitioning to a mixed-mode of segregation and intrusive processes as the pingo matures and the overburden undergoes brittle failure. We identify a positive correlation between morphological maturity and internal cryostratigraphic complexity, yet find that topographic prominence alone is a poor predictor of subsurface heterogeneity. Furthermore, structural patterns such as slope-parallel resistive anomalies support a thin elastic plate model for pingo formation, where intrusive ice is emplaced along flank failure surfaces rather than exclusively via basal accretion. These findings suggest that monolithic ice cores represent a rare endmember, and that a distributed segregation-injection framework better explains pingo evolution on Earth and analogous planetary environments.

---

**A >360 myr history of polymetallic vein formation at Cobalt–Gowganda, Ontario, and relevance to Nova Scotia, Canada**

---

DANIEL J. KONTAK<sup>1</sup>, JACOB J. HANLEY<sup>2</sup>, JOSHUA JACKMAN<sup>2</sup>, ROBERT A. CREASER<sup>3</sup>, AND JONATHON TOMA<sup>4</sup>

1. Harquail School of Earth Sciences, Laurentian University, Sudbury, Ontario P3E 2C6, Canada
2. Department of Earth Science, Saint Mary's University, Halifax, Nova Scotia B3H 3C3, Canada  
<joshua.jackman@smu.ca>
3. Department of Earth and Atmospheric Sciences, University of Alberta, Edmonton, Alberta T6G 2E3, Canada
4. Department of Earth and Planetary Sciences, Yale University, New Haven, Connecticut 06520-8109, USA

The world-class “five-element” vein systems of Cobalt–Gowganda (Ontario) are among the highest-grade deposits

of Ag and Co (15.5 billion g Ag, 20.5 million kg Co) globally. Their proximity to a regional mafic sill complex (Nipissing diabase) spurred the assumption that the vein mineralization was coincident with and genetically related to the intrusive event at ca. 2.22 Ga. Results from Re–Os analysis of pre- and syn-ore-stage sulfarsenide phases from several polymetallic vein settings indicate a greatly protracted history for mineralization spanning >360 myr which precedes the Nipissing diabase by >245 Ma. These results require alternate model(s) that invoke both an extended time frame and various events to explain the polymetallic mineralization. These findings are perhaps relevant to polymetallic vein systems in Nova Scotia (e.g., Nictaux Falls Dam, Lansdowne, Cape St. Marys) as they share some similarities to those of Cobalt–Gowganda in metallogeny (e.g., Co, Ni, Ag, Sb, Pb), and peculiar mineralogy (e.g., Sb-, Ag-sulfosalts, native Ag, Sb, cobaltite). The latter geological best assumption for deposit genesis at Cobalt–Gowganda informed exploration, with production ceasing when deposits could no longer be found near the favoured setting of diabase and country rock contact. Whereas the documentation of this protracted history of mineralization does not preclude a role for the diabase intrusion for some deposits, it suggests that mineralization also occurred before and after, due to other processes which may also include other undiscovered locations. Similarly, the polymetallic vein deposits of Nova Scotia are considered in isolation, and with emphasis on their host formations. Instead, these deposits may represent an extended, more complicated system of events.

---

**What's REALLY under the Meguma terrane?  
A fresh look at xenoliths from the Eastern Shore dykes**

---

REBECCA JAMIESON<sup>1</sup>, SANDRA M. BARR<sup>2</sup>, CHRIS E. WHITE<sup>2</sup>, DEANNE VAN ROOYEN<sup>2</sup>, AND DONNELLY ARCHIBALD<sup>3</sup>

1. Department of Earth and Environmental Sciences, Dalhousie University, Halifax, Nova Scotia B3H 4R2, Canada  
<beckyj@dal.ca>
2. Department of Earth and Environmental Science, Acadia University, Wolfville, Nova Scotia B4P2R6, Canada
3. Department of Earth and Environmental Sciences, St. Francis Xavier University, Antigonish, Nova Scotia B2G 2W5, Canada

Xenoliths hosted by the late Devonian Eastern Shore lamprophyre dyke swarm are the only samples of what lay beneath the Meguma terrane when the dykes were intruded (ca. 370 Ma). The xenoliths represent a wide variety of lithologies and mineral assemblages and have been the focus of several studies since their discovery 40 years ago. Nevertheless, some long-standing interpretations deserve further scrutiny, including suggestions that: (i) pelitic lithologies are not potential source rocks for peraluminous

magmatism; (ii) mafic granulites are an important component of the xenolith suite; and (iii) their geochemical and isotopic characteristics indicate that the Meguma terrane is underlain by Avalonian lower crust. We report preliminary results from a re-examination of these important samples undertaken to document their characteristics further and to test existing hypotheses concerning their origin and significance. Most xenolith samples are pelites or pelitic migmatites with complex polymetamorphic histories. An early kyanite + garnet + biotite-bearing assemblage is strongly overprinted by sillimanite associated with corundum + spinel and sapphirine-bearing leucosome. The observations are consistent with high-temperature biotite dehydration melting; preserved melt pockets are clearly peraluminous and quartz-poor. Later retrogression produced distinctive spinel + gedrite coronas around garnet. Mafic xenoliths display pristine igneous textures involving pyroxene + plagioclase ± amphibole but lacking either garnet or olivine. They are interpreted as gabbro and diorite; samples resembling metamorphic mafic granulite have not yet been recognised. Based on these observations we question some previous interpretations regarding the nature and significance of the pelitic and mafic xenoliths. Work is in progress to determine the petrogenetic affinity of the mafic igneous rocks and P-T conditions for melting of the pelitic rocks.

---

### Gold enrichment in porphyry Cu-Au systems: insights from low-melting chalcophile elements behavior in the Late Triassic Zhongdian arc, Sanjiang region, SW China

---

JIAWEN JIANG<sup>1,2</sup>, HAIJUN YU<sup>3</sup>, DAVID LENTZ<sup>2</sup>,  
AND WEN-CHANG LI<sup>4</sup>

1. School of Earth Resources, China University of Geosciences, Wuhan 430074, China <j7c3j@unb.ca>
2. Department of Earth Sciences, University of New Brunswick, Fredericton, New Brunswick E3B 5A3, Canada
3. Key Laboratory of Sanjiang Metallogeny and Resources Exploration and Utilization, Yunnan Geological Survey, China
4. Kunming University of Science and Technology, Kunming 650093, China

Gold is efficiently concentrated in subduction-related porphyry copper-deposits (PCDs), which are estimated to host nearly 20% of the world's gold resources. Although subduction and its duration are fundamental drivers of magma fertility in PCDs, the specific mechanisms responsible for Au enrichment—particularly the role of low-melting chalcophile elements (LMCEs) and nanoscale formation pathways—remain poorly understood. This study focuses on the giant Late Triassic Pulang and Lannitang porphyry Cu-Au deposits, located in the Zhongdian terrane,

Sanjiang region, SW China, which are interpreted to have formed in an island-arc-related subduction setting, but preserve contrasting structural levels and thermal histories. Through a comparative multi-analytical approach—integrating reflected light microscopy, scanning electron microscopy (SEM), electron probe microanalysis (EPMA), laser ablation-inductively coupled plasma time-of-flight mass spectrometry (LA-ICP-TOF-MS), and high-resolution transmission electron microscopy (HRTEM)—we demonstrate that Au enrichment at Pulang was efficiently scavenged during the early magmatic-hydrothermal stage associated with Bi-Te-As mineralization, when its LMCE phases formed under high-temperature and oxidized conditions and promoted nanoscale gold nucleation. These gold nanoparticles were subsequently preserved within arsenopyrite and rooseveltite during progressive cooling and oxidation. In contrast, the structurally shallower Lannitang system, characterized by a thicker wall-rock, records extensive late-stage hydrothermal overprinting, in which earlier LMCE-mediated Au reservoirs were largely re-equilibrated or destroyed, resulting in dominantly free native gold precipitation during argillic alteration. These contrasting features reflect differences in structural level and thermal state rather than distinct metallogenic processes. We therefore propose that subduction-related settings create the essential thermo-chemical window—marked by high oxidation states, delayed sulfide saturation, and elevated temperatures—that promotes the enrichment and saturation of LMCEs. These LMCEs act as highly efficient gold scavengers, representing a primary control on gold endowment in PCDs. This LMCE-driven mechanism offers a new mechanistic and nanoscale framework for exploration targeting.

---

### A provisional stratigraphic revision of the Neoproterozoic–Cambrian succession in the Caledonian Highlands, southeastern New Brunswick, Canada

---

SUSAN JOHNSON<sup>1</sup>, AYALEW GEBRU<sup>2</sup>, ADRIAN PARK<sup>2</sup>, AND  
STEVEN ROSSITER<sup>1</sup>

1. New Brunswick Department of Natural Resources, Sussex, New Brunswick E4E 5L2, Canada <Susan.Johnson@gnb.ca>
2. New Brunswick Department of Natural Resources, Fredericton, New Brunswick E3B 5H1, Canada

The Caledonian Highlands encompass an area 30 km wide by 130 km long bordering the northern Bay of Fundy from Saint John to Alma. Neoproterozoic and Cambrian rocks of the Caledonian Highlands represent one of several Avalonian terranes in the northeastern Appalachian orogen, along with northern Nova Scotia and southern Cape Breton Island, southeastern Newfoundland, and southeastern Newfoundland,

and southeastern New England, USA. Since the last bedrock mapping was conducted in the late 1980s–early 1990s a wealth of new data has been published; however, no attempt has been made to reconcile these data with the published set of bedrock geology maps. Towards this aim, a multi-year bedrock mapping project was initiated by the New Brunswick Geological Surveys Branch in 2015. The Neoproterozoic succession in the highlands was previously divided into two groups of Ediacaran age: an older Broad River Group (ca. 620–615 Ma) and younger Coldbrook Group (ca. 550 Ma). The more recent recognition of a suite of ca. 690 Ma felsic intrusions that were emplaced into metavolcanic and volcanoclastic rocks assigned to the Broad River Group has led to significant stratigraphic revisions and the delineation of a new Cryogenian succession we have named Lumsden Group. This new group hosts the Lumsden base-metal deposit, which is similar in age and metallogeny to the Stirling deposit on Cape Breton Island (Mira terrane). Some other major revisions include reducing the rank of several stratigraphic formations in the Coldbrook Group to members, thereby decreasing the number of formations in the group from twelve to five. In the structurally complex eastern highlands, we have expanded the distribution of the Coldbrook Group and re-assigned the clastic meta-sedimentary rocks of the Goose River Formation (formerly Broad River Group) to the Cambrian Saint John Group, indicating that the Avalonian overstep sequence is more extensive in the eastern highlands than previously thought.

---

### **Preliminary investigation of the metamorphic gradient in the southwestern Meguma terrane, Nova Scotia, Canada**

---

JILLIAN KENDRICK AND CHARBEL ABOU-KHALIL

*Department of Earth Science, Saint Mary's University, Halifax, Nova Scotia B3H 3C3, Canada <jillian.kendrick@smu.ca>*

The Meguma terrane in southwestern Nova Scotia exhibits a low-pressure metamorphic gradient, locally reaching suprasolidus conditions near the Devonian Barrington Passage Pluton. Excellent exposure, particularly along the coast, allows metasedimentary units of the Goldenville and Halifax groups to be traced across metamorphic isograds. This region therefore provides an opportunity to study fundamental aspects of the metamorphic evolution of metasedimentary sequences, including (1) the influence of variable bulk compositions on pelitic phase equilibria, and (2) the mobilization of trace metals through metamorphic reactions. Here, we present preliminary field and petrographic observations from coastal exposures near Villagedale and Sandy Point in Shelburne County. Migmatite outcrops near Villagedale

are foliated with patchy leucosome and abundant muscovite porphyroblasts, which appear to be retrograde after the peak assemblage of sillimanite + biotite + feldspar + quartz + magnetite ± garnet. Rare tourmaline is also texturally late. Near Sandy Point, the influence of changing pressure-temperature (P-T) conditions and bulk composition is apparent. Metamorphic grade decreases southeast along the coast, indicated by staurolite disappearing from the assemblage and the garnet distribution becoming sporadic in pelitic layers. However, this sequence is interrupted by a package of green metasiltsstones containing biotite and chlorite. Future work will focus on high-resolution mapping of shoreline exposures in this region, with documentation of variations in bulk composition. Geochronology of multiple minerals in the migmatite (e.g., zircon, monazite) will be used to constrain the timing of partial melting and melt crystallization, potentially providing a clearer link between magmatism and metamorphism in the region.

---

### **Deformation and metamorphism in the southernmost Chebogue Point shear zone of the Meguma terrane, Nova Scotia, Canada**

---

ROWAN KENNEDY, DEANNE VAN ROOYEN,  
AND CHRIS E. WHITE

*Department of Earth and Environmental Science,  
Acadia University, Wolfville, Nova Scotia B4P2R6, Canada  
<162276k@acadiau.ca>*

The Chebogue Point shear zone (CPSZ) is a major intra-terrane high-strain zone in the Meguma terrane in southwestern Nova Scotia. It juxtaposes Cambrian to Ordovician rocks of the metasedimentary Goldenville Group (Green Harbour, Government Point and Moshers Island formations) and Halifax Group (Cunard Formation) with Silurian and younger metavolcanic and metasedimentary rocks of the Rockville Notch Group (White Rock Formation and associated suites of mafic sills and dykes) that only occur northwest of the shear zone. At a coastal section at Chebogue Point the stratigraphy is well-exposed and units young towards the NW. At the base of the section the Green Harbour Formation consists of massive metasandstone and grades upwards to the Government Point Formation which consists of cross-bedded siltstone with minor sandstone layers. The overlying high-Mn Moshers Island Formation consists of siltstone with spessartine-rich nodules and lenses. The Cunard Formation consists of black, sulphide-rich chloritic slate with rare sandstone layers. The White Rock Formation in this section is represented by interlayered siltstone, tuff, and amphibolite. The section shows a pronounced deformation gradient increasing in intensity towards the NW. From the base of the section to the top,

$S_0$  and  $S_1/D_1$  spaced cleavage defined by weakly foliated muscovite and chlorite become increasingly overprinted by  $S_2/D_2$  foliation that folds and rotates the  $S_0/S_1$  structures into a composite transposition foliation and vertical and rootless isoclinal folds defining the CPSZ. Based on the presence of chlorite and muscovite in the  $S_2$  structures (foliation, pressure shadows around diagenetic pyrite, axial planar cleavage in folds, and mineral lineations) the  $S_2/D_2$  shearing took place at lower greenschist facies conditions, in contrast to amphibolite-facies shearing recorded further inland along strike. Kinematic indicators show local SE-directed motion overprinted by the main NW-directed oblique sinistral motion that defines the CPSZ.

---

### Rare earth and Nb enrichment in Kimberlite perovskite: Insights from the Lac de Gras kimberlites, Canada

---

SAHROZ KHAN AND YANA FEDORTCHOUK

*Department of Earth and Environmental Sciences,  
Dalhousie University, Halifax, Nova Scotia B3H 4R2, Canada  
<sahroz.khan21@gmail.com>*

Kimberlites are mantle-derived igneous rocks known for their diamond content, but they also host significant rare earth elements (REEs) and critical metals (e.g., Ni, Co, and Nb) in primary and secondary REE mineral hosts. Perovskite and apatite are the primary REE-bearing minerals; however, perovskite is generally the dominant REE host, with crystallisation sequence exerting a strong control on REE distribution. This study presents an investigation of different lithological facies from six Lac de Gras (LDG) kimberlites, with a focus on perovskite and compares it with kimberlite data from other global localities. The samples were examined using petrographic, scanning electron microscopes (SEM), and electron microprobe (EMP) to characterise mineral textures and compositions. In hypabyssal kimberlites (Grizzly and Leslie), perovskite occurs as euhedral to subhedral crystals with sharp grain boundaries and compositional zoning. In contrast, perovskite in the volcanoclastic kimberlites (Beartooth, Koala, Misery, and Panda) displays irregular, embayed, and fragmented morphologies, consistent with resorption and fragmentation. The highest REE and Nb concentrations are found in the few perovskites from the Leslie, wherein abundant groundmass apatite often contains Ce-La and REE-rich apatite inclusions. In Grizzly, the abundant perovskite is the primary REE host and exhibits Nb-rich rims of up to 5  $\mu\text{m}$  thickness. Perovskite from Panda and Koala shows REE and Nb concentrations comparable to the core of perovskite from Grizzly. Perovskite in samples from Misery and Beartooth has lower REE and Nb concentrations. In Beartooth, REE-rich apatite occurs in association with groundmass carbonate, and monazite occurs as overgrowth on

ilmenite. REE minerals, including monazite and ancylite, occur sporadically in both hypabyssal and volcanoclastic kimberlites. Compared with other African and Canadian kimberlites, LDG perovskite has a higher REE concentration and shows Nb enrichment. The variation in the perovskite composition between the LDG kimberlites and worldwide localities reflects variation in melt composition and emplacement mechanism.

---

### Critical mineral potential of historical antimony tailings in Atlantic Canada: a case study from the Lake George mine, New Brunswick, Canada

---

NIYAYESH KHORSHIDI<sup>1</sup>, MICHAEL B. PARSONS<sup>2</sup>,  
AND DAVID R. LENTZ<sup>1</sup>

- 1. Department of Earth Sciences, University of New Brunswick, Fredericton, New Brunswick E3B 5A3, Canada  
<niya.khorshidi@unb.ca>*
- 2. Geological Survey of Canada, Natural Resources Canada, Dartmouth, Nova Scotia B3B 1A6, Canada*

Between 1998 and 2023, Canada ranked as the ninth-largest producer of primary antimony (Sb); however, Canadian Sb deposits now contribute minimally to global supply. Atlantic Canada hosts three Sb deposits (Clarence Stream and Jones Brook, NB, and Beaver Brook, Central Zone, NL), six past producers (including the Beaver Brook (NL), Lake George (NB), and West Gore (NS) mines), and 35 significant Sb occurrences, indicating strong potential for future exploration. Historical mine tailings across the region represent an underexplored secondary source of critical minerals and a potential resource for both recovery and reclamation at former mine sites. This study examines the distribution of Sb deposits, occurrences, and Sb-bearing tailings in Atlantic Canada, with particular attention to abandoned Sb mines that may retain economically relevant concentrations of Sb and associated critical minerals. Among these, the Lake George Sb mine in New Brunswick—formerly the largest Sb producer in North America—stands out due to its notable content of Sb and gold (Au). Mineralogical analyses of the Lake George tailings identified primary Sb minerals such as stibnite [ $\text{Sb}_2\text{S}_3$ ] and berthierite [ $\text{FeSb}_2\text{S}_4$ ], along with secondary weathering products including Sb oxides and Sb-Fe oxides. Although Au occurs in microscopic or invisible forms, bulk geochemical analyses reveal elevated Au concentrations in the tailings (mean 415 ppb) and substantial residual Sb (mean 3,015 ppm), indicating incomplete historical recovery and highlighting the potential for modern reprocessing. The results of this study show that a detailed understanding of the mineral hosts, spatial distribution, and geochemical behaviour of critical minerals is essential for identifying tailings zones with the highest recovery potential and for assessing

associated environmental risks. Evaluating the critical mineral content of mine waste at abandoned mines in the Atlantic Provinces can support targeted exploration, strengthen secondary resource development, and contribute to Canada's broader strategy of securing domestic critical mineral supplies.

---

**Feldspars as monitors of fluid-mediated ore processes in magmatic-hydrothermal critical-metal deposits: theory and application**

---

DANIEL J. KONTAK

*Harquail School of Earth Sciences, Laurentian University, Sudbury, Ontario P3E 2C6, Canada <dkontak@laurentian.ca>*

The alkali feldspars (AFs), here simply albite (Ab) and K-feldspar (Kf), and quartz dominate the modal mineralogy of host rock granites and pegmatites for many critical metal (CM) ore deposit settings (i.e., Sn, W, Ta, Nb, Li, REEs), in many cases forming >90–95% of the mode. Whereas quartz is relatively robust in regard to fluid-rock interaction, which involves heated variably saline alkali-chloride solutions, the AFs are highly reactive with coupled dissolution precipitation (CDP) reactions producing textural and chemical changes that provide the means to monitor fluid interaction. Here the theory relevant to these changes in the AFs is first reviewed followed by its application to a variety of CM settings to illustrate its pragmatic application. A key textural feature noted is formation of perthite textures in Kf (e.g., film, flame, bleb) commensurate with a change from sanidine to orthoclase with development of pitted textures due to the presence of fluid inclusions (FIs) in newly precipitated Kf via CDP processes. Textural modification of Ab via CDP also occurs with FIs preserved along with growth of secondary apatite in P-rich settings. Synchronous with these textural changes is the progressive transition from magmatic to metasomatic chemistry of the AFs, thus Or<sub>70</sub> to Or<sub>95–100</sub>, and Ab<sub>85–90</sub> to Ab<sub>100</sub>. An important aspect of these latter changes is that primary magmatic textures are preserved, thus formation of metasomatic rocks that texturally look igneous as demonstrated by X-ray maps. Using this foundation, examination of several ore deposit settings reveals a strong correlation between CM mineralization and pervasive development of the textural and chemical features noted above. In fact, the CDP process is often so efficient that the host metasomatic rocks have and continue to be erroneously referred to as magmatic in nature.

---

**Rocks in the classroom: insights from teacher workshops and school visits**

---

LOUISE LESLIE<sup>1</sup> AND AARON C. ELSER<sup>2</sup>

*1. GeoLearns, Two Islands, Nova Scotia B0M 1S0 Canada  
<louise@geolearns.com>*

*2. Cobequid Educational Centre, 34 Lorne Street,  
Truro, Nova Scotia B2N 3K3, Canada*

A comprehensive rock kit was designed for rock identification activities and used at two teacher workshops and seven school visits. Fifteen kits were assembled with rock samples donated by the Nova Scotia Department of Natural Resources and materials that were purchased with grants from the Nova Scotia Mineral Resources Development Fund and the Canadian Geological Foundation. Each kit contains rock samples representing all rock types, materials for rock identification activities, and individual student materials, including a hand lens, ruler, reference guide and worksheets. Insights from the first teacher workshop led to redesigning the kit with a smaller number of rock samples that had textures that were more easily identifiable and increasing the number of kits used in the classroom to accommodate up to five groups of four to six students. Teachers at the second workshop experienced more hands-on time and activities that engaged them in greater depth than the first workshop using a similar format. School visits included: four different grade 4 classes, one day-long visit with several small groups of mixed primary to grade 6 students, one grade 7 class and one Geology 12 class. Insights from these visits led to minor changes in the kit's content, improvements in the demonstrations and the creation of detailed lesson plans redesigned into three modules consisting of a knowledge review, learning observational skills and naming rocks. The kit has proven to be an invaluable resource for teachers that engages students with learning rock identification skills and is applicable for several grade levels, starting at elementary and up to high school level.

---

**A review of carbonate cementation in the syn-rift strata of the Jeanne d'Arc Basin, Grand Banks, Newfoundland, Canada: challenges, models, and the potential for scalability using hyperspectral data**

---

DAVID LOWE, NOAH SLANEY, HODGSON JAIMEE, ERIC HUTTON, BRETTE HARRIS, AND HILARY CORLETT

*Department of Earth Sciences, Memorial University of Newfoundland, St. John's, Newfoundland and Labrador  
A1B 3X5, Canada <dlowe@mun.ca>*

The Jeanne d'Arc Basin on the Grand Banks contains significant potential for subsurface CO<sub>2</sub> storage within syn-rift marginal to shallow-marine Jurassic and Cretaceous sandstone reservoirs. However, uncertainties remain related to reservoir heterogeneity and reactivity to injected CO<sub>2</sub>, particularly concerning the distribution of carbonate cements. Carbonate cementation reduces reservoir capacity and connectivity, and may dissolve or reprecipitate during CO<sub>2</sub> injection, altering reservoir properties. Previous studies have identified multiple generations of carbonate cements in the Jeanne d'Arc Basin formed before and after burial compaction. Cements precipitated under near-surface conditions reflect strong paleoenvironmental controls. In the fluvio-deltaic Hibernia Formation, dolomite cementation is linked to seawater retention during sea-level regression, whereas calcite cementation corresponds to transgression and increased marine fluid flux into marginal-marine sediments. Similarly, in the storm-dominated Ben Nevis Formation, early calcite cements are associated with bioclast dissolution and early diagenetic fluid migration during marine transgression. Subsequent dissolution of early cements and precipitation of late ferroan calcite in both formations likely occurred during migration of basinal brines and/or meteoric fluids along faults. Despite this detailed reservoir-scale understanding, extrapolating carbonate cement distribution to the basin scale remains challenging due to limitations in identifying and mapping carbonate cements using conventional core, petrophysical, or seismic datasets. Hyperspectral imaging provides a promising basin-scale approach, as calcite and dolomite exhibit diagnostic absorption features in the short-wave (2325–2345 nm), mid-wave (3920–3960 nm), and long-wave (1114–1130 nm) infrared ranges. We have acquired hyperspectral data from over 7000 m of drill core spanning all cored clastic reservoir intervals in the Jeanne d'Arc Basin, with the goal of developing supervised and unsupervised classification workflows to map carbonate cement phases, calibrated against detailed sedimentological, petrographic, and geochemical analyses. This integrated approach aims to constrain carbonate cement paragenesis and evaluate the controls on cement precipitation/dissolution across the basin.

---

### Who takes a natural disasters class? (And where does it take them?)

---

JASON LOXTON

*Department of Math, Physics and Geology, Cape Breton  
University, Sydney, Nova Scotia B1M 1A2, Canada  
<jason\_loxton@cbu.ca>*

Like many institutions, Cape Breton University offers an introductory natural disasters course designed to serve as both

a recruitment tool for geology and a general introduction to science for Arts students. Over the past five years, students enrolled in this course have completed a pre-course survey to assess their needs, motivations, and expectations. This presentation discusses data gathered from over 300 students. Although some general trends emerge, e.g., enrollment was predominantly domestic and female-identifying, there was significant diversity in academic major, year of study, comfort level with science, and academic background. Participants generally fell into three subpopulations: upper-level science students seeking an engaging elective, general-interest students, and Arts or pre-Education students fulfilling a science requirement. Each of these groups reported unique needs and expectations upon entering the class. These are discussed, along with course changes implemented based upon these responses. End of course comments and teaching evaluations frequently highlighted an increased interest in science after taking our natural disasters class. Between 2021–2025, nearly 50% of the students enrolled in our two introductory courses had also taken our disasters course, demonstrating its effectiveness as a recruitment tool.

---

### Can free software produce high-quality Virtual Outcrop Models?

---

GRAYDON MACISAAC, JASON LOXTON,  
AND EDWARD J. MATHESON

*Department of Math, Physics and Geology, Cape Breton  
University, Sydney, Nova Scotia B1M 1A2, Canada  
<cbu23bkgs@cbu.ca>*

The use of Structure from Motion (SfM) photogrammetry has grown in the geoscience community over the past decade as drones have become affordable to the average consumer. Structure from Motion is a remote sensing technique that works by inputting overlapping photographs from different angles and using algorithms to triangulate the images. The technology enables geoscientists to create high-resolution 3D models of outcrops that can be annotated and interpreted from a remote location. The Cape Breton Sedimentology Research Group is in the process of creating a virtual library of 3D outcrop models from across Cape Breton Island. This poster documents our benchmarking process to determine photogrammetric best practices for in-house model creation. Complete results will be shared with the broader geoscience community to advance best practices and enhance accessibility of 3D SfM modelling. Our project required user-friendly software that generated high-resolution textures and meshes. We considered nine potential photogrammetry software packages. Two were chosen for detailed testing: Agisoft Metashape, the current industry standard for photogrammetric processing, and

RealityScan, a free software not widely mentioned in remote sensing literature. We conducted standardized tests using an HP Z2 G9 PC with a GeForce RTX 4060 graphics card and 32 GB of RAM. Tests included model processing time, point cloud density, mesh and texture quality, and software functionality. RealityScan is more versatile than the standard edition of Metashape, due to the shorter model generation time, reduced hardware requirements, lower cost, greater functionality, and higher-quality textures and meshes. However, Metashape is more compatible with some visualization software, as it less frequently requires model decimation to upload to 3D libraries. This research demonstrates that RealityScan is a viable low-cost alternative to Metashape that could facilitate more widespread adoption of SfM in geoscience research and education.

---

**Context matters: examining soft-sediment deformation structure terminology with insights from the Pomquet Formation, Cape Breton Island, Nova Scotia, Canada**

---

JESSICA MACISAAC<sup>1</sup>, EDWARD J. MATHESON<sup>1</sup>,  
JASON LOXTON<sup>1</sup>, AND MO E. SNYDER<sup>2</sup>

1. *Department of Math, Physics and Geology,  
Cape Breton University, Sydney, Nova Scotia B1M 1A2, Canada*  
<cbu21ftd@cbu.ca>

2. *Department of Earth and Environmental Science, Acadia  
University, Wolfville, Nova Scotia B4P2R6, Canada*

Classification of soft-sediment deformation structures (SSDS) can be difficult, as descriptive, diagnostic, and interpretive terminology and criteria vary widely among authors. This is particularly true when examining purported seismites, sedimentary deformation caused by earthquakes. Definitions of specific SSDS are inconsistently applied in the literature, reflecting differences in regional geology, scale of observation, and outcrop perspective/exposure, the possible trigger mechanisms for deformation inferred to be at play. This study examines pseudonodules and dish structures in the Pomquet Formation (Mabou Group) located near Creignish, Cape Breton Island, Nova Scotia, using detailed outcrop observation, ground- and drone-based photo analysis, and comparison to published global SSDS examples. The evidence suggests that classification and description of SSDS is strongly dependent on bed composition and architecture, the scale of features, and viewing orientation (e.g., plan-view vs. cross-section). These contextual controls are rarely emphasized in existing classification schemes, yet they strongly influence description and when compared with other reports of these SSDS, differences in scale and context are clear. These differences can make it difficult to determine how to classify the Pomquet Formation SSDS, because the pseudonodules and dish structures analyzed by

other geologists are not exactly the same as those at the Creignish section. These observations demonstrate that broadly defined SSDS categories may encompass structures with different origins and that generalized descriptive criteria should not be used in isolation to support specific genetic interpretations. Superficially similar SSDS may not share the same formative processes. As such, this study emphasizes the importance of context-driven descriptions of SSDS through careful analysis and side-by-side comparison of global SSDS to those closer to home, the latter of which illustrates the liquidization that was occurring in the Maritimes Basin, into the deposition of the Mabou Group.

---

**The human element: why emotion leads geoscience communication**

---

ALEX MACKAY

*Nova Scotia Department of Natural Resources Core Library,  
Westville, Nova Scotia B0K 1S0, Canada*  
<alex.mackay@novascotia.ca>

Public resistance to mining contrasts sharply with society's dependence on mined materials. Traditional outreach—explaining where minerals come from or calling out hypocrisy—has failed to connect. *The Human Element* display, developed in partnership with Geologize Inc., takes a different path: it begins with feeling, not facts. This presentation explores how evidence-based communication principles make geology relatable and meaningful. The exhibit starts with familiar human experiences: a grandmother hears ocean waves again thanks to rare earth elements in her hearing aids; a father fights an invisible hazard—radon gas—in his basement to protect his family; a community finds hope after a deadly stretch of highway is rebuilt with local aggregate and steel; and a senior traces family history through gypsum walls that have stood for generations. These narratives, told in Nova Scotians' own tone and cadence, reveal how geology quietly underpins health, safety, and heritage—without promoting industry, but by highlighting relevance. By framing geoscience as part of everyday life, this approach transforms outreach into an empathetic dialogue, bridging the gap between science and society.

---

**Research opportunities in Nova Scotia geoscience: hyperspectral and RGB drill core imaging for advanced research**

---

ALEX MACKAY

*Nova Scotia Department of Natural Resources Core Library,  
Westville, Nova Scotia B0K 1S0, Canada  
<alex.mackay@novascotia.ca>*

In 2023, a hyperspectral drill core scanning pilot project was completed at the Stellarton Drill Core library. The work, carried out by Scient Analytics and funded through the Mineral Resource Development Fund, scanned 30 drill holes from diverse geological settings across Nova Scotia. Using Short-Wave infrared (SWIR), Near-Wave infrared (NWIR), and high-resolution RGB imaging, the project captured detailed mineralogical and textural information. The hyperspectral data has been processed into ENVI-compatible files, creating a robust dataset ready for advanced interpretation. This resource offers significant opportunities for collaborative research—from undergraduate honors projects to Ph.D. studies—focused on mineral mapping, alteration characterization, and assessing the applicability of hyperspectral techniques across different geological environments, to name a few. The poster lists all scanned drill holes, including West Gore, Coxheath Hills, New Ross, East Kemptville, Walton, Brazil lake, and others. We invite researchers to engage with this dataset to identify where hyperspectral and RGB imaging provide the greatest geological insight and where limitations exist, fostering innovation in mineral exploration and geoscience applications.

---

### Quaternary erosion of the Canadian Shield from a 250-m $^{36}\text{Cl}$ depth profile in the Sudbury Igneous Complex, Ontario, Canada

---

LAUREN MACLELLAN<sup>1</sup>, JOHN GOSSE<sup>1</sup>, AND ALAN HIDY<sup>2</sup>

*1. Department of Earth and Environmental Sciences,  
Dalhousie University, Halifax, Nova Scotia B3H 4R2, Canada  
<lauren.maclellan@dal.ca>*

*2. Center for Accelerator Mass Spectrometry, Lawrence Livermore  
National Laboratory, Livermore, California 94550, USA*

Quantifying crustal erosion rates over million-year timescales in cratons remains challenging because tectonic stability, resistant lithologies, and low relief result in slow time-averaged erosion and poor preservation of erosional landforms. Low-temperature thermochronology, although powerful in tectonically active regions, typically models pre-Quaternary cooling histories in cratons and integrates exhumation over Phanerozoic timescales. limited constraints on late Cenozoic erosion in stable cratons leave a temporal gap between short-term (millennial) erosion methods and longer-term thermochronological exhumation estimates. Here, we present preliminary results from a novel approach being developed at CRISDal laboratory that uses the depth-dependent patterns of rare isotopes

produced in minerals by cosmic ray interactions to determine a Quaternary erosion rate for the Canadian Shield. in situ  $^{36}\text{Cl}$  concentrations were measured in feldspar separates from seventeen samples collected along a 250 m-deep drill core through compositionally homogeneous granophyre in the Sudbury igneous Complex. At these depths,  $^{36}\text{Cl}$  production transitions from neutron-dominated interactions near the surface to muon-dominated production, reducing sensitivity to short-lived surface effects such as ice sheet history and transient shielding owing to ephemeral lakes or sediment cover. Measured depth-dependent  $^{36}\text{Cl}$  concentrations were compared with predicted profiles calculated from semi-empirical formulas under varying erosion-rate scenarios, allowing Quaternary erosion rates to be estimated from the best-fitting models. Proximity to SNOIAB enables future integration of measured muon fluxes with physics-informed codes (e.g., MUTE and GEANT4), while extending the method to additional cosmogenic nuclides with differing half-lives will allow erosion histories to be examined across multiple timescales. This research provides a means to estimate erosion thickness over the next 1 myr above a proposed underground site in the Canadian Shield currently being evaluated for the long-term storage of Canada's nuclear waste.

---

### The oldest known dipluran associated with a new diverse assemblage of invertebrate body fossils preserved in a lycopod stump from the Carboniferous (Moscovian) Clifton Formation of New Brunswick, Canada

---

BAILEY MALAY<sup>1</sup>, LUCY L SMITH<sup>2</sup>, MATTHEW R STIMSON<sup>1,3</sup>, ROBERT A GASTALDO<sup>4,5</sup>, BRIAN L HEBERT<sup>6</sup>, MARGARET L SCOTT<sup>7</sup>, DUNCAN MCLEAN<sup>8</sup>, R. ANDREW MACRAE<sup>3</sup>, JOHN H CALDER<sup>3</sup>, STEVEN J HINDS<sup>9</sup>, ADRIAN F PARK<sup>9</sup>, PAGE QUINTON<sup>10</sup>, MICHAEL RYSEL<sup>10</sup>, TETSUTO MIYASHITA<sup>11</sup>, AND HILLARY MADDIN<sup>12</sup>

*1. Steinhammer Paleontological Laboratories, Natural Science Department, New Brunswick Museum, Saint John, New Brunswick E2K 1E5, Canada <baileymalay27@gmail.com>*

*2. Zoology Section, Natural Science Department, New Brunswick Museum, Saint John, New Brunswick E2K 1E5, Canada*

*3. Department of Earth Science, Saint Mary's University, Halifax, Nova Scotia B3H 3C3, Canada*

*4. Department of Geology, Colby College, Waterville, Maine E 04901, USA*

*5. Department of Paleobiology, Smithsonian National Museum of Natural History, Washington, DC 20560, USA*

*6. Fundy Treasures, 107 Highway 242, Joggins, Nova Scotia B0L 1A0, Canada*

*7. Department of Earth Sciences, Memorial University of Newfoundland, St. John's, Newfoundland and Labrador A1B 3X5, Canada*

8. *MB Stratigraphy Ltd., Sheffield, S9 5EA, UK*
9. *New Brunswick Department of Natural Resources, Fredericton, New Brunswick E3B 5H1, Canada*
10. *Department of Earth and Environmental Sciences, SUNY Potsdam, Potsdam, New York 13676, USA*
11. *Canadian Museum of Nature, Ottawa, Ontario K1P 6P4, Canada*
12. *Department of Earth Sciences, Carleton University, Ottawa, Ontario K1S 5B6, Canada*

A diverse assemblage of invertebrate body fossils has been discovered in the Pennsylvanian Clifton Formation (northern New Brunswick), which is composed exclusively of terrestrial strata. Palynostratigraphy of the formation places this rock in the “*Torispora Biozone*” indicating an upper Moscovian (315–307 Ma) age. Despite previous documentation in the Maritimes Basin of a diverse continental paleontological record, terrestrial invertebrates are rare in the Pennsylvanian. The invertebrate body fossils in the Clifton Formation are entombed in lycopod stumps that were rooted in a mineral paleosol. The taphonomic window is similar to that at the Joggins Fossil Cliffs, Nova Scotia. At Joggins, myriapods (*Xyloliulus*), eurypterid cuticle, and terrestrial gastropods (*Dendropupa*) were first discovered during the 19<sup>th</sup> century. The Clifton specimens are preserved on, and in, an oxidized mudstone associated with raindrop impressions, vertebrate skeletons, and invertebrate trace fossils. The specimens originate from three lycopsid stumps. One is a complete, 3-dimensionally preserved specimen of a Diplura, a basal hexapod lacking eyes, wings, and external mouthparts. Other invertebrates in the lycopod trees have yet to be described but include: xyloiuoidea myriapods with a compound eye; a terrestrial isopod; a paleocarid shrimp; and scorpion or eurypterid cuticle. To date, the indisputable dipluran record extends back only to the Cretaceous Period. A single, purported specimen is reported from Mazon Creek, Illinois, USA, but the interpretation is contentious. Hence, the Clifton dipluran is the oldest unequivocal example of the group and extends its fossil record back to the Pennsylvanian. These new invertebrate body fossils offer insight into the Carboniferous fauna of the Maritimes Basin in New Brunswick before a period of significant paleoenvironmental change that witnessed the onset of the Kazimovian rainforest collapse and aridification of the Maritimes Basin.

---

### Age and petrogenesis of Devonian peraluminous plutons in southwestern Newfoundland, Canada

---

KIERSTY M MALAY<sup>1</sup>, DONNELLY B ARCHIBALD<sup>2</sup>,  
JAMES CONLIFFE<sup>3</sup>, AND JAMES L CROWLEY<sup>4</sup>

1. *Department of Earth Sciences, Memorial University*

- of Newfoundland, St. John's, Newfoundland and Labrador A1C 5S7, Canada <kmmalay@mun.ca>*
2. *Department of Earth and Environmental Sciences, Saint Francis Xavier University, Antigonish, Nova Scotia B2G 2W5, Canada*
3. *Mineral Deposits Section, Geological Survey of Newfoundland and Labrador, St. John's, Newfoundland and Labrador A1B 4J6, Canada*
4. *Department of Geosciences, Boise State University, Boise, Idaho 83725, USA*

The Couteau Brook and Peter Snout plutons in southern Newfoundland are a complex composite intrusion of biotite-muscovite granite that formed from multiple magmatic pulses and possible remelting events. The plutons are tourmaline- and garnet-bearing and crop out over an area of approximately 10 km<sup>2</sup>. LA-ICP-MS and CA-ID-TIMS zircon dating of three samples from the Peter Snout pluton yielded ages between 419 ± 10 and 392 ± 10 Ma. Two samples of Couteau Brook pluton yielded ages between 397.49 ± 0.17 and 396 ± 9 Ma. Geochemical data show that both plutons are strongly peraluminous and show a negative REE slope with a negative Eu anomaly. Trace element patterns for both plutons show a slightly negative trend with elevated concentrations of elements such as Cs, Rb, and Ta. Tourmaline and mica compositions show Peter Snout is more geochemically evolved than Couteau Brook. Tectonic classification diagrams show that both plutons are syn-collisional and Nd-Sr isotopic data reflect a derivation from a continental crust source. Based on these data, Couteau Brook and Peter Snout are likely part of the same magmatic system, with some parts of the Peter Snout pluton representing a remelted version of older intrusions. Taken together, these plutons provide better age and tectonic constraints of this region of Ganderia in the Devonian.

---

### Deciphering the post-Grenvillian geological history in the Long Range Inlier, western Newfoundland, Canada: implications on gold mineralization

---

KEATON MARKHAM<sup>1</sup>, SHAWNA WHITE<sup>1</sup>, MO SNYDER<sup>2</sup>,  
AND SEBASTIAN KOMMESCHER<sup>3</sup>

1. *Department of Earth Science, Saint Mary's University, Halifax, Nova Scotia B3H 3H3, Canada <keaton.markham@gmail.com>*
2. *Department of Earth and Environmental Science, Acadia University, Wolfville, Nova Scotia B4P 2R6, Canada*
3. *Core Science Facility, Memorial University of Newfoundland, St. John's, Newfoundland and Labrador A1C 5S7, Canada*

The geologic history of Appalachian orogenesis in Newfoundland, involving Paleozoic episodic accretion of microcontinents and terranes, is the primary reason for gold occurrences across the province. Although most occurrences occur in central Newfoundland in the peri-Gondwanan terranes, recent exploration has uncovered occurrences within

the Mesoproterozoic (ca. 1.0 Ga) Long Range Inlier (LRI), a Laurentian massif of crystalline rocks. Gold showings are spatially associated with regional, northwest-trending LiDAR lineaments, which parallel, and merge with, basement faults farther west. These observations suggest structural control on gold occurrences, demonstrating significant economic potential; however, locating deposits is hindered by a lack of geoscientific research in the LRI. This study therefore combines geological mapping and geochronology to better constrain the inlier's geological history. Results show a metasedimentary sequence, previously grouped with Mesoproterozoic rocks, nonconformably overlies crystalline basement. Mafic units intruded metasedimentary rocks sub-parallel to bedding. These rocks, originally correlated with Mesoproterozoic amphibolite, are classified as dioritic-gabbroic, and their extent and textural variability suggests a plutonic suite origin. Preliminary U–Pb dates from LA-ICP-MS analysis of zircon from the gabbro are ca. 425 Ma, similar to the ca. 430 Ma Taylors Brook Gabbro, the only known Silurian mafic unit in the region. LiDAR lineaments correlate with sinistral faults with pervasive gouge, breccia, and slickensides, and mineralized veins and altered host rock are pervasive within the deformation zone. Ongoing K/Ar dating of illite in gouge will constrain the timing of fault motion. This work refines the geological history of the LRI by identifying a Paleozoic stratigraphic succession that is used as a marker horizon to track deformation post 1.0 Ga. A major sinistral fault system offsets these marker beds and clearly focused late mineralizing fluids. Recognition of Silurian magmatism, combined with recently documented Silurian deformation in western Newfoundland, implies that the Salinian orogeny may be farther reaching than previously thought.

---

### Minerals and metals critical for the energy transition in Canada

---

TANIA MARTINS

*Manitoba Geological Survey, Winnipeg, Manitoba R3G 3P2, Canada <vania.martins@gov.mb.ca>*

Critical minerals are clearly a current topic of interest and research. The Government of Canada released a list of 34 elements and raw materials considered vital for the sustainable economic success, national security, decarbonization and strategic interests of the country and its allies. Additionally, Canada has demonstrated to be committed to grow domestic and global value chains for the green and digital economy by creating funding research opportunities and infrastructure investment around critical minerals. This is a trend also observed outside of Canada, with many nations in-

terested in fostering the search for critical minerals and their research, especially giving the importance of the exploration and mining industry to any country's economic development. Of Canada's list, one of the considered critical and strategic elements is lithium. This element has important physical and chemical properties that are suited to a wide range of applications including pharmaceuticals, glass and ceramics, and aerospace technologies. Furthermore, lithium is currently considered one of the most important critical metals for the energy transition due to its importance in the lithium-ion battery technology for electric vehicles (EVs), portable electronic devices, electric tools, and energy grid storage applications. Canada is a hard-rock lithium producer from granitic pegmatites and has extensive reserves of this resource, but it also holds significant resources associated with brines. In Nova Scotia, Brazil Lake is the most prospective area for hard-rock lithium with recent resource drilling and new discoveries being made. Other elements are also critical for these same low-carbon technologies such as copper, graphite, nickel and cobalt to name a few. All these critical elements and raw materials occur in a variety of deposits, geological settings and mineral associations. Canada is well endowed in these types of commodities and well positioned to offer a competitive and reliable supply.

---

### Your path to professional geoscience: education, licensure, and careers

---

MAUREEN MATTHEW<sup>1</sup>, NATASHA MACADAM<sup>2</sup>,  
AND MAYLIA PARKER<sup>3</sup>

1. *BGC Engineering, Halifax, Nova Scotia B3J 2C6, Canada <mmatthew@bgcengineering.ca>*
2. *Nova Scotia Department of Energy, Halifax, Nova Scotia B3J 3M8, Canada*
3. *Geoscientists Nova Scotia, Dartmouth, Nova Scotia B3B 1H7, Canada*

Geoscience is a regulated profession in most of Canada, with legislative acts restricting the practice of geoscience to individuals and companies registered with self-governing professional associations. In Nova Scotia, this role is fulfilled by Geoscientists Nova Scotia, formally known as the Association of Professional Geoscientists of Nova Scotia (APGNS). In some jurisdictions, like New Brunswick and Newfoundland, geoscience is jointly regulated with engineering practice under APEGNB and PEGNL, respectively; however, all associations share common requirements and mandates set by Geoscientists Canada. APGNS and its counterparts across Canada are responsible for upholding high standards of geoscience practice and education through the regulation of admissions, provision of disciplinary oversight, and protection of

the professional title “geoscientist” from misuse by unlicensed practitioners. For students studying the geosciences, the purpose of professional associations, the academic requirements for licensure, and the path toward professional registration can be difficult to navigate. This presentation will help clarify the role of associations, outline the educational requirements for professional membership, and provide guidance on course selection to support future licensure. Early-career mentoring and experience requirements that precede full professional registration will be highlighted. In addition to regulatory and academic considerations, the presentation will explore what geoscience careers can look like in practice, drawing on examples from academia, consulting, industry, and the public sector. By linking licensure requirements with real-world career pathways, the session aims to help students better understand how their academic choices translate into professional opportunities. Time will be reserved for open discussion and questions to ensure students leave with a clear understanding of their course requirements and the tools needed to develop individualized plans for a successful early career in the geosciences.

---

**Evaluating the cosmogenic nuclide catchment erosion rate method on the Margaree River, Cape Breton Island, Nova Scotia, Canada**

---

EMMA MCCARRON AND JOHN GOSSE

*Department of Earth and Environmental Sciences,  
Dalhousie University, Halifax, Nova Scotia B3H 4R2, Canada  
<em517837@dal.ca>*

Terrestrial cosmogenic nuclides (TCNs) are widely used to establish tributary to mountain range-scale differences in stream catchment erosion owing to tectonic and climatic forcing over millennial timescales. Increasingly, this method is used to establish pre-industrial baseline erosion rates of catchment slopes to evaluate landscape responses to anthropogenic climate change. The catchment-wide average erosion rates (CWAER) can be readily converted to total sediment discharge (bed, suspended, and dissolved load). The approach assumes that catchment slopes have experienced sufficient exposure to cosmic radiation to produce the expected exponential decrease in nuclide concentration with soil depth. However, in recently deglaciated catchments, exposure to TCN irradiation may be too short for this assumption to hold. Insufficient exposure, combined with glacial sediments containing variable inherited TCN concentrations, can yield depth profiles with low concentrations that are easily overprinted by inherited signals. Despite this limitation, the validity of CWAER methods under these conditions has

not been systematically tested. To address this gap, we will determine CWAERs for the Northeast Margaree River using cosmogenic  $^{10}\text{Be}$  in quartz isolated from coarse stream-sand samples from two first-order tributaries, a steep downstream second-order tributary, and the mainstem in Inverness County. Two soil samples - a saprolite exposed on the upland barrens and a till 5 km upstream of the mainstem - will be used to evaluate the inherited nuclide concentrations in slope sediment. Based on catchment slope, area, and stream discharge, we expect the two upper catchments to exhibit relatively low  $^{10}\text{Be}$  erosion rates, the third a substantially higher erosion rate owing to steep long slopes, and the mainstem to yield an intermediate erosion rate, reflecting its mixed slope conditions. If successful, this study will provide Nova Scotia's first baseline erosion rates and suggest that the method can inform investigations of recent climate and anthropogenic impacts on stream bank erosion.

---

**Age and tectonic evolution of the Clyburn Brook Formation, Cape Breton Island, Nova Scotia, Canada**

---

DYLAN J. MCKEEN, DEANNE VAN ROOYEN, SANDRA M. BARR, CHRIS E. WHITE, AND ROBERT P. RAESIDE

*Department of Earth and Environmental Science,  
Acadia University, Wolfville, Nova Scotia B4P 2R6, Canada  
<0304624m@acadiau.ca>*

The Eastern Highlands Shear Zone (EHSZ) in the Cape Breton Highlands juxtaposes the Ganderian Aspy terrane consisting primarily of Ordovician to Silurian metavolcanic and metasedimentary rocks and abundant middle Paleozoic granitoid plutons with the Ganderian Bras d'Or terrane that consists mainly of Ediacaran metasedimentary rocks and Ediacaran to Cambrian plutons. The Clyburn Brook Formation (CBF) is an assemblage of interlayered, low-grade, bimodal metavolcanic, and metasedimentary rocks which straddles the northeastern part of the EHSZ. Based on lithological and geochemical similarities and interpreted age, the CBF has been correlated with Silurian units in the Aspy terrane which are arc-related based on chemical characteristics. New LA-ICP-MS detrital zircon analysis indicate maximum depositional ages of ca. 420 Ma from calcareous metaarkose and ca. 410 Ma from subarkosic metasandstone in the lower and upper epiclastic members, respectively, suggest that the CBF formed as a syn-convergent pull-apart basin that received detrital material from Silurian arc-related igneous activity that post-dated initial convergence along the EHSZ. Silurian volcanic activity at 430 to 420 Ma is well-documented in the Sarach Brook, Calumruadh Brook, and Money Point formations of the adjacent Aspy terrane. Based on the MDAs reported here the CBF may be correlative with the upper parts of

those units. The CBF was intruded by largely undeformed Cameron Brook granodiorite (CBG) previously dated at  $402 \pm 3$  Ma (U–Pb TIMS zircon), suggesting that terrane juxtaposition was accompanied by Devonian syn-collisional magmatism. This interpretation is supported by a new LA-ICP-MS zircon igneous crystallization age of  $395.8 \pm 2.2$  Ma from felsic metatuff in the middle volcanic member interpreted as the volcanic equivalent of the CBG, suggesting that the CBF mostly post-dates the initial collision between terranes interpreted to have begun between ca. 430 and 420 Ma. The CBF records syn-convergent deformation and folding, as well as post-collisional brittle deformation, resulting from transpressional deformation during protracted convergence of the Aspy and Bras d’Or terranes.

---

**Zircon chemistry-geochronology and melt inclusion constraints on the origin and critical metal endowment of late Devonian granitoid rocks in the Port Mouton Pluton, Meguma Lithotectonic Zone, Nova Scotia, Canadian Appalachian orogen\***

---

GEENA M. MORSE<sup>1</sup>, ANNA TEREKHOVA<sup>1</sup>, JACOB HANLEY<sup>1</sup>, JAMES CROWLEY<sup>2</sup>, AND BRENDAN MURPHY<sup>3</sup>

1. *Department of Earth Science, Saint Mary’s University, Halifax, Nova Scotia B3H 3H3, Canada <geena.morse@smu.ca>*
2. *Department of Geosciences, Boise State University, Boise, Idaho 83725, USA*
3. *Department of Earth and Environmental Sciences, St. Francis Xavier University, Antigonish, Nova Scotia B2G 2W5, Canada*

Late Devonian satellite plutons (Shelburne, Barrington Passage, and Port Mouton ‘PMP’) in southwestern Nova Scotia, coeval with the much larger South Mountain Batholith, show felsic magma interaction with coeval mafic magmas. In the PMP, field and petrographic observations, bulk rock geochemistry, and EPMA characterized mafic “enclaves” to constrain the conditions during interaction of mafic magmas with coeval felsic magmas, and influence on the critical metal endowment of late Devonian granitoid magmas. Oxythermobarometry data from peritectic amphibole (after pyroxene) indicate that values of  $fO_2$  in the mafic magmas shifted from FMQ+4 to FMQ+1 over a cooling window from  $\sim 850$ – $700^\circ\text{C}$  during this interaction, at pressure of up to  $\sim 390$  MPa (14.7 km), much deeper than that of final magma emplacement. LA-ICP-MS zircon geochemistry and U–Pb dating show that PMP zircon populations shift in predominant age populations across lithologies. Zircons from mafic lithologies (rounded, corroded, inherited, or xenocrystic) at Forbes Point returned exclusively Neoproterozoic ages (ca. 550 Ma), whereas early granodiorites have predominantly Neoproterozoic ages (ca. 560–620 Ma) and rare late Devonian (ca. 390 Ma) ages (consistent with PMP emplacement), with the balance shifting almost

entirely to 370–390 Ma zircons in late-stage granodiorites and hybrid tonalites (formed by mixing of mafic and felsic magmas). PMP-coeval zircon from the felsic-intermediate rocks yield: (i) REE-HFSE ratios that diverge from typical magma fractionation trends, and (ii) values of  $fO_2$  that are widely variable in PMP-coeval zircon ranging from FMQ+1 to FMQ-5, decreasing with progressive fractionation. These findings suggest re-equilibration and thermal disturbance of small volumes of crystallizing felsic magmas by mafic magmas and strongly reducing wall-rock lithologies during deep magma gestation. Melt inclusions in zircon in the tonalite have low granophile critical metal (Sn, W, Mo) concentrations, comparable to average upper crust. The dominant Neoproterozoic ages in mafic rocks and early granodiorites in the PMP sparks new questions about the timing and source of mafic end-member magmas.

*\*Winner: AGS Rupert MacNeill Award for best undergraduate student oral presentation*

---

**Post-accretionary emplacement of granitoid plutons in the Meguma terrane, Nova Scotia, Canada: transition from dextral transpression to transtension?**

---

J. BRENDAN MURPHY<sup>1</sup>, ROB A STRACHAN<sup>2</sup>, WILLIAM J COLLINS<sup>3</sup>, DONNELLY B ARCHIBALD<sup>1</sup>, JACOB HANLEY<sup>4</sup>, AND GEENA M MORSE<sup>4</sup>

1. *Department of Earth and Environmental Sciences, St. Francis Xavier University, Antigonish, Nova Scotia B2G 2W5, Canada <bmurphy@stfx.ca>*
2. *School of Earth and Environmental Sciences, University of Portsmouth, Portsmouth PO1 2EG, UK*
3. *School of Earth and Planetary Sciences, Curtin University, Perth, Western Australia 6102, Australia*
4. *Department of Earth Science, Saint Mary’s University, Halifax, Nova Scotia B3H 3H3, Canada*

Continental crust forms primarily where processes associated with subduction transfer mass and heat from the mantle to the base of the crust. These processes are encapsulated in the concept of trans-crustal magmatic systems (TCMS), in which mafic underplating along the base of the crust energizes a deep crustal hot zone (DCHZ), from which buoyant silicic magma rises to form a vertically interconnected network of reservoirs. We provide a synthesis of recent studies on the origin and emplacement of the voluminous peraluminous granitoid magmas in the Meguma terrane. Magma genesis began at ca. 400 Ma in the DCHZ and gestated for 25–30 myr before emplacement in the middle-to-upper crust in discrete episodes. Protracted magmatism in the lower crust was the product of ongoing subduction, coeval with regional metamorphism and dextral transpressional polyphase deformation of upper crustal rocks. Ascent of granitoid magma through the crust, beginning at ca. 380 Ma and continuing until

ca. 370 Ma, occurred during the transition from transpression to dextral transtension, distributed strain to focused strain, and the development of through-going shear zones. These shear zones were a combination of Reidel and P-shears which produced an interconnected network of conduits that facilitated the magma ascent and emplacement at depths ranging from 10–20 km. Granitoid magmas created rheological barriers, impeding the ascent of coeval mafic magmas except along the shear zones. Field relationships, age data, and geochemistry of these volumetrically subordinate mafic intrusions are evidence of mafic underplating and imply coeval emplacement of mantle-derived magma in a continental arc setting. Clasts of Meguma-derived granitoid rocks in ca. 360 Ma unconformably overlying continental clastic strata of the Horton Group imply ~10 km exhumation of the Meguma terrane between 370 and 360 Ma during protracted regional dextral transtension culminating in the composite late Devonian–Early Permian Maritimes Basin.

---

### Using apatite chemistry to distinguish Li-Cs-Ta-fertile and barren pegmatites

---

CLAUDE NAMBAJE<sup>1</sup>, CHRISTOPHER R.M. MCFARLANE<sup>1</sup>, TANIA MARTINS<sup>2</sup>, MAX KACZMER<sup>3</sup>, AND LEE A. GROAT<sup>4</sup>

1. *Department of Earth Sciences, University of New Brunswick, Fredericton New Brunswick E3B 5A3, Canada*  
<claudenambaje@unb.ca>

2. *Manitoba Geological Survey, Winnipeg, Manitoba B R3G 3P2, Canada*

3. *New Age Metals Inc., Richmond, British Columbia V6Y 1M3, Canada*

4. *Department of Earth, Ocean and Atmospheric Sciences, University of British Columbia, Vancouver British Columbia V6T 1Z4, Canada*

Apatite major- and trace-element chemistry offers a powerful tool for assessing the fertility of granitic pegmatites and distinguishing Li-Cs-Ta-bearing systems from barren counterparts. Across representative pegmatites examined in this study—including a case-study suite from the Bird River greenstone belt, Manitoba, Canada—apatite compositions systematically record magmatic differentiation and volatile enrichment processes linked to rare-element mineralization. Apatites from fertile pegmatites are characterized by elevated Mn and higher Mn/Fe ratios, together with enrichment in Li, U, and Th. These are accompanied by lower Eu<sub>N</sub> values, subdued total rare-earth-element abundances, pronounced negative Eu anomalies, and subtle but diagnostic REE tetrad effects, reflecting advanced melt evolution. By contrast, apatites from barren pegmatites typically show higher ΣREE, elevated Eu<sub>N</sub> values and weaker Eu anomalies, lower Mn/Fe and Li contents. integrated compositional thresholds—parti-

cularly Mn enrichment, low Eu and ΣREE, and Li-U-Th enrichment—provide a reproducible geochemical fingerprint for fertile pegmatites. These results demonstrate that apatite is an applicable recorder of magmatic-hydrothermal evolution and an effective discriminator of pegmatite fertility, with direct implications for exploration targeting of critical-metal systems.

---

### Assessing the impacts of river-based carbon sequestration: effects of river alkalinity enhancement on dissolved metal concentrations in the West River, Nova Scotia, Canada

---

ROWAN NORRAD<sup>1</sup>, SHANNON STERLING<sup>2</sup>, AND IAN SPOONER<sup>1</sup>

1. *Department of Earth and Environmental Science, Acadia University, Wolfville, Nova Scotia B4P 2R6, Canada*  
<renorrad@gmail.com>

2. *Department of Earth and Environmental Sciences, Dalhousie University, Halifax, Nova Scotia B3H 4R2, Canada*

Low pH conditions in the West River, Pictou County, Nova Scotia, are a result of both natural and anthropogenic processes that have resulted in elevated metal concentrations which can be toxic to aquatic flora and fauna. River alkalinity enhancement (RAE) is a process that has been applied to the West River in which crushed alkaline minerals are added to the river to has the potential to sequester atmospheric carbon dioxide. RAE also enhances habitat quality by raising river pH which theoretically reduces dissolved metals. In this study, two years (2024–2025) of water chemistry and quality data were collected from three sites along the West River. Sites were selected upstream of the application location, directly downstream of the dosing location, and at the head of tide. Towards the end of winter 2025 (March through May), data were collected while RAE was taking place within the river. Water samples were collected using polyethylene bottles and submitted to the Dalhousie Clean Water lab where metals were measured using ICP-MS. Temperature, pCO<sub>2</sub>, pH and conductivity data were also collected using continuous autonomous in-situ monitoring stations. Preliminary comparisons between the application window in the control year (2024) and treatment year (2025) show that concentrations of dissolved Al, Fe, Cu, Mn, and Zn vary greatly. in 2025 only Al concentrations appeared to vary between upstream and downstream sites. Data indicates that the severe drought in 2025 had an impact on dissolved metal concentrations in the West River. However, the lack of significant trends between collection sites may demonstrate that the application of RAE for carbon dioxide removal does not have a pronounced effect on the concentration of dissolved metals. These results have the potential

to provide transportable risk assessment guidance for RAE projects. Future work includes the integration of hydrogeological parameters in metal concentration assessment.

---

### Dissolved inorganic carbon stable and radiocarbon isotope distributions in slope waters of Nova Scotia and southern Newfoundland, Canada

---

LAUREN T. O'REILLY<sup>1</sup>, BRETT D. WALKER<sup>2</sup>,  
AND OWEN A. SHERWOOD<sup>1</sup>

1. *Department of Earth and Environmental Sciences,  
Dalhousie University, Halifax, Nova Scotia B3H 4R2, Canada  
<lauren.oreilly@dal.ca>*

2. *School of Physical Science, University of California,  
Irvine, California 92697-3100, USA*

Marine dissolved inorganic carbon (DIC; ~38 000 GtC) is the largest reservoir of oceanic dissolved carbon. Carbon isotopes ( $\delta^{13}\text{C}$  and  $\Delta^{14}\text{C}$ ) of marine DIC are useful for tracing water mass advection and mixing, residence times, and carbon cycling. Few DIC  $\delta^{13}\text{C}$  and  $\Delta^{14}\text{C}$  depth profiles have been published for the Northwest Atlantic off Nova Scotia and southern Newfoundland, which is one of the fastest-warming regions globally. We report slope water DIC stable ( $\delta^{13}\text{C}$ ) and radiocarbon ( $\Delta^{14}\text{C}$ ) depth profiles from the Gully Marine Protected Area off Nova Scotia, and a coral conservation area (Division 30) off the southwest Grand Banks of Newfoundland. The subsurface water masses in these areas comprise Warm Slope Water (WSW:  $\theta = 10^\circ\text{C}$ ,  $S = 35.2$ ), which is associated with the northward flowing Gulf Stream, and Labrador Slope Water (LSW:  $\theta = 6^\circ\text{C}$ ,  $S = 34.9$ ), which is formed in part from outflow from the Canadian Arctic Archipelago. We find that these water masses have distinct DIC  $\delta^{13}\text{C}$  and  $\Delta^{14}\text{C}$  isotopic signatures, with WSW having higher DIC  $\Delta^{14}\text{C}$  values and lower DIC  $\delta^{13}\text{C}$  values than LSW, suggesting a larger contribution of “bomb”  $^{14}\text{C}$  and higher anthropogenic carbon inputs to WSW. Lower DIC  $\Delta^{14}\text{C}$  values in LSW are consistent with dilution effects of deep convection in the Labrador Sea and atmospheric isolation of Arctic halocline waters. These results, when paired with proxy bioarchival data, will better constrain our understanding of water mass circulation, carbon cycling, and AMOC variability in the Northwest Atlantic.

---

### Recording peak metamorphic conditions in garnet-rich schists from the Dalradian highlands of Scotland

---

KARI OSFOLK AND RICHARD COX

*Department of Earth and Environmental Sciences,  
Dalhousie University, Halifax, Nova Scotia B3H 4R2, Canada  
<osfolk.kari@gmail.com>*

Determining the timing of metamorphism is key to understanding the relationship between deformation, mineral growth, and crustal evolution during mountain building. This study investigates peak-metamorphic assemblages within the garnet zone of the Dalradian Supergroup in the West Highlands of Scotland. Samples were mapped and collected from Rubha Cladaich (Loch Leven Schist) and Cononish Glen (Ben Lui Schist), two areas that record distinct deformation histories. Garnet growth occurred syn- $D_2$  at Rubha Cladaich and post- $D_1$  to pre- $D_2$  at Cononish Glen, providing an opportunity to examine how metamorphism and deformation were linked during different stages of the Caledonian Orogeny. Petrographic and microstructural analyses are being used to characterize garnet textures, inclusion trails, and mineral assemblages in order to interpret pressure-temperature conditions and the timing of metamorphic events. Planned analytical work includes trace-element mapping of garnet (REE and Y zoning), garnet end-member determination, and thermobarometric analysis, including application of the garnet-biotite and garnet-muscovite thermometers to estimate peak temperatures. By comparing metamorphic features between these two Dalradian localities, this research aims to identify variations in metamorphic timing and intensity across the region. The results will help clarify how garnet growth relates to deformation phases and crustal thickening during the Caledonian orogeny. Ultimately, this work contributes to a better understanding of the metamorphic history and tectonic development of the Scottish Highlands.

---

### Characterizing the evolution and magmatic endowment of the Mount Douglas and McDougall Brook granite suites, southwestern New Brunswick, Canada

---

CHLOE OXNER, JACOB HANLEY, SARAH SPEIGHT, AND  
GEENA MORSE

*Department of Earth Science, Saint Mary's University, Halifax,  
Nova Scotia, B3H 3C3, Canada <Chloe.Oxner@smu.ca>*

Polymetallic (Sn-W-Mo-In-Zn-Bi) deposits occur in the Mount Pleasant Granites (MPG) within the Mount Pleasant Caldera Complex, on the northern margin of the Saint George Batholith (SGB) in southwestern New Brunswick. Polymetallic (Sn-W-Mo) mineral occurrences are also hosted in the Mount Douglas Granite (MDG), a granitic suite in the SGB. The intracaldera McDougall Brook Granites (MBG) represent barren, precursor intrusions to the MPG and are genetically related to the min-

eralized MPG based on similarities in textures, geochemistry, crystallization ages, and mineralized assemblages. The MDG may be a parental magma source for the MBG, and coeval with mineralization in the MPG, although relationships remain unresolved. Zircon from the MDG is likely overprinted and reworked, commonly seen in “cold” granites, and inherited zircon cores result in an overlap in reported age ranges for emplacement and crystallization of the MDG and MBG. To address these uncertainties, this study integrates petrography, whole rock geochemistry, oxythermobarometry, and halogen systematics from accessory mineral chemistry (apatite, titanite, zircon) to constrain the crystallization and emplacement P-T conditions, degassing history, and oxygen fugacity of these granitic magmas. The overall objective is to resolve and characterize processes that influenced the variable magmatic mineralization potential of these granitoids. Results from petrographic analysis document multiple processes relevant to the metal endowment in the MDG, including magma mixing, volatile saturation, sulphide saturation, and late magmatic oxidation. As well, preliminary whole rock geochemical results show pronounced depletions in Eu, Sr, Ba, and Ti in MDG samples, indicating that magmas experienced extensive melt fractionation with significant crystal accumulation prior to emplacement, a key process for metal enrichment.

---

### Tectonic framework for Metasomatic Iron Alkali-Calcic (Iron Oxide Copper Gold) mineralization systems in southern New Brunswick, Canada

---

ADRIAN F. PARK<sup>1</sup>, STEVEN J. HINDS<sup>1</sup>, SUSAN C. JOHNSON<sup>2</sup>,  
STEVEN L. E. ROSSITER<sup>2</sup>, AND AYLALEW GEBRU<sup>1</sup>

1. *New Brunswick Department of Natural Resources, Fredericton, New Brunswick E3B 5H1, Canada* <adrian.park@gnb.ca>
2. *New Brunswick Department of Natural Resources, Sussex, New Brunswick E4E 5L2, Canada*

MIAC (Metasomatic Iron Alkali-Calcic) systems are large-scale hydrothermal alteration systems related to mineralization involving many critical elements. Within MIAC systems IOCG (Iron Oxide Copper Gold) deposits can also host important reserves of Cu, U, Th and REEs, and Five Element vein deposits can host important reserves of Cu, Co, Ni, Ag, and Bi. A thermal model for MIAC systems posits a series of deposit types from high to low temperature related to various ‘alteration facies.’ The challenge of mapping out the geometry of these facies is only compounded when deformation has occurred during and after MIAC systems form. Southern New Brunswick illustrates an example of how deformation related to major strike-slip faults reorganizes (frag-

ments) the products of a MIAC system(s). The same faults provide the fluid pathways necessary for this crustal-scale hydrothermal activity. Positive flower structures (transpressive) especially when transtensional relaxation follows their formation, are related to several examples of MIAC alteration. In southern New Brunswick the Bellisle, Kennebecasis, and offshore Cobequid–Chedabucto faults are the principal structures involved, with associated flower structures recognized from the Mascarene Peninsula to Quaco Head. In eastern Caledonia large flower structures have been modified by strike-slip fault overprinting and extensive salt movement. Extensive quartz-mylonite zones derived from granitoid and felsic volcanic protoliths show evidence of silica flooding and Fe, K, Ca, and Na depletion, whereas massive specular hematite, siderite, and magnetite mineralization with associated Cu and Au in shear zones and brecciated zones represent the mineralization. Five Element veins are also represented. Stratabound redox boundary deposits represent later fluids channeled by faults, derived from dewatering in the Cumberland Basin. Geochronological constraints suggest that the systems were operative from the early Devonian to middle Permian. Positive flower structures with evidence of late transtensional relaxation from Tournaisian to late Permian are potentially the most prospective areas.

---

### Critical mineral distribution and resource potential in tailings from the abandoned Stirling Zn-Pb-Cu-Au-Ag Mine, Nova Scotia, Canada

---

MICHAEL B. PARSONS<sup>1</sup>, DUANE C. PETTS<sup>2</sup>, JEANNE B. PERCIVAL<sup>2</sup>, TERRY C. CHENG<sup>3</sup>, AND JULIE DERIAZ<sup>3</sup>

1. *Geological Survey of Canada, Natural Resources Canada, Dartmouth, Nova Scotia B3B 1A6, Canada* <michael.parsons@nrcan-rncan.gc.ca>
2. *Geological Survey of Canada, Natural Resources Canada, Ottawa, Ontario K1A 0E8, Canada*
3. *Canmet MINING, Natural Resources Canada, Ottawa, Ontario K1A 0G1, Canada*

Increasing global demand for critical minerals and lengthening timelines for permitting new mines have led to strong interest in reprocessing legacy mine wastes. There are more than six billion tonnes of tailings at active and abandoned metal mines across Canada; however, many questions remain concerning the economics, technical feasibility, and regulations associated with reprocessing. The goal of this study is to evaluate the critical metal content and mineralogy of tailings at the former Stirling Mine in Cape Breton, the spatial distribution and recoverability of critical metals, and the potential for reprocessing to help reduce environmental risks. The Stirling Mine operated from 1927–1931, 1935–1938, and 1951–1956, extracting Zn, Pb, Cu, Au, and Ag from a vol-

canogenic massive sulphide deposit and generating approximately 930 000 tonnes of tailings. In 2023 and 2024, we collected tailings from hand-dug pits to investigate the effects of weathering in near-surface tailings. In June 2024, we drilled 33 boreholes through the tailings, recovering samples for geochemical, mineralogical, particle size, and metallurgical testing. Geochemical analyses of the tailings using ICP-ES/MS following 4-acid digestion (and fire assay for Au) show the following median metal concentrations (n = 152): 1.19 wt.% Zn, 0.35 wt.% Pb, 0.18 wt.% Cu, 0.50 g/t Au, and 21 g/t Ag. A subset of tailings was analyzed using ICP-ES/MS following sodium peroxide fusion, revealing the following median concentrations of potential by-product critical elements (n = 51): 6.6 mg/kg Ga, 1.1 mg/kg Ge, and 0.80 mg/kg In. Critical minerals in the tailings are hosted by sulphide, sulphosalt, and silicate minerals including sphalerite, chalcopyrite, galena, tennantite, chlorite, muscovite, and talc. Metal-bearing, weathering-related secondary phases are present in the near-surface oxidized tailings. Results from this study are being used to assess the viability of reprocessing historical base metal tailings to recover valuable elements and reduce risks to local ecosystems and surrounding communities.

---

### Glacial dispersal from the Brazil Lake lithium-cesium-tantalum pegmatites, southwest Nova Scotia, Canada

---

ROGER PAULEN<sup>1</sup>, BETH McCLENAGHAN<sup>1</sup>, CHRIS BECKETT-BROWN<sup>2</sup>, DENISE BRUSHETT<sup>3</sup>, AND JESSEY RICE<sup>1</sup>

1. Geological Survey of Canada, Natural Resources Canada,  
Ottawa, Ontario K1A 0E8, Canada

<roger.paulen@nrcan-rncan.gc.ca>

2. Ontario Geological Survey, Sudbury, Ontario P3E 6B5, Canada

3. Nova Scotia Department of Municipal Affairs, Halifax,  
Nova Scotia B3J 2M4, Canada

The Brazil lake lithium-cesium-tantalum (LCT)-type pegmatites (North, South, and Army Road) in southwestern Nova Scotia were the focus of a detailed till study to document glacial dispersal through several documented phases of glaciation. A total of 84 bulk (7–20 kg) till samples, from mostly subglacial traction till, were collected from 77 sites near and down ice of the three pegmatites. Glacial dispersal of indicator minerals from the pegmatites is best defined by spodumene (the primary Li ore mineral) in the 0.25–0.5 mm fraction of the 2.8–3.2 specific gravity till concentrates, and its presence in the local till (100s to 1000s of grains per sample) confirms the existence of nearby LCT pegmatites. Results indicate that spodumene dispersal is the product of multiple ice-flow events with varying trajectories, and formed a broad, fan-shaped footprint extending at least 12 km down ice that is considerably larger than the pegmatite subcrop

areas. Other useful LCT pegmatite indicator minerals identified during this study include apatite and tourmaline (10 to 100s of grains), and lesser amounts of columbite-tantalite, cassiterite, and scheelite (1 to 10s of grains), although their spatial distribution in till is limited compared to spodumene. Geochemical analyses of three till size fractions was completed including coarse sand (1.0–2.0 mm), silt + clay (<0.063 mm), and clay (<0.002 mm) using various digestion and fusion methods to identify best methods to detect LCT pegmatites. Lithium dispersal from the Brazil lake pegmatites is best defined by the coarse fraction (Na-peroxide fusion), which contains up to 531 ppm Li and forms narrow, ribbon-shaped dispersal train extending 4 km down ice. The results presented here demonstrate that drift prospecting methods are effective tools for Li exploration in southwestern Nova Scotia, as well as in other prospective glaciated regions in Canada.

---

### Depth controls on metamorphic decarbonation: insights from metacarbonate rocks from the Sierra Nevada arc, California, USA

---

BROOK PENNER AND ZHENHAO ZHOU

Department of Earth and Environmental Science,  
Acadia University, Wolfville, Nova Scotia B4P 2R6, Canada  
<0309030p@acadiau.ca>

Metamorphic decarbonation has been recently regarded as a source of carbon that may contribute to the global carbon cycle. This is attributed in part to the temporal coincidence between fluctuations in total length of continental arcs and climate intervals throughout geological history. Most field-based studies to date focus on metacarbonate rocks from shallow depths (<2 kbar equivalent), because thermodynamic modelling reveals that decarbonation reactions are suppressed at high pressures. However, this potential depth effect has not been revealed using natural samples. This research examines this gap by investigating metacarbonate rocks near the Sierra Nevada Batholith (SNB) in California, USA. The SNB intruded carbonate platforms at a range of depths causing metamorphism of the carbonate protolith from 2 kbar to at least 8 kbar. We hypothesize that carbonate protolith at mid-crustal levels experiences less decarbonation than its shallower equivalents, and that decarbonation reactions at different pressures occur under distinct physiochemical conditions. This hypothesis is being tested using petrography, bulk-rock chemistry, mass balance analysis and thermodynamic modelling. Preliminary results show that a sample formed at >8 kbar consists almost entirely of carbonates, whereas a sample formed at 2 kbar preserves only 10 vol. % carbonate minerals. This result suggests that higher pressure suppresses the extent of decarbonation. Ob-

served assemblages correspond to model conditions at 5 kbar and 2.5 kbar, with the higher-pressure assemblage indicating higher temperature and lower  $X_{\text{CO}_2}$  (i.e., more water-rich fluid). This suggests with increasing pressure, decarbonation reactions require higher temperatures and greater water availability to be triggered. These results and the ongoing mass balance work will provide new constraints on how pressure, temperature, and fluid availability regulate metamorphic decarbonation. Comparing the loss of carbon with that from other modern tectonic settings will enhance our understanding of how metamorphic decarbonation contributes to the global carbon cycle.

that altered the underlying basalt and led to hydrothermal zeolite precipitation. The thick upper flow unit re-activated hydrothermal flow through the Blomidon Formation and mixed with zeolite-rich fluids, producing agate with an outer rim of zeolite, in some cases reactivating or fragmenting earlier agate veins. Reactivated veins and deformed goethite flames demonstrate syn-emplacement tectonism. Oxidizing groundwater following Early Cretaceous inversion of the Fundy Basin led to alteration of original magnetite and hematite flames to goethite. The beauty and variety of Fundy agate were created by the specific details of the volcanic hydrothermal system.

---

### Mineralogy and origin of Bay of Fundy agate, Nova Scotia, Canada

---

GEORGIA PE-PIPER<sup>1</sup>, DAVID J.W. PIPER<sup>2</sup>, BRIAN HEBERT<sup>3</sup>,  
AND ANGELIKI PAPOUTSA<sup>4</sup>

1. *Department of Earth Science, Saint Mary's University,  
Halifax, Nova Scotia, B3H 3C3, Canada*

*<gpiper@smu.ca>*

2. *Geological Survey of Canada, Natural Resources Canada,  
Dartmouth, Nova Scotia B3B 1A6, Canada*

3. *Fundy Treasures Gifts and Tours, 107 Highway 242,  
Joggins, Nova Scotia B0L 1A0, Canada*

4. *Department of Geology and Geoenvironment,  
University of Athens, Athens 10679, Greece*

This study investigates the distinctive agates in the latest Triassic North Mountain Basalt and proposes a model for their formation. Ten representative agate samples were investigated by petrographic microscope, scanning electron microscope with energy dispersive spectroscopy analysis of mineral inclusions, and Raman spectroscopy analysis of silica and iron minerals. Agate veins typically show three zones: (1) an outer zone microgranular quartz and irregular goethite flames; (2) banded chalcedony; and (3) a central zone of coarse-grained polycrystalline quartz. Similar zonation is reported from many hydrothermal vein agates elsewhere, interpreted as forming in situ from an initial impure amorphous silica sol, by processes of self-organization and Ostwald ripening. The North Mountain Basalt in the Fundy Basin comprises >150 m thick lower and upper inflated flow units and a middle unit of multiple thinner flows. The base of the lava pile is saucer shaped. Hydrothermal circulation driven by the lower flow unit dissolved halite from the underlying Blomidon Formation and was hot enough (>300°C) to dissolve silica, copper, and iron. The resulting silica sol migrated up dip to the southern margin of the basalt, where actively deforming faults along the Minas Fault Zone provided pathways and accommodation. The thinner flows of the middle unit created smaller hydrothermal systems

---

### Interpreting coastal landscapes for the Cliffs of Fundy Geopark, Nova Scotia, Canada: new geoscience insights

---

DAVID J.W. PIPER

*Geological Survey of Canada, Natural Resources Canada,  
Dartmouth, Nova Scotia B3B 1A6, Canada*

*<david.piper@canada.ca>*

Interpretation of coastal landscapes and their impact on human activities is important for public outreach in the Cliffs of Fundy UNESCO Global Geopark. Landscapes are influenced by bedrock distribution, glaciation and post-glacial sea-level changes and many previous workers have evaluated their impacts. This contribution proposes new interpretations of some landscape features and attempts a coherent synthesis. Before the opening of the basalt barrier between Cape Split and Cape d'Or in the earlier Quaternary, first allowing marine transgression into Minas Basin, the meandering dry valley occupied by Highway 2 west of lower Five islands was cut by a west-flowing river that entered the Bay of Fundy near Port Greville. Only after the sea entered Minas Basin did the rivers draining the Cobequid Highlands have a direct southward route to the sea. Once breached, the basalt barrier was widened and deepened by glacial erosion. ice-streaming through Minas Passage is recorded in the roche moutonnee form of Partridge Island and Cape d'Or. Farther inland, slot canyons and incised meanders in south-flowing rivers were cut subglacially, not proglacially. ice retreat to the present coastline at 16 ka allowed raised beaches, wave-cut platforms, sea stacks and fluvioglacial terraces to form, up to 40 m above present sea level. Relative sea level fell progressively until the at -30 m lowstand at 8 ka, when Minas Basin was a river plain with incised river talwegs leading to a delta located in Minas Passage. During the subsequent rise in relative sea level, the old delta was re-worked into a barrier beach that blocked Minas Passage. A fresh or brackish lake occupied Minas Basin. At ca. 3400 BP, the barrier beach was breached in a storm and macrotidal marine waters first entered Minas Basin.

The gravel remnants of the beach are found at Diligent River and Black Rock.

MICHAEL POWELL AND JAMES BRENNAN

*Department of Earth and Environmental Sciences,  
Dalhousie University, Halifax, Nova Scotia B3H 4R2, Canada  
<michael.powell@dal.ca>*

---

**Arsenic accumulation in alder trees at a legacy gold mine tailings impoundment in southwest Nova Scotia, Canada: implications for environmental risk**

---

EVE POLE<sup>1</sup>, IAN SPOONER<sup>1</sup>, SARAH HINES<sup>2</sup>,  
AND NELSON O'DRISCOLL<sup>1</sup>

*1. Department of Earth and Environmental Science,  
Acadia University, Wolfville, Nova Scotia B4P 2R6, Canada  
<0300502p@acadiau.ca>*

*2. K.C. Irving Environmental Science Centre,  
Acadia University, Wolfville, Nova Scotia B4P 2R6, Canada*

Legacy gold mine tailings contaminated with arsenic pose significant ecological risks, particularly through bioaccumulation in terrestrial food webs. This study investigated patterns of arsenic accumulation in alder trees (*Alnus* spp.) growing on legacy gold mine tailings in southwestern Nova Scotia. Alder was selected as the target species due to its ecological importance as a food source, tolerance to disturbed environments, and to better understand their potential role in the phytostabilization of arsenic. Field sampling involved collecting alder tissues, such as leaves, catkins, and branches, as well as associated soil and water samples. Metal concentrations were analyzed using X-ray fluorescence (XRF) and inductively coupled plasma spectrometry (ICP-MS). Arsenic levels in tailings soil (430–3200 ppm) and associated groundwater (350–5600 ppb) exceeded baseline values and CCME guidelines. Arsenic levels in alder growing on tailings greatly exceed concentrations in non-tailings alder samples. Catkins on the tailings exhibited the highest arsenic concentrations (mean 1300 ppb; maximum 2140 ppb) and branches the lowest (mean 290 ppb; maximum 420 ppb). Relatively low concentrations in leaves and branches compared to tailings sediments and groundwater suggest that alder may contribute to arsenic phytostabilization at contaminated sites. Though alder is not considered to be an arsenic hyperaccumulator, our data indicates that elevated arsenic concentrations in catkins may pose a potential arsenic bioaccumulation risk for birds and other wildlife that consume reproductive tissues.

Recent market conditions, coupled with government investment and regulatory relief, have led to an exploration rush in Nova Scotia. Among the desirable units for exploration in Nova Scotia is the South Mountain Batholith (SMB), the largest extant granitic intrusion of the Appalachian orogen. However, despite the presence of a previously operating Sn mine and dozens of known mineralizations for critical mineral resources (Li, W, U, etc.), there have been no advanced operations developed in the SMB for over 30 years. To promote the discovery of new resources, we develop a mineral prospectivity map of the SMB using a rasterized cosine similarity approach. This approach leverages our newly determined mineral proportions and oxide trace element geochemical data, combined with previous measurements of whole rock geochemistry and biotite trace elements, to reveal the mineralogical and geochemical zonation of the SMB. Geochemical and mineralogical data are interpolated at both the batholith and pluton scale using an inverse distance weighting. Interpolated values at known mineralizations are used to calculate an average multivariate geochemical vector representing target pixels which are most likely to bear mineralization. Cosine similarity between this target vector and the interpolated vector at any given pixel quantifies their directional similarity in multivariate geochemical space, with the angular separation indicating (dis)similarity. The resulting prospectivity map highlights regions whose geochemical signatures closely resemble, or strongly contrast with, those of known mineralizations, thereby identifying areas that are highly prospective for new resources. Results are framed in the context of a cooling granitic magma experiencing fractional crystallization as the dominant differentiation mechanism.

---

**Designing a geoscience strategy board game**

---

SCOTT PRESTON AND KATIE McDONOUGH

*Department of Earth Sciences, University of New Brunswick,  
Fredericton New Brunswick E3B 5A3, Canada  
<spreston@unb.ca>*

Step into 19<sup>th</sup> Century Saint John and become a founding member of the Steinhammer Club. Play as ambitious young citizen-scientists curious to unravel the remarkable secrets hidden beneath their city's stony hills and jagged coastline. In this board game, you will acquire knowledge and collect fossils while exploring the world-famous geology of the Bay of Fundy and venture to other locales across the Maritime

---

**Mineralogical and geochemical zonation of the South Mountain Batholith, Nova Scotia, Canada with applications to mineral resource prospectivity by cosine similarity mapping**

---

Provinces. Race to collect sets of cards to give public lectures and publish scientific papers before your opponents. Befriend renowned scientists and help the museum build a collection of the fossil record from the Precambrian to the Quaternary. *Steinhammer: Dawn of Geoscience on Canada's East Coast* is a board game being developed by Resolve, the University of New Brunswick's applied game design research group led by Dr. Scott Preston. Resolve designs games that educate and inform. Since 2021, fifteen different students have participated. This project began in Fall 2024 after Dr. Preston met with Catrina Russell to discuss collaborating on a new science-focused game. Those early meetings identified the need for a light-strategy game aimed at older teens and young adults focused on geoscience literacy, preferably something that highlighted the incredible geology of the Saint John area. The story of the real-life Steinhammer Club (a group of young citizen-scientists based in the city in the 19<sup>th</sup> century) became the game's hook early on and the project took shape as not only a science-based game, but also, as the scope of the game changed to include NS and PEI, a game about the beginnings of geology and paleontology in Eastern Canada. Dr. Preston and Katie McDonough would be thrilled to bring a prototype of *Steinhammer* to AGS this winter, present a talk about the game, and provide gameplay opportunities for those in attendance.

---

**Metasomatic alteration around pegmatites in the Cat Lake–Winnipeg River pegmatite field, southeastern Manitoba, Canada: insights from exocontact cesium enrichment**

---

ZANE ALDEEN RAHABI<sup>1</sup>, DAVID LENTZ<sup>1</sup>,  
TANIA MARTINS<sup>2</sup>, AND LEE GROAT<sup>3</sup>

1. Department of Earth Sciences, University of New Brunswick,  
Fredericton New Brunswick E3B 5A3, Canada

<zanealdeen.rahabi@unb.ca>

2. Manitoba Geological Survey,

Winnipeg, Manitoba R3G 3P2, Canada

3. Department of Earth, Ocean and Atmospheric Sciences,

University of British Columbia, Vancouver,

British Columbia V6T 1Z4, Canada

The Cat Lake–Winnipeg River pegmatite field in southeastern Manitoba is prospective for many metals considered critical including Li, Cs, Ta, and Sn. Outside of known occurrences at the Tanco pegmatite and High-Grade dyke, recent exploration shows that pegmatites in this region do not contain significant Cs mineralization. However, core assay data show that ortho-amphibolite and mica-cordierite schist wallrock intervals around these Cs-poor pegmatites locally contain Cs concentrations up to 5100 ppm, and calibrated pXRF measurements on samples from these intervals yield Cs concentrations up to 2.3 wt%.

Detailed petrography reveals three wallrock alteration assemblages, from proximal to distal: (1) tourmaline + muscovite + apatite ± Li-Mn-Fe phosphates; (2) biotite ± holmquistite; and (3) epidote + chlorite ± calcite ± titanite ± holmquistite. A comparison of pXRF results and petrography shows a correlation between Cs concentrations and biotite modal abundance, with values for both peaking 20 cm away from the pegmatite-wallrock contact and decreasing to below detection limits beyond 60 cm. Micro-XRF analysis confirms that wallrock Cs is mainly contained within biotite. Alteration assemblages around the studied pegmatites strongly resemble those described by previous workers at the nearby Tanco pegmatite, which include B (± Li) tourmalinization, K-Rb-Cs-F (± Li) biotitization, and propylitic alteration (Hbl + Pl → Ep + Chl + Ttn + Cal + clay). Previous interpretations of these assemblages at Tanco suggest that they are caused by B- and F-rich fluids released during late pegmatite crystallization and enriched in Cs and other alkalis by sericitization of K-feldspar. Similar processes could therefore be responsible for Cs enrichment around these studied pegmatites, even in the absence of pegmatite Cs mineralization. However, the observed distribution and association among tourmaline, muscovite, apatite, and Li-Mn-Fe phosphates suggests that Li, P, K, and Rb behaved differently in these pegmatites than in the Tanco pegmatite system.

---

**Petrogenetic analysis of Neoproterozoic niobium, yttrium, and fluorine (NYF) pegmatites found in Perch Falls quarry, south-central New Brunswick, Canada: insight from petrology, lithochemical signatures, and zircon analysis**

---

JENNA REES, DAVID LENTZ, AND FAZI YOUSEFI

Department of Earth Sciences, University of New Brunswick,  
Fredericton New Brunswick E3B 5A3, Canada

<jrees3@unb.ca>

In the Perch Falls area of south-central New Brunswick, eight steeply dipping pegmatitic dykes strike NE–E that intruded the Neoproterozoic Perch Falls Granodiorite. These pegmatites represent a late-stage igneous event and are composed of quartz, plagioclase, K-feldspar, accessory phases, and locally quartz-rich cores. The presence of magmatic epidote and rare allanite in these calc-alkaline granitic dykes suggests crystallization near the wet solidus at moderate crustal pressures (>300 MPa) consistent with low normative quartz, with minor late-stage hydrothermal alteration. Whole-rock geochemistry shows the pegmatites have silica-rich compositions (SiO<sub>2</sub> = 71.1–77.6 wt.%), moderate Al<sub>2</sub>O<sub>3</sub> (12.3–15.6 wt.%), and elevated alkalis (Na<sub>2</sub>O = 2.65–4.12 wt.%; K<sub>2</sub>O = 3.01–5.62 wt.%), with elevated Nb–Y contents consistent with

a NYF affinity, but show variable Zr concentrations (23 to 420 ppm), reflecting low crystallization T (<650°C). Normative quartz calculations further support considerable magmatic fractionation. Overall geochemical signatures and normative values closely overlap those of the nearby Prince of Wales Granite, meaning these dykes likely were derived from it. MicroXRF reveals a few larger zircons (100–600 µm) present in the pegmatitic dykes. SEM-BSE analysis reveals these zircons are predominantly subhedral and weakly unzoned. These zircons locally exhibited metamictization and these zircons host micron-scale Th-rich inclusions (thorite, 5–10 µm), based on SEM-EDS. SEM-BSE imaging showed that zircons are present in the adjacent Prince of Wales Granite. Unlike the pegmatitic dykes, the zircons in the Prince of Wales Granite lacks thorite inclusions, are notably zoned, subhedral shaped, and are small-sized (5–100 µm). However, in both granitic units, these zircons preferentially occur within feldspar-rich domains. Previous regional geochronology constrains these units to the late Neoproterozoic, interpreted as the youngest magmatic event. The combined petrological, geochemical, and zircon textural evidence indicates magmatic fractionation with incompatible-element enrichment, and minor magmatic-hydrothermal modification within weakly evolved NYF-type pegmatites.

surface and dissipate heat, but where there is extensive salt withdrawal, e.g., in the Atholl Syncline, gradients reach over 25°C/km. Drilling costs with these temperature-depth profiles present economic challenges for closed-loop or induced-fracture power generation and it is unlikely that open-loop systems in Carboniferous sandstones are practicable, based on extensive core data and hydrocarbon production that requires fracking. In the Sable Island area, we provide similar animations of Triassic/Jurassic salt (Argo Formation) followed by late Jurassic/Early Cretaceous successions (Mic Mac and Missisauga formations) that were deposited by the Sable Island Delta across the continental shelf and in isolated shelf-margin depocentres above listric growth-faults (“expansion trends”). In this setting salt diapirs that have pierced the footwalls of listric faults to 4000–5000 m subsea have enhanced temperatures in overlying high-quality aquifers with geothermal gradients exceeding 30°C/km. Standalone geothermal production is unlikely, but integration of energy transition opportunities should be considered. Wind power and hydrocarbon exploration are currently being licensed offshore where stranded gas resources of ~2 TCF are well-established. Both basins have energy storage potential in salt diapirs. Government funding (~\$40 million) for gas, geothermal energy, and carbon storage research was announced in 2025.

---

**3D geocellular modeling linking sedimentary, structural, and diapiric architecture to geothermal potential: Cumberland Basin and Sable Sub-basin, Nova Scotia, Canada**

---

BILL RICHARDS<sup>1</sup>, FRASER KEPPIE<sup>2</sup>, HELEN CEN<sup>2</sup>, AND GRANT WACH<sup>1</sup>

1. Department of Earth and Environmental Sciences, Dalhousie University, Halifax, Nova Scotia B3H 4R2, Canada <billrichards888@hotmail.com>
2. Nova Scotia Department of Energy, Halifax, Nova Scotia B3J 3P7, Canada

We present structural-stratigraphic and geothermal geocellular models of the onshore, Paleozoic, Cumberland Basin and the offshore, Mesozoic, Sable Sub-basin. In each basin, we show 3D animations of sedimentary successions above autochthonous and allochthonous salt and relate geothermal power potential to the vertical extent of salt diapirism. The structural-stratigraphic models are constructed from 2D and 3D seismic data and petroleum wells, with outcrop data incorporated at Cumberland. Temperature is modeled using Fourier’s law at Cumberland and populated from wells at Sable. The Cumberland Basin is the highest ranked onshore basin in the province for geothermal energy. We model geothermal gradients as low ~15°C/km where high thermal conductivity Windsor Formation salt diapirs reach

---

**Quantitative assessment and risk analysis of geologic carbon storage potential in the Triassic-Jurassic Fundy Sub-basin, Atlantic Canada**

---

BILL RICHARDS<sup>1</sup>, FRASER KEPPIE<sup>2</sup>, HELEN CEN<sup>2</sup>, AND GRANT WACH<sup>1</sup>

1. Department of Earth and Environmental Sciences, Dalhousie University, Halifax, Nova Scotia B3H 4R2, Canada <billrichards888@hotmail.com>
2. Nova Scotia Department of Energy, Halifax, Nova Scotia B3J 3P7, Canada

Carbon storage potential in the Fundy Basin was addressed qualitatively in 2010 and 2015 based on legacy 2D seismic mapping and two offshore wells that penetrated saline aquifers near onshore emissions sites. In this presentation, we provide quantitative assessment of multiple scenarios in the Fundy Sub-basin: range ~25 Mt–3 Gt, with storage efficiencies of 0.5%–5%. We provide risk analyses of probability against confidence for each play element: capacity, injectivity, containment, and pressure-space. The Fundy Basin comprises three contiguous sub-basins formed during the rifting of Pangea. The Fundy Sub-basin is a half-graben with three components: a crestally faulted rollover anticline above the Grand Manan–Headlands listric growth-fault, a synclinal basin and a southeastern flank that rises to onshore outcrops. To the northeast, the Chignecto Sub-basin is

transitional to the Minas Sub-basin, which is flanked by the Cobequid–Chedabucto strike-slip fault system. To assess capacity, we built a geocellular model from published maps incorporating four formations. Triassic Wolfville and Blomidon alluvial-fluvial-aeolian-playa deposits are separated stratigraphically from similar Scots Bay clastics by Jurassic North Mountain basalts. Aquifer porosities were populated from sonic logs with multiple realisations dependent on lithofacies/diagenetic models. Mud-weights from drilling records and onshore aquifer studies indicate long-term pressure equilibration facilitated by extensive fracturing, particularly in basalts, which raises containment risk if 10 000 years sequestration is stipulated. Conversely, uncertainty in primary aquifer continuity leads to uncertainty in short-term injectivity and pressure-space, with risk of unacceptable pressure build-up and project shut-down. Furthermore, if suitable aquifers are confined to basin margins by basinward development of fine-grained playa facies and increased cementation with depth, then there are contamination risks to onshore freshwater aquifers and crestal containment risks. Near Cape Spencer O-79, basalts “sea-crop” above sandstones that could leak gaseous CO<sub>2</sub> transitioning from supercritical phase. Interpreted shallow faults near Chinampas N-37 risk similar cross-fault leakage of high-buoyancy gas to the seabed.

---

### Exploring the influence of rock geochemistry on biofilm composition in a lithium pegmatite system using metabarcoding

---

LAURA RIENDEAU, DONNELLY ARCHIBALD,  
AND JESSE McNICHOL

*Department of Earth and Environmental Sciences, St. Francis Xavier University, Antigonish, Nova Scotia B2G 2W5, Canada  
<x2020api@stfx.ca>*

Microbes inhabit almost every environment on Earth and their community compositions are highly dependent on abiotic environmental gradients. These abiotic environments include different rock types that can have very different chemical compositions. The objective of this research is to investigate the influence of rock geochemistry on distributions in microbial communities, with a focus on lithium, an important chemical element for decarbonization. Biological applications to mineral exploration are growing in popularity and will be a valuable tool as the demand for critical minerals increases. Microbial biodiversity is being investigated using metabarcoding of rock surface biofilms on outcrops in the Hayman Hill area located near St. Stephen, New Brunswick. The study area is located in the St. Croix terrane of Ganderia and the local geology consists of plutonic rocks, including mainly diorite, granodiorite and musco-

vite-biotite granite, and spodumene pegmatite dykes that intruded metasedimentary rocks. Small (<2cm diameter and <0.5cm thick) rock chips were collected from different rock types for 3-domain metabarcoding, a process that generates a total community fingerprint of eukaryotic and prokaryotic rRNA barcodes that can identify organisms down to the species level and determine their relative abundances. Absolute abundances per unit of exposed surface area were determined by normalizing to three internal standards. Preliminary results show a diverse biofilm community of prokaryotic and eukaryotic microbes and success of a novel DNA extraction method from rock chips. Current work is focused on comparing the rock surface metabarcodes to whole-rock geochemical data from the same samples. Taken together, this study uncovers the relationships between rock type, geochemistry, and microbial community composition on rock surfaces in the study area.

---

### Fundy Basin reservoir characterization for geological carbon storage in eastern Canada

---

COADY ROACH<sup>1</sup>, EMMA WEHBE<sup>1</sup>, EMILY THEBEN<sup>1</sup>,  
LAUREN WHEATON<sup>1</sup>, DAVE KEIGHLEY<sup>2</sup>, AND GRANT WACH<sup>1</sup>

*1. Department of Earth and Environmental Sciences,  
Dalhousie University, Halifax, Nova Scotia B3H 4R2, Canada  
<lauren.wheaton@dal.ca>*

*2. Department of Earth Sciences, University of New Brunswick,  
Fredericton New Brunswick E3B 5A3, Canada*

For a project titled “Basin Evaluation for Offshore Carbon Storage in Atlantic Canada”, in association with Memorial University, University of New Brunswick, and University of Waterloo, we have been completing work regarding reservoir characterization of Atlantic sedimentary basins. This study has three major focus areas, delineating the Mesozoic strata for candidate reservoirs for storage of CO<sub>2</sub>; macro-characterization of reservoirs in Nova Scotia and New Brunswick, micro-characterization of these reservoirs, and static and dynamic models incorporating the data to estimate parameters such as geobody analysis, porosity, and permeability for long term CO<sub>2</sub> storage. Fieldwork has been completed along the Bay of Fundy, collecting data on potentially suitable formations within the Fundy Basin. Through outcrop, wellbore, and analog analysis, geobody and architectural elements can be deduced for reservoir modelling, 3D reservoir facies can be defined, and lithofacies analysis can be correlated to subsurface formations, incorporating well log and 2D seismic data. Through field and lab work data collection, including measurements from scintillometers, permeameter, and drone mapping, key formations are characterized based on lithology and

stratigraphic relationships, gamma ray logs and elemental analysis (XRF, XRD), and permeability. Thin sections from samples collected in the field allow petrographic analyses to define details of reservoir mineralogical composition, texture, and heterogeneity of reservoir strata, while gross depositional environment mapping gives a broader view of the systems of formation. Macro- and micro-characterization will be input into static (stratigraphic) and dynamic (fluid flow) models of the representative storage reservoirs.

---

### Looking for a deeply buried new Walton deposit - across Iapetus and into the Carboniferous basins of eastern North America

---

PETER ROGERS

*Chavin Consulting, 10902 Highway 215, South Maitland, Nova Scotia B0N 2H0, Canada <chavin\_consulting@yahoo.ca>*

Historically the Maritime Provinces have the greatest number of mines per unit area and Nova Scotia the second greatest mineral production per square kilometre of all Canadian provinces. The lower Carboniferous carbonate-hosted barite/base metal/silver deposits are significant occurrences. A new Walton Mine (5 M tonnes of barite and 0.4 M tonnes silver) would have a present-day value in the range of over \$1B CDN. Current mineral development in lower Carboniferous rocks in Nova Scotia is roughly equivalent to where Ireland was some 25 years ago. The Navan array of ore bodies (\$12B in ground value) was discovered in the early 1970s as a near-surface deposit using standard geochemical techniques. The next discovery of a “new Walton” will undoubtedly involve innovative techniques to find buried deposits of ‘Irish-type’ that are typically found in clusters along identifiable lineaments at 8 km intervals. To date sustained activity in Ireland has outlined approximately 10 additional economic deposits with the more recently discovered ones being deeply buried. The lower Carboniferous basins in the Maritime Provinces have both complex structural geology and a deep masking clay overburden with 4 multi-directional glacial events that have limited exploration success. Enzyme leach studies in the Tennycape area revealed an impactful combination anomaly of significant size—larger than that of a buried porphyry Cu deposit in the Atacama Desert of Chile. The Halogen elements define a distinct complete low Cl, Br, Ba, Se, Ge, Ti + Mn anomaly seen as a 600 m × 400 m feature trending NW to SE. Ba occurs as a distinct trough, a possible fault/plumbing trace. The elements As, V, Cu, and Ga define a drill target whereas Pb, Zn, Cd, Co, and Ni occur in the halo. REE anomalies including La and Ce along with Th and U occur as a spatially coherent high in the west, possibly an intrusive body.

---

### Nova Scotia critical minerals opportunity

---

PETER ROGERS<sup>1</sup> AND ROBERT STEWART<sup>2</sup>

1. *Chavin Consulting, 10902 Highway 215, South Maitland, Nova Scotia B0N 2H0, Canada <chavin\_consulting@yahoo.ca>*

2. *FGC, Independent Geoscience Consultant, Bedford, Nova Scotia B4A 3Z2, Canada*

Rare earth elements (REEs) consist of 15 lanthanide elements along with yttrium and scandium and generally fall into 2 groups of light rare earths (LREE) or heavy rare earths (HREE). Their end use and largest demand is found in the world’s fastest growing markets: clean energy and high technology. The global REE market size is USD 3.74 billion in 2024 and expected to grow to USD 8.14 billion by 2032 - <https://www.fortunebusinessinsights.com>. Specifically, many REEs are expected to be in short supply in the next 15 years include neodymium, europium, terbium, dysprosium, and yttrium. Globally ionic Absorption (IA) REE deposits are the predominant source for heavy REE (HREEs) and yttrium with most exploited deposits confined almost exclusively to areas underlain by granitic rocks in southern China. IA ores are defined when weathered granites contain more than ~50% ion-exchangeable REY (REE + Y), relative to whole-rock REY. Present day ore grades range from 140 to 6500 ppm REY with the most valuable REE ores having significant HREE enrichment. For exploration purposes the weathering profile of the deposits can be divided into a REE-leached zone in the upper part of the profile, with a positive Ce anomaly, and a REE accumulation zone with the IA ores in the lower part of the profile that is characterized by a negative Ce anomaly. Within Nova Scotia the area with most REE potential is Cape Breton Island which contains several granitic bodies of interest especially in the Aspy terrane. Recent vegetation sampling has indicated areas of exploration interest for REE that are potential economic drivers for development in partnership with local indigenous groups.

---

### Evolving shorelines: update on Nova Scotia’s coastal monitoring program

---

ANNA RYAN

*Nova Scotia Department of Natural Resources, Halifax, Nova Scotia B3J 2T9, Canada <anna.ryan@novascotia.ca>*

Nova Scotia has a dynamic coastline extending over 13 000 km, boasting a wide range of shoreline types from rocky cliffs to sandy beaches. To monitor coastal change across this diverse environment, the Nova Scotia Geological Survey established a Coastal Monitoring Program comprised

of over 120 sites that are periodically surveyed using Remotely Piloted Aircraft (RPA) to collect high-resolution LiDAR and photogrammetry data. Comparison of annual surveys enables visualization and quantification of shoreline change, while analysis of historical aerial imagery provides context for longer term coastal evolution. Case studies were conducted for three separate monitoring sites to identify and interpret the relative influence of different factors driving coastal change, including the bedrock and surficial geology, geographic setting, coastal cell dynamics, and anthropogenic influences. Recognizing the complex, site-specific nature of coastal change along Nova Scotia's shoreline is fundamental to inferring future change and guiding shoreline management.

---

### Carbon sequestration and geological storage: Atlantic Canada's route to net-zero

---

SANA SALEHI, LAUREN WHEATON, AND GRANT WACH

*Department of Earth and Environmental Sciences,  
Dalhousie University 1355 Oxford Street,  
Halifax Nova Scotia B3H 4R2, Canada  
<lauren.wheaton@dal.ca>*

With federal and provincial emissions reductions goals set for 2030, the due date for progress is quickly approaching, requiring innovation and action to achieve targets and continue on the path to net-zero. Carbon capture and storage (CCS) is one such innovation with potential to reduce future emissions through utilization of existing geological resources in Atlantic Canada. By injecting CO<sub>2</sub> into suitable subsurface geological formations, long term sequestration of emissions can be achieved, reducing atmospheric and climatic impacts. Reservoir characterization projects are now underway to evaluate basin potential to store CO<sub>2</sub> in subsurface formations offshore Atlantic Canada. As potential for CCS grows, federal and provincial regulations must be reviewed and revised to clear the way for successful projects in Atlantic Canada.

---

### Subduction initiation within the Penobscot back-arc basin: evidence from the Great Bend Complex, central Newfoundland, Canada

---

GABRIEL SANTOS, IVAN YIP, AND HAMISH SANDEMAN

*Geological Survey of Newfoundland and Labrador,  
St. John's, Newfoundland and Labrador A1B 4J6, Canada  
<gabrielsantos@gov.nl.ca>*

The Great Bend Complex (GBC) is one three freq-

uently correlated ophiolitic complexes at the base of the Exploits Subzone, a major tectonic division in central Newfoundland. This presentation describes new whole-rock and mineral chemistry from five representative samples of dunite from the GBC, collected during the 2024 field season. The samples are medium-grained, strongly serpentinized, and consist of olivine, disseminated euhedral to subhedral Cr-spinel grains, and minor interstitial clinopyroxene. Whole-rock proxies are consistent with formation of the dunite via extensive melting of peridotite in the spinel stability field. The compositions of olivine (Fo<sub>90-92</sub>, NiO 0.1–0.35 wt%), clinopyroxene (Mg# ~95, TiO<sub>2</sub> <0.05 wt%, Al<sub>2</sub>O<sub>3</sub> ~1 wt%), Cr-spinel (Mg# 12–50, Cr# 60–80), and (Pt/Ir)<sub>N</sub> values are all consistent with the formation of dunite via a replacive reaction at low pressures between a depleted peridotite and an exotic percolating boninite melt. Platinum-group element data further corroborate the reactive origin of the GBC dunite. The record of boninitic magmatism, typically preserved in ophiolites formed in forearc settings, is inconsistent with prevailing tectonic models that assume the GBC represents a back-arc basin ophiolite. Our data suggest at least one previously unrecognized episode of basin inversion and subduction initiation within the Penobscot back-arc basin between its formation ca. 494 Ma and ophiolite obduction ca. 476 Ma. An episode of ridge inversion during induced subduction in the Penobscot back-arc basin would be consistent with the absence of a mature intra-oceanic arc and the short timeline between basin formation, inversion, boninite magmatism, and ophiolite obduction.

---

### Distribution and form of gold-silver in the Paleoproterozoic Lalor VMS deposit, Manitoba, Canada: preliminary analysis of precious-metal enrichment processes

---

MANJU SATI<sup>1</sup>, TANIA MARTINS<sup>1, 2, 3</sup>, CHRIS MCFARLANE<sup>1</sup>,  
DAVID LENTZ<sup>1</sup>, AND PURA ALFONSO<sup>4</sup>

*1. Department of Earth Sciences, University of New Brunswick,  
Fredericton New Brunswick E3B 5A3, Canada  
<manju.sati@unb.ca>*

*2. Manitoba Geological Survey, Winnipeg,  
Manitoba R3G 3P2, Canada*

*3. Department of Earth, Ocean and Atmospheric Sciences,  
University of British Columbia, Vancouver, V6T 1Z4, Canada*

*4. Department of Mining, Industrial and ICT Engineering,  
Universitat Politècnica de Catalunya Barcelona Tech (UPC),  
Barcelona 08034, Spain*

The Lalor base-metal deposit, in the Snow Lake area of the Paleoproterozoic Flin Flon domain, Manitoba, Canada,

is a structurally complex, gold-rich volcanogenic massive sulfide (VMS) system. The deposit is comprised of an auriferous Zn-Cu-bearing sulfide lenses, with variable gold-enriched zones, and enveloping hydrothermal alteration zones within amphibolitic host rocks. As the largest of the VMS deposits in the Snow Lake area the host rocks in Lalor are affected by syn-genetic alterations followed by intense polyphase deformation and amphibolite facies metamorphism. As a result, original ore mineral assemblages and individual phases are extensively modified, re-crystallized to coarser phases, and partly remobilized, such that syngenetic gold is remobilized locally leading to differential enrichment of Au-Ag. The lithologies, stratigraphy, and alteration patterns, supported with litho-geochemical analysis, of the Lalor VMS system have been investigated earlier; however, the trace-elements signature of ores and gangue minerals, especially in relation to gold enrichment, remains inadequately constrained. This study aims to systematically characterize trace-element distributions in various ore sulfides and associated gangue minerals from the Lalor VMS deposit to better understand their pre- to post-deformational form, distribution and saturation processes. This detailed approach helps to appreciate the post-ore metamorphic changes using combined analytical methods i.e., reflected-light petrography, micro-X-Ray fluorescence energy dispersive spectroscopy ( $\mu$ XRF-EDS), and scanning electron microscopy (SEM) coupled with EDS and backscatter electron (BSE) imaging, followed by laser ablation-inductively coupled plasma-mass spectrometry (LA-ICP-MS). LA-ICP-MS will quantify major- and trace-element concentrations in minerals from each assemblage, providing constraints on developing an understanding of critical trace element and gold incorporation into these phases, which helps to interpret early (syn-genetic) to late (deformation-related) fluid evolution and physicochemical conditions in this VMS system.

obscures the complex thermal history and vertical structure of Arctic soils, particularly a temperature-sensitive “transition zone” that experiences episodic freezing and thawing over decadal to centennial timescales as active layer thickness adjusts to shifts in climate, vegetation, and surface soil layers. identifiable by its high suspended (“ataxic”) ice content, this zone likely concentrates organic matter, mineral nutrients, and microbial assemblages percolated through soil water. As a result, the transition zone represents a disproportionately reactive substrate and a potentially important, underrepresented source of greenhouse gas emissions. We propose that applying change point detection to high-resolution radiocarbon measurements of bulk soil resolves five distinct layers—accumulation, mixing, transient, intermediate, and isothermal permafrost—that record thaw history and carbon dynamics within the upper two metres of the landscape. The active layer comprises an accumulation horizon, marked by nuclear bomb-spike carbon, and a homogenized mixing layer shaped by cryoturbation and inputs of modern dissolved organic matter and root exudates. Within the transition zone, the transient layer marks the maximum depth reached by the modern active layer each summer, whereas the intermediate layer stores millennial-aged carbon deposited during Holocene warm intervals. isothermal permafrost, isolated from the contemporary carbon cycle for millennia, exhibits substantially older radiocarbon ages and a more gradual radiocarbon depth profile. Today, gradual thaw as well as disturbances such as wildfires are causing ground-ice melt and transition zone collapse, exposing much older permafrost carbon to the atmosphere. Therefore, expanding depth-resolved datasets is essential to capture the full vertical structure of the soil-permafrost complex and improve projections of Arctic landscape stability and carbon release.

---

### Are we digging deep enough to get permafrost carbon?

---

MELISSA S. SCHWAB<sup>1</sup>, CLAUDIA I. CZIMCZIK<sup>2</sup>, SHAWN A. PEDRON<sup>2</sup>, AND JEFFREY M. WELKER<sup>3</sup>

1. *Department of Earth and Environmental Sciences, Dalhousie University, Halifax, Nova Scotia B3H 4R2, Canada*  
<melissaschwab@gmx.net>

2. *School of Physical Science, University of California, Irvine, California 92697-3100, USA*

3. *University of Alaska, Anchorage, Alaska 99508, USA*

Permafrost soils contain vast carbon reservoirs that emit greenhouse gases upon thaw. Yet current climate projections rely on simplified two-layer models distinguishing only the seasonally thawing “active layer” from underlying permanently frozen ground (“permafrost”). This framework

---

### Ichnofaunal diversity of the Pennsylvanian Clifton Formation, including the first documented trace fossil assemblage from within a hollowed lycopod stump, Clifton, New Brunswick, Canada

---

MARGARET SCOTT<sup>1,2</sup>, MATT STIMSON<sup>3,4</sup>, BAILEY MALAY<sup>4</sup>, BRIAN HERBERT<sup>5</sup>, OLIVIA KING<sup>6</sup>, LYNN DAFOE<sup>7</sup>, ROBERT GASTALDO<sup>8,9</sup>, JOHN CALDER<sup>3</sup>, MIKE RYGEL<sup>10</sup>, PAGE QUINTON<sup>10</sup>, LUCY SMITH<sup>11</sup>, KAYLA GERARD<sup>12</sup>, LUKE ALLEN<sup>12</sup>, KEENAN RICHARD<sup>13</sup>, SPENCER LUCAS<sup>14</sup>, AND HILARY MADDIN<sup>6</sup>

1. *St. Francis Xavier Alumni Association, Antigonish Nova Scotia, B2G 2X5, Canada* <margaretscott824@gmail.com>

2. *Department of Earth Sciences, Memorial University of Newfoundland, St. Johns, Newfoundland and Labrador A1B 3X5, Canada*

3. *Department of Earth Science, Saint Mary's University,*

Halifax, Nova Scotia B3H 3C3, Canada

4. *Geology and Paleontology Section, Natural Science Department, New Brunswick Museum, Saint John, New Brunswick, E2K 1E5, Canada*

5. *Fundy Treasures, 107 Highway 242, Joggins, Nova Scotia B0L 1A0, Canada*

6. *Department of Earth Sciences, Carleton University, Ottawa, Ontario K1S 5B6, Canada*

7. *Geological Survey of Canada, Natural Resources Canada, Dartmouth, Nova Scotia B2Y 4A2, Canada*

8. *Department of Earth Sciences, Colby College, Waterville, Maine 04901, USA*

9. *Department of Paleobiology, National Museum of Natural History, Smithsonian Institution, Washington, D.C. 20560, USA*

10. *Department of Earth and Environmental Sciences, State University of New York at Potsdam, Potsdam, New York 13676, USA*

11. *Zoology Section, Natural Science Department, New Brunswick Museum, Saint John, New Brunswick E2K 1E5, Canada*

12. *Department of Earth Science, University of New Brunswick, Fredericton New Brunswick. E3B 5A3, Canada*

13. *54 Cole Ave S, Moncton, New Brunswick E1C 8E8, Canada*

14. *New Mexico Museum of Natural History and Science, Albuquerque, New Mexico 87104, US*

The Maritimes Basin is renowned for its near-complete record of late Paleozoic tetrapod tracks (Early Carboniferous to early Permian). The Moscovian Clifton Formation in northern New Brunswick is well known for its paleobotanical record including in-situ lycopod trees. The lycopods are rooted in mineral paleosols, and were once hollowed-out stumps, episodically infilled with layers of sediment. Mud cracks, raindrop impressions and trace fossils are preserved on a bedding plane between strata that contain vertebrate and invertebrate body fossils. Preservation of traces within stumps are rare within the Paleozoic, with this report representing the first invertebrate assemblage, and second record of vertebrate ichnofossils, preserved inside a Pennsylvanian lycopod tree. The ichnoassemblage within the tree is diverse and dominated by invertebrate traces that includes the ichnogenera *Gordia*, *Helminthoidichnites*, *Helminthopsis*, *Circulichnus*, *Planolites*, *Cruziana*, *Striatichnium*, *Diplopodichnus*, and a single vertebrate attributed trace of cf. *Characichnos*, interpreted as a small tetrapod swim trace. The Clifton Formation also preserves a sparse ichnofaunal assemblage from the flanks of channel bodies/floodplain deposits and within abandoned channel deposits. Invertebrate ichnoassemblages (outside of lycopods) includes: *Diplichnites*, *Stiaria*, *Hexapodichnus*, *Cochlichnus*, *Planolites*, *Taenidium Gordia*, and *Helminthoidichnites*. The tetrapod track assemblage includes large *Characichnos* (swimming stem tetrapods), *Batrachichnus*, *Matthewichnus*, *Limnopus* (temnospondyls/microsaur), and *Hylopus* (anthracosaurs/reptiliomorphs). Trace fossils can be compared with body fossils of vertebrates and invertebrates also found preserved in the lycopods of the Clifton Formation; a unique

preservational occurrence. Trace fossils inside of a lycopod have implications for microhabitats in tree hollows within the Carboniferous and potentially expand the known paleobiodiversity of this time.

---

### Geochronological and microstructural analysis of the Cobequid–Chedabucto deformation corridor, Nova Scotia, Canada

---

DANA ŠILEROVÁ<sup>1</sup>, DAWN KELLETT<sup>1,2</sup>, SHAWNA E. WHITE<sup>1</sup>, CATHERINE MOTTRAM<sup>3</sup>, AND KYLE P. LARSON<sup>4</sup>

1. *Department of Earth Science, Saint Mary's University, Halifax, Nova Scotia B3H 3C3, Canada*

<dana.silerova@smu.ca>

2. *Geological Survey of Canada, Natural Resources Canada, Dartmouth, Nova Scotia B2Y 4A2, Canada*

3. *School of the Environment and Life Sciences, University of Portsmouth, Portsmouth PO1 3QL, UK*

4. *Earth and Environmental Sciences, University of British Columbia Okanagan, Kelowna, British Columbia C V1V 1V7, Canada*

The South Mountain Batholith (SMB), the largest composite granitoid body in the Appalachian orogen, outcrops over a large portion of central and western Nova Scotia. The SMB comprises 13 plutons, classified into generally barren Stage 1 and commonly mineralized Stage 2. The Stage 2 Davis Lake Pluton (DLP) of the SMB consists of several subplutons (East Kemptville, Dog Lake, Solomon Lake) contrasting in texture and composition, varying from muscovite leucogranite to leucomonzogranite to muscovite-biotite monzogranite. Metal enrichment varies within the DLP subplutons, with the East Kemptville subpluton hosting the world-class East Kemptville Sn greisen deposit, but additional Cu-Zn-In-Li-W resource potential is also recognized. No detailed chemical comparisons of these subplutons have been conducted to elucidate the reasons for their contrasting mineralization potential. In this study high spatial resolution microanalytical methods [laser ablation-inductively coupled plasma mass spectrometry (LA-ICP-MS), scanning electron microscopy, electron microprobe analysis] are combined with zircon host U–Pb age-dating and trace element analysis, and detailed petrography to quantify and compare the metal content and oxygen fugacity of magmatic liquids from the subplutons of the broader DLP of the SMB. Preliminary results of the LA-ICP-MS of silicate melt inclusions trapped within non-inherited zircon growth domains record significant compositional variations within, and between, the DLP subplutons. Large variations are observed in total alkali silica content, metal concentrations (e.g., Cu, Zn, Sn, W) and extent of differentiation, not observed previously in bulk rock analyses. Overall, the data

are consistent with the composition of silicate melt inclusions from other Stage 2 plutons within the SMB (e.g., New Ross, West Dalhousie Pluton). Notably, despite showing only subtle differences from other subplutons geochemically, silicate melt inclusions from the East Kemptville subpluton record some of the highest concentrations of Sn and W among the analyzed samples (up to ~500 ppm of Sn and W).

---

### **Integrating coastal environments into our understanding of major Mesozoic carbon cycle perturbations**

---

RICARDO L. SILVA

*Department of Earth Sciences and BETY Lab, Clayton H. Riddell  
Faculty of Environment, Earth, and Resources, University of  
Manitoba, Winnipeg, Manitoba R3T 2N2, Canada  
<ricardo.silva@umanitoba.ca>*

The Mesozoic witnessed several major carbon cycle perturbations associated with the deposition of organic-rich sediments in many locations worldwide. These episodes of environmental upheaval are commonly known as oceanic anoxic events (OAEs), often explained by drastic changes in carbon storage in the marine realm. However, there is a wide knowledge gap regarding the responses of terrestrial and coastal areas to the environmental drivers of OAEs and the impact of changing fluxes of carbon in and out of these depositional environments on the global carbon cycle. This presentation discusses the potential implications of terrestrial and coastal environments on global organic matter storage and atmospheric carbon feedback during the Mesozoic. The integration of our findings from both sides of the Atlantic margins with current modelling efforts will allow us to clarify the impact of coastal environments on climate, carbon-cycle perturbations, and OAEs, and to provide a Deep Time perspective on the factors contributing to present-day issues of coastal anoxia expansion and tidal wetland loss.

---

### **The Cambrian Hawke Bay Event of the eastern Laurentian margin in western Newfoundland, Canada**

---

NOAH SLANEY, DAVID LOWE, AND NICOLAS GOMEZ

*Department of Earth Sciences,  
Memorial University of Newfoundland, St. Johns,  
Newfoundland and Labrador A1B 3X5, Canada  
<Naslaney@mun.ca>*

The Laurentian Iapetan margin formed from the extension and breakup of the supercontinent Rodinia during

the Neoproterozoic and early Paleozoic. Strata deposited during late rifting through the onset of passive-margin sedimentation include the Labrador Group of the Laurentian autochthon in the Humber Domain of western Newfoundland. At the base of the Labrador Group, the Hawke Bay Formation is interpreted to represent a middle Cambrian (ca. 510–505 Ma) regional clastic regressive event termed the Hawke Bay Event (HBE). However, the tectonostratigraphic significance of this event along the Laurentian margin remains debated. To resolve details of the HBE and examine the effects of inherited rift structures on the tectonostratigraphic evolution of the Laurentian margin, facies and provenance analyses were conducted on the Hawke Bay Formation. Results of the facies analyses support the occurrence of a Cambrian Series 2 to early Miaolingian normal marine regression coinciding with a highstand systems tract followed by a falling-stage systems tract, possibly linked to post-rift isostatic rebound of the Laurentian margin following the opening of the Taconic Seaway. Highstand marine regression coincided with development of a wave-to-tide-dominated embayed clastic shelf succession, with progressive tidal reworking of quartz sand. This succession is capped by coastal aeolian facies and a deflation lag, comprising the falling-stage systems tract. Detrital zircon provenance suggests cratonic-scale sediment supply, but with a more significant contribution from Grenville sources than in the Hawke Bay Formation farther south in western Newfoundland, indicating sediment drainage along two separate sediment routing systems within the St. Lawrence Promontory. These separate routes were likely controlled by antecedent rift structures of a low-angle detachment rift system, which also likely increased tidal amplification along the regressed Laurentian margin.

---

### **A history mystery: “who the heck is Carl Steinhammer?!”**

---

LEA M.A. ST. JOHN AND CATRINA E. RUSSELL

*Department of Earth Sciences, University of New Brunswick,  
Fredericton New Brunswick E3B 5A3, Canada  
<catrina.elen@gmail.com>*

The Steinhammer Club of Saint John, New Brunswick, brought together a group of young men fascinated by the region's geological and palaeontological history. Among its members were Charles Frederic Hartt and George Frederick Matthew, both of whom went on to become celebrated geologists. Under the mentorship of John William Dawson, the Steinhammer Club evolved into the Natural History Society of New Brunswick. At this time, the original Steinhammer Club records were sealed, not to be opened again until after the deaths of all founding members. While this evolution expanded the study of New Brunswick's

natural history, one question has endured for more than a century - why was the name Steinhammer chosen in the first place? Some have suggested it refers to a geologist's primary tool, the "stone" or "rock" hammer. Others have jokingly proposed a connection to another geological pastime, indulging in steins of beer. Was the German word chosen to reflect respect for scientific scholarship in Germany at the time? Or was there an individual named "Steinhammer" whom the founders intended to honour? Our investigation into this history mystery began when our first author, Lea, discovered an 1863 article published by a "Carl Steinhammer." Assuming him to be an early member of the Steinhammer Club, she shared the reference with Catrina, whose immediate response was, "Who the heck is Carl Steinhammer?!" This talk follows our efforts to uncover the identity of Carl Steinhammer while tracing the origins of the Steinhammer Club's name. Ongoing digitization of archival evidence ultimately led to a decisive breakthrough, and we have been able to reveal a long-overlooked connection that demonstrates how a single forgotten name can unlock an entirely new chapter of the Steinhammer Club's story.

---

### Closure in concentration data can be measured in relative terms

---

CLIFF STANLEY

*Department of Earth and Environmental Science,  
Acadia University, Wolfville, Nova Scotia B4P 2R6, Canada  
<cliff.stanley@acadiau.ca>*

Closure poses interpretative challenges when evaluating compositional data as it forces component concentrations to sum to unity. This mathematical dependency contributes spurious variability to concentrations beyond those caused by material transfer. Fortunately, measuring the magnitude of closure's spurious variability is possible using the definition of a concentration:  $x = X/S$ , where  $x$  and  $X$  represent a component's concentration and amount in a system, and  $S$  represents the size of the system. By taking this equation's derivative, dividing both sides by  $x$ , and simplifying the result, we obtain a fundamental equation:  $dx/x = dX/X - dS/S$ , demonstrating that the relative variability of a component concentration ( $dx/x$ ) is a function of the relative variability of the component's amounts ( $dX/X$ ) and system sizes ( $dS/S$ ). Clearly,  $dX/X$  describes the relative amount of material transfer that took place. Thus,  $dS/S$ , the relative change in the size of the system, must be the amount of closure affecting the system's concentrations. Substituting a conserved component ( $Z$ ), where  $dZ = 0$ , for  $X$  in our fundamental equation yields:  $dz/z = -dS/S$ , so we must conclude that  $dS/S$  is a measure of the amount of closure, as  $Z$  undergoes no material transfer. Moreover, closure is a

function of the relative size of the system, so its effect is the same for every component. Examining concentrations of component ( $Y$ ) that were affected by material transfer ( $dY > 0$ ), we can also measure the relative amount of material transfer that took place ( $dY/Y$ ). Substituting the amount of closure ( $dS/S = -dz/z$ ) into our fundamental equation in  $Y$  and simplifying, we obtain:  $dY/Y = dy/y - dz/z$  and can now compare the magnitudes of closure and material transfer in compositional systems. Examples comparing the magnitudes of closure and material transfer are presented using lithochemical data from several different compositional systems.

---

### Nova Scotia hurricanes leave no Peggy's Cove boulders unturned

---

CLIFF STANLEY AND ANNA KIEFTE

*Department of Earth and Environmental Science,  
Acadia University, Wolfville, Nova Scotia B4P 2R6, Canada  
<cliff.stanley@acadiau.ca>*

Nova Scotia Prospectors Association participants on a recent fieldtrip to Peggy's Cove recognized a large, displaced boulder of biotite monzogranite near the lighthouse with a xenolith in a complementary position to a xenolith on a nearby outcrop. The similar lithology, size, orientation, and location of the xenolith on the boulder and outcrop confirmed the displaced boulder's origin. When the boulder moved 15.18 m (rotating 51°), either as a glacial erratic, or via meteorological forces (hurricanes, tropical storms, or "northeasters"), or both, was not clear until "time-bracketing" air photos proved the boulder moved between Jun. 2004 and Mar. 2009, likely during Tropical Storm Ophelia (2005-09-18), Hurricane Noel (2007-11-04), or Hurricane Kyle (2008-09-29). By fitting an irregular heptagon around the boulder, measuring its side lengths and orientations, and the boulder height at each vertex, the boulder volume was estimated (~128 m<sup>3</sup>). Multiplying this by the average biotite monzogranite density (2.70 kg/m<sup>3</sup>) yields a mass of ~346 tonnes, equal to a Boeing 777F (freighter) aircraft at take-off, a Union Pacific steam locomotive, an Eocene whale (*Perucetus colossus*, twice size of a blue whale), a 797B Caterpillar Mine Haul Truck payload, or 10 000 talents of gold (Chronicles 1, 29:7)! Furthermore, the work done by the waves to move the rock was ~3.62 megajoules, and the power expended was 0.52 megawatts, ~1/3 of a typical Nova Scotia windmill that can power 500 homes. We know precisely when smaller boulders (12.5 m<sup>3</sup>, 31.3 tonnes) at Peggy's Cove moved (Aug. 23, 2009,) during Hurricane Bill (between 1416 and 1418 hours) because of time lapse photography. Additionally, high-resolution satellite images track how other boulders south of the lighthouse have

moved “to” and “fro” via storms between 2015, 2021, and 2023. The masses of these boulders are stunning, providing ample appreciation for the power of Mother Nature’s fury.

---

**Provenance of metasedimentary units in the Laurentian Long Range Inlier, west Newfoundland, Canada: evidence of a shared history with the peri-Laurentian Dashwoods block**

---

CONNOR STEELE<sup>1</sup>, SHAWNA WHITE<sup>1</sup>, KEATON MARKHAM<sup>1</sup>,  
AND SEBASTIAN KOMMESCHER<sup>2</sup>

1. *Department of Earth Science, Saint Mary’s University,  
Halifax, Nova Scotia B3H 3C3, Canada*

*<Connor.Steele@smu.ca>*

2. *Department of Earth Sciences,  
Memorial University of Newfoundland, St. Johns,  
Newfoundland and Labrador A1B 3X5, Canada*

The Long Range Inlier (LRI) in the western Newfoundland (NL) Appalachians is a massif of Mesoproterozoic crystalline rocks correlated with the (ca. 1.0 Ga) Grenville Province of eastern Laurentia, uplifted to its current structural position during Devonian Acadian orogenesis. Recent exploration in the southern inlier has resulted in discoveries of high-grade gold occurrences, demonstrating significant economic potential. However, locating a deposit is impeded by lack of detailed geological mapping and geoscientific research within the inlier. This study, part of a larger mapping project aimed at resolving the inlier’s geological history, is focused on outcrop analysis and detrital zircon geochronology of metasedimentary units, traditionally grouped with Mesoproterozoic Grenville basement. Field results show metasedimentary units sit nonconformably atop (therefore younger than) 1.0 Ga crystalline basement. U–Pb detrital zircon ages range from Silurian to Paleoproterozoic and define a maximum depositional age (MDA) of  $705 \pm 15$  Ma, consistent with field observations. Lithology and age of these rocks point to correlation with the Paleozoic margin exposed throughout the Laurentian realm. Although younger than the MDA, a small Silurian population, with a ca. 428 Ma peak, is inferred as metamorphic due to a newly dated gabbro, which intruded the metasedimentary units. Major age peaks occur at ca. 1.0, 1.8, and 2.7 Ga, typical of Laurentian derived units; however, a significant proportion of Tonian (ca. 700–900 Ma) and ca. 2.0–2.2 Ga zircons are atypical and generally attributed to Gondwanan sourcing. Although not observed elsewhere in the NL Laurentian realm, these atypical ages are reported in inherited and detrital zircon populations in intrusive and Laurentian margin units, respectively, in the peri-Laurentian Dashwood terrane. These results not only aid in resolving the geological history of the LRI but demonstrate a connection between

Laurentian massifs and peri-Laurentian elements during Neoproterozoic to Cambrian margin formation which has implications in paleogeographic reconstructions of Rodinian rifting and Iapetan margin geometry.

---

**Strengthening geochemical provenance studies of pre-contact L’nu artifacts through the chemical homogeneity of natural copper samples at Cape d’Or, Nova Scotia, Canada, using laser ablation-inductively coupled plasma-mass spectrometry**

---

JODIE STEPHENS<sup>1</sup>, JACOB HANLEY<sup>1</sup>,  
CATHERINE COTTREAU-ROBINS<sup>2</sup>, AND TIM FEDAK<sup>2</sup>

1. *Department of Earth Science, Saint Mary’s University,  
Halifax, Nova Scotia B3H 3C3, Canada*

*<jodie.stephens@smu.ca>*

2. *Nova Scotia Museum of Natural History,  
Halifax, Nova Scotia B3H 3A6, Canada*

For thousands of years before European contact, the Wabanaki peoples - including the L’nu of the Maritimes Basin, used copper for sacred ceremonial gatherings such as the Mawiomí, gift and trade, and personal adornment as well as for functional objects including awls and fishhooks. Previous studies have found that Cape d’Or, Nova Scotia was likely a key source of native (elemental) copper for the Wabanaki peoples. In this study, microscopically-destructive laser ablation inductively coupled plasma-mass spectrometry (LA-ICP-MS) is used to further expand a large existing copper source database by analyzing the composition and homogeneity of native copper samples from the Cape d’Or underground workings at the historically mined Hanway lode, a location previously unstudied, as well as an additional location on Spencers Island near Cape d’Or. Analysis of a large number of copper fragments from a large volume of mined rock provides a means to determine how chemically homogeneous this source of copper is, thereby reducing uncertainty in provenance studies that aim to differentiate Bay of Fundy copper sources from other important copper procurement sites in eastern and central North America. Through the work of LA-ICP-MS the goal of this study is to ultimately answer key questions concerning the movement and trade of culturally and spiritually significant pre-contact artifacts of the Wabanaki peoples of the far northeast, thereby further strengthening future provenance knowledge in archaeological studies.

---

**Zircon geochronology and geochemistry in variably endowed subphases of the Davis Lake Pluton, South Mountain Batholith, Nova Scotia, Canada**

---

ANNA TEREKHOVA<sup>1</sup>, JACOB J. HANLEY<sup>1</sup>, AND JIM CROWLEY<sup>2</sup>

1. *Department of Earth Science, Saint Mary's University, Halifax, Nova Scotia B3H 3C3, Canada*  
<anna.terekhova@smu.ca>
2. *Department of Geosciences, Boise State University, Boise, Idaho 83725, USA*

The late Devonian South Mountain Batholith (SMB) consists of 13 plutons distinguished as early, barren Stage 1 or late, mineralized Stage 2 containing intrusion-related Sn-W-Mo-Cu-Zn-In-Li deposits. The Stage 2 Davis Lake Pluton (DLP) outcrops in the SW part of the SMB and is divided into 4 subplutons: main Davis Lake (“DL” leucomonzogranite), East Kemptville (“EK” leucogranite, hosting the EK Sn greisen deposit), Dog Lake (“DG” leucogranite) and Solomon Lake (“SL” muscovite-biotite monzogranite). The contrasting critical metal endowment of the subplutons is poorly understood, with the DL and EK subplutons being most prospective. Silicate melt inclusions in zircon show similar metal concentration ranges and extents of melt fractionation in all subplutons. Notably, Rb shows a positive correlation with Sn in melt inclusions from only the DL and EK subplutons, providing direct evidence that magma fractionation led to Sn enrichment. Ongoing characterization of zircon further elucidates the contrasting metal endowment and the evolution of the parental magmas. Through LA-ICP-MS analysis of subpluton zircons, combined with U–Pb age dating and detailed zircon petrography, we quantify and compare magma temperature and oxygen fugacity, and zircon trace element and metal concentrations (including REE, HFSE, and Sn–W) which constrain the degree of fractionation and magma metal enrichment. Additionally, we investigate the impact other crystallizing accessory phases have on zircon composition (e.g., titanite, apatite, monazite, etc.) and their relationship to processes controlling metal endowment. Preliminary results show that zircons from productive subplutons (EK and DL) record: (i) higher temperature ranges throughout their cooling history; (ii) lower values of magma  $fO_2$ ; and (iii) trace element ratios that record a disrupted fractionation trend, attributed to elevated concentrations of fluorine in the magmas that increased zircon and other accessory mineral solubilities preventing them from crystallizing in the later stages of the productive subpluton evolution.

---

**Defining the Geminid Deposit, northeastern Ontario, Canada: investigating mineral assemblages, textures, and Ni distribution**

---

EMILY THEBEN<sup>1</sup>, RICHARD COX<sup>1</sup>, JAMES BRENNAN<sup>1</sup>, AND MATTEA MCRAA<sup>2</sup>

1. *Department of Earth and Environmental Sciences, Dalhousie University, Halifax, Nova Scotia B3H 4R2, Canada*  
<emily.theben@dal.ca>
2. *Gold Candle Ltd., Toronto, Ontario M5J 2M2, Canada*

The Geminid Nickel Deposit was discovered in 2023 approximately 2 km east of the historic Kerr-Addison Mine, an underground gold mine that produced over 11M oz of gold from 1938 to 1996. Located in northeastern Ontario 40 km east of Kirkland Lake, the deposit sits along the Larder Lake - Cadillac Break (LLCB), within the Abitibi Greenstone belt, a region renowned for its mineral endowment. The initial discovery was made during exploration drilling along the LLCB, when the core intersected an approximately 23 m wide section of millerite mineralization with an average grade of 1.9% nickel. Follow-up drilling defined an approximately 300 × 700 m mineralized zone with vertical planar geometry. The Ni-ores are millerite-dominated and are hosted mainly in the ultramafic facies of the Larder Lake group, below the contact with Timiskaming sediments. At present, no comprehensive study has investigated the mineral assemblages, textures, or nickel distribution within the Geminid Deposit, limiting models describing its formation, potential economic value and Ni recovery. The aim of this study is to characterize the magmatic and secondary sulphide mineral phases and textures, evaluating their relationship to nickel distribution. Analytical methods include: (1) detailed petrographic analysis of magmatic and secondary mineral assemblages; (2) XRD analysis to determine mineral phases present; (3) SEM and  $\mu$ XRF for phase mapping and phase distributions at different scales within selected samples; (4) LA-ICP-MS analysis to investigate minor and trace element contents including PGEs. The results will be used to: (1) characterize ore types based on mineral assemblages, textures, and Ni tenor; (2) investigate factors governing nickel mobilization; (3) develop questions for further investigation. This study will provide a foundation for future genetic interpretations and economic evaluations. Beyond advancing the understanding of the Geminid Deposit, this research addresses a broader gap in knowledge of secondary (altered) sulfide systems and millerite dominated mineralization.

---

## A Late Carboniferous myriapod from the Joggins Fossil Cliffs UNESCO World Heritage Site, Nova Scotia, Canada

---

SYDNEY THOMPSON<sup>1</sup>, LEXIE ARNOTT<sup>1</sup>, AND JADE ATKINS<sup>2</sup>

1. *Department of Earth and Environmental Sciences,  
Dalhousie University, Halifax, Nova Scotia B3H 4R2, Canada  
<sy493765@dal.ca>*

2. *Joggins Fossil Institute, 100 Main St., Joggins,  
Nova Scotia B0L 1A0, Canada*

The Joggins Fossil Cliffs is a UNESCO World Heritage Site known for its Late Carboniferous fossils. The approximately 15 km coastal exposure preserves the fossiliferous ca. 315–310 Ma Joggins Formation, which was a tropical wetland-forest paleoenvironment. Approximately 200 different species have been discovered and identified from the Joggins Formation. Despite the high preservation potential in this formation, animal body fossils are still uncommon, which makes each body fossil an exciting find that allows us to learn more about Late Carboniferous animals. In 2016, a specimen, known as NSM016GF018.006, was discovered along the shoreline at the Joggins Formation. Specimen NSM016GF018.006 is an incomplete myriapod (an arthropod group that includes millipedes and centipedes) body fossil suspected to be a part of the Diplopoda class (millipedes) that is preserved in a concretion. During the middle- to Late Carboniferous Period arthropods were an abundant and diverse terrestrial species due to the many available niches on land, making NSM016GF018.006 an important addition when piecing together the Joggins Formation's paleoecology. The purpose of the present research is to (1) describe NSM016GF018.006 morphology, (2) compare NSM016GF018.006 morphology to other Late Carboniferous myriapods, and (3) investigate how NSM016GF018.006 fits into the Joggins Formation paleoenvironment. To accomplish this, NSM016GF018.006 was  $\mu$ CT-scanned in 2025 at  $\mu$ -VIS at Southampton University. The resulting 3D model was digitally segmented using Dragonfly's 3D analysis software. By using this non-destructive method, the digital segmentation of NSM016GF018.006 revealed morphological features that were hidden by the concretion's hard and difficult to prepare matrix. These features, such as tergites, diplosegments, and legs, will contribute to our understanding of NSM016GF018.006 morphology. Therefore, this research will also help to create a clearer understanding of myriapods during the Late Carboniferous and the paleoenvironments they inhabited.

---

## Provenance and tectonic evolution of the Meguma terrane, northern Appalachian orogen

---

CEES VAN STAAL<sup>1,2</sup>, SANDRA BARR<sup>3</sup>, CHRIS WHITE<sup>3</sup>,  
DEANNE VAN ROOYEN<sup>3</sup>, AND JOHN WALDRON<sup>4</sup>

1. *Geological Survey of Canada, Natural Resources Canada,  
Vancouver, British Columbia V6B 5J3, Canada  
<cees.vanstaal@NRcan-RNcan.gc.ca>*

2. *Department of Earth and Environmental Sciences,  
University of Waterloo, Waterloo Ontario N2L 3G1, Canada*

3. *Department of Earth and Environmental Sciences, Acadia  
University, Wolfville, Nova Scotia B4P 2R6, Canada*

4. *Department of Earth and Atmospheric Sciences,  
University of Alberta, Edmonton, Alberta T6G 2E3, Canada*

The Meguma terrane represents a piece of the Armorican terrane assemblage transferred to Avalonia during Cambrian to Ordovician rifting and opening of the Rheic Ocean or was accreted to East Avalonia as an isolated terrane or as a Gondwanan promontory following subduction of the Rheic Ocean. Early Devonian dextral translation moved the Meguma terrane opposite West Avalonia. Emsian to Famennian contraction between a cold and stiff promontory on West Avalonia in Nova Scotia and the Meguma terrane produced the Kejimkujic orogeny ( $D_1$ ), Kejimkujic orogenesis was first induced by a period of flat subduction of the Rheic Ocean beneath the Meguma terrane and composite Laurentia. Flat subduction inhibited formation of Early to Middle Devonian upper plate magmatism and placed the upper plate under compression, which led to folding. Contraction culminated in late  $D_1$  thrusting of the Meguma terrane over West Avalonia during Famennian indentation. Sinistral shear occurred in the northeastern part along ENE to NE-striking shear zones with dextral shear in the present southwestern part of the Meguma terrane and adjacent parts of West Avalonia. Progressive steepening of the Rheic slab in the Late Devonian re-established a metasomatized mantle wedge and formation of hydrous, subduction-related magmas, which assimilated significant amounts of Meguma terrane metasedimentary rocks that led to formation of peraluminous S-type magmas; more anhydrous A-type magmas formed in the distal backarc region where fluxing by hydrous slab-derived fluids was absent. Latest Devonian to Carboniferous translation of the Meguma terrane towards the west using the E-striking Minas Fault zone as a dextral transfer fault culminated in  $D_2$  sinistral oblique reverse slip along middle Carboniferous (Alleghenian) shear zones, which had progressively rotated anticlockwise into a NE-strike during oroclinal bending. This process led to further underthrusting of composite Laurentia beneath part of the Meguma terrane.

---

**Deconstructing the “Iapetus suture”:  
Diachronous accretion of terrane assemblages  
in the Appalachian–Caledonide collage**

---

JOHN WALDRON<sup>1,2</sup>, SANDRA BARR<sup>1</sup>, CHRIS WHITE<sup>1</sup>,  
DEANNE VAN ROOYEN<sup>1</sup>, AND PHIL MCCAUSLAND<sup>3</sup>

1. *Department of Earth and Environmental Sciences, Acadia University, Wolfville, Nova Scotia B4P 2R6, Canada*  
<waldronjwf@gmail.com>
2. *Department of Earth and Atmospheric Sciences, University of Alberta, Edmonton, Alberta T6G 2E3, Canada*
3. *Department of Earth Sciences, Western University, London, Ontario, N6A 3K7, Canada*

Maps of the Appalachian–Caledonide orogen have traditionally sought to identify a unique Iapetus suture marking either the collision between Laurentian and Gondwanan continental crust, or the closure of the last remnant of the Iapetus Ocean. This seemingly straightforward task is fraught with perils in an orogen that was built by the accretion of multiple fragments over substantial time. Orogen syntheses based in Britain and Ireland typically show the Iapetus suture as Silurian; those based in Newfoundland show a Late Ordovician suture; those in Cape Breton Island have no record of an Iapetus suture, whereas those in southern New England show the Iapetus closing in the Early Ordovician. The provenance and the timing of accretion can be examined using detrital zircon distributions and stratigraphic relationships. For example, the approach of a Ganderian terrane to the Laurentian margin is typically marked by an influx of ca. 1 Ga zircon from the Grenville orogen. The end of accretion is typically bracketed by an angular unconformity, above which forearc basin sedimentary and volcanic rocks contain both Laurentian and non-Laurentian zircon. This approach allows identification of multiple sutures arranged en echelon, ranging from Early Ordovician to Devonian. Terranes derived from peri-Gondwanan Ganderia arrived diachronously, such that the Laurentia–Gondwana boundary is marked by sutures of different age along the orogen. Traced along orogen strike, a suture of a given age may separate: (1) two Gondwana-derived terranes; (2) a Laurentia-derived terrane and a Gondwana-derived terrane; or (3) two Laurentia-derived terranes. Naming a single Iapetus suture is further complicated by recent results that show a second Devonian suture, within Britain, and other results that show the Munsungun inlier of northern Maine as Laurentian, not Gondwanan. We therefore argue that efforts to portray and discuss a single Appalachian–Caledonide “Iapetus suture” should be abandoned.

---

**Significance of the first documented Devonian  
sedimentary manganese mineralization in the Lower  
Seboomook Group in northern Maine, USA**

---

CHUNZENG WANG<sup>1</sup>, DAVID LENTZ<sup>2</sup>, AMBER WHITTAKER<sup>3</sup>,  
AND ROBERT MARVINNEY<sup>3</sup>

1. *Earth and Environmental Studies, University of Maine at Presque Isle, Presque Isle, Maine 04769, USA*  
<chunzeng.wang@maine.edu>
2. *Department of Earth Science, University of New Brunswick, Fredericton New Brunswick. E3B 5A3, Canada*
3. *Maine Geological Survey, 93 State House Station, Augusta, Maine 04333, USA*

The Devonian Lochkovian Seboomook Group, a flysch sequence of turbiditic slate and sandstone, is a major part of the Connecticut Valley–Gaspé Trough in northern–northwestern Maine. Recent mapping shows that the Seboomook rests unconformably on highly mineralized Cambrian–Silurian inliers, with this regional unconformity marking the end of the Salinic orogeny and the onset of Acadian foreland basin development. The lower Seboomook comprises a laterally continuous “trinity” of basal conglomerate, fossiliferous limestone and calcareous sandstone, and maroon-red slate-siltstone. The trinity locally contains multiple basalt flows erupted in a continental-arc setting, recording the onset of Acadian arc volcanism in northern Maine. Units previously mapped as separate or Silurian formations are now dated by conodonts as Lochkovian and assigned to the Seboomook Group; for example, the Frost Pond Formation of red slate northeast of Ripogenus Lake is reassigned to be the maroon-red slate-siltstone unit. This unit of hematitic iron formation is typically 10–15 m thick (locally up to 100 m) and indicates a critical early Devonian marine oxidation event. In 2025, a nearly 2-m-thick manganese-rich interval was discovered within this unit in a borrow pit near Hay Brook, in T7 R9 WELS township. The interval consists of alternating red, pink, and black Fe–Mn laminae of very fine-grained minerals; iron phases are predominantly hematite and minor magnetite and manganiferous phases dominated by Ca-rhodochrosite, with abundant cross-cutting rhodonite veinlets (generally <2 mm) and surficial black manganese-oxide films produced by supergene processes. Whole-rock analyses of two samples yield very high MnO content (45.8 and 47.1 wt.%) and low Fe<sub>2</sub>O<sub>3T</sub> content (3.09 and 2.97 wt.%). MicroXRF scanning confirms high Mn content. A slightly positive Ce anomaly and a convex-upward PAAS-normalized REY pattern support a hydrogenous origin for manganese mineralization. This occurrence, the “Hay Brook manganese occurrence”, represents the first documented Devonian sedimentary manganese mineralization in the northern Appalachian orogen.

---

## The re-established Lochkovian Seboomook Group and the onset of Acadian arc volcanism in the northern Maine Appalachian orogen

---

CHUNZENG WANG<sup>1</sup>, AMBER WHITTAKER<sup>2</sup>,  
AND ROBERT MARVINNEY<sup>2</sup>

1. *Earth and Environmental Studies, University of Maine at Presque Isle, Presque Isle, Maine 04769, USA*  
<chunzeng.wang@maine.edu>

2. *Maine Geological Survey, 93 State House Station, Augusta, Maine 04333, USA*

As a major part of the Connecticut Valley–Gaspé Trough, the Lower Devonian Lochkovian Seboomook Group is widespread in northern northwestern Maine. It is a flysch sequence of turbiditic slate and sandstone deposited in a foredeep due to advancing Acadian deformation. Recent detailed and reconnaissance mapping around Cambrian to Silurian Munsungun, Winterville, East Branch, Lunksoos Lake–Weeksboro, and Number Nine Mountain inliers demonstrates that the Seboomook Group rests unconformably on these inliers, with this regional unconformity marking the end of the Salinic orogeny and the inception of Acadian basin development. The lower Seboomook Group comprises a conformable, laterally continuous “trinity” of basal conglomerate, fossiliferous limestone and calcareous sandstone, and maroon-red slate-siltstone. The latter locally hosts laminated manganese-iron deposits. These units were previously mapped as separate or even Silurian formations but now are constrained by new conodont data to be Lochkovian and to be the lower succession of the Seboomook Group. Within the trinity, multiple basalt flows and minor mafic pyroclastic rocks (including the “West Branch Volcanics” redefined in our study) in the headwaters of the East Branch Penobscot and Aroostook rivers and near Ripogenus Lake provide a key record of early Acadian magmatism. Whole-rock analyses of 14 basalt samples reveal a remarkably homogeneous calc-alkaline geochemical signature characterized by pronounced negative Nb-Ta anomalies, enrichment in LILE and LREE, depletion in HFSE, low Nb/Y (average 0.22), and elevated La/Nb (average 2.58), Th/Nb (average 0.48), and Th/La (average 0.18) ratios, all typical of subduction-related continental-arc basalt. These data indicate that the basalts mark the onset of Acadian arc volcanism in earliest Devonian; this followed the final Salinic submarine mafic arc-like volcanism preserved in the ca. 425–424 Ma Spider Lake and Churchill Ridge formations in the Munsungun inlier, the rapid uplift and erosion of Salinic basement, and the earliest Acadian marine transgression in northern Maine.

---

## On the discovery of Triassic invertebrate and vertebrate ichnofossils from Melvin Beach, New Brunswick, Canada

---

EMMA M. WEHBE<sup>1</sup>, MATTHEW R. STIMSON<sup>2</sup>,  
COADY C. ROACH<sup>1</sup>, LAUREN E. WHEATON<sup>1</sup>, EMILY  
THEBEN<sup>1</sup>, DAVE KEIGHLEY<sup>3</sup>, PAUL OLSEN<sup>4</sup>, STEVEN J.  
HINDS<sup>5</sup>, ADRIAN PARK<sup>5</sup>, AND GRANT D. WACH<sup>1</sup>

1. *Department of Earth and Environmental Sciences, Dalhousie University, Halifax Nova Scotia B3H 4R2, Canada*  
<emma.wehbe@hotmail.com>

2. *Natural History Department, New Brunswick Museum, Saint John New Brunswick E2M 2K8, Canada*

3. *Department of Earth Sciences, University of New Brunswick, Fredericton, New Brunswick E3B 5A3, Canada*

4. *Earth and Environmental Sciences, Lamont-Doherty Earth Observatory, Columbia University, Palisades, New York 10964, USA*

5. *New Brunswick Department of Natural Resources, Fredericton, New Brunswick E3C 2G6, Canada*

The Fundy Basin of Atlantic Canada represents an aulacogen that preserves a thick succession of continental clastic fluvial and lacustrine deposits in both the Nova Scotian and New Brunswick coastlines. The basin accommodates sediments deposited between the latest Permian and the early Jurassic periods. To date, most paleontological research has been conducted in Nova Scotia, including skeletal remains of Canada’s oldest dinosaurs (Prosauropods), and abundant late Triassic and early Jurassic vertebrate ichnofossils, predominantly that of dinosaurs (i.e., *Otozoum*, *Grallator*, *Anomoepus*). Here, we present the first discoveries of vertebrate footprints, along with vertebrate and invertebrate burrows from the Echo Cove Formation at Melvin Beach New Brunswick, along the Fundy Trail Provincial Park. These traces represent the first evidence of vertebrate life from the upper Triassic (Carnian stage), and the first known Mesozoic vertebrate burrows in New Brunswick. The vertebrate tracks here are assigned to the ichnogenus *Chirotherium* ichsp, which others interpreted to be made by chirotheriid archosaurs, and to the ichnogenus *Characichnos* ichsp, assignable to swimming tetrapods. Here, we describe the first known occurrence of the ichnogenus (*Katarrhedites*) alongside several additional vertebrate burrows left in open nomenclature from the Triassic and noting the potential trace maker archosauriforms or another Triassic tetrapod. Invertebrate burrows such as *Taendium*, were observed within an isolated sandstone bed representing infaunal invertebrate deposit feeders. Trace fossils are here explored as an option to refine age constraints, where other conventional age-diagnostic information such as palynology or radiometric dates are limited or absent. Vertebrate footprint ichnobiochrons can be used based on global records of Triassic ichnofossils. In addition to the

systematic description of these new trace fossils, this study uses sedimentological and stratigraphic analyses of the Echo Cove Formation from data collected at coastal outcrops to better understand its depositional setting, that once supported the vertebrate and invertebrate activity.

---

**Microscale deformation and metamorphism of Neoproterozoic granitoid rocks in Stuart Mountain area, northern Caledonia Highlands, New Brunswick, Canada**

---

ALEX M. WELLES<sup>1</sup>, JOSEPH WHITE<sup>1</sup>, STEVEN HINDS<sup>2</sup>, CHRIS MCFARLANE<sup>1</sup>, AND ADRIAN PARK<sup>1</sup>

1. *Department of Earth Sciences, University of New Brunswick, Fredericton, New Brunswick E3B 5A3, Canada*  
<AlexWelles55@gmail.com>

2. *New Brunswick Department of Natural Resources, Fredericton, New Brunswick E3C 2G6, Canada*

Two suites of granitoid intrusions are emplaced into the Ediacaran volcanic-sedimentary succession in the northern Caledonia Highlands, New Brunswick: the Point Wolfe River Suite (ca. 625–615 Ma) and Bonnell Brook Suite (ca. 555–550 Ma). Both suites have been subjected to deformation commencing shortly after the intrusion of the Point Wolfe River suite and continuing episodically until Carboniferous time. Recently, the New Brunswick Geological Surveys Branch has undertaken re-mapping of this part of the Caledonia Highlands terrane with the aim of expand on the earlier published work using new exposures and U–Pb zircon dating. In aid of the latter, a detailed analysis of newly accessible outcrops in the Kent Hills Wind Farm (45.833297° N, 64.877544° W), currently included as part of the Bonnell Brook Suite, has been undertaken. This component of the project focuses on characterizing the intrusive body and its relationship to regional geology, relating the microstructure, both deformational and metamorphic/retrogressive, to evolution of the intrusion and U–Pb zircon and Rb–Sr white mica dating. Geochronological determinations are intended to constrain the protolith age, as well as establish more exact ages of deformation. The granodioritic host rock is characterized by calcic plagioclase, hornblende and quartz. Large relict hornblende grains are crossed by numerous K-feldspar veinlets. Subsequent deformation/metamorphism is exhibited by dynamic recrystallization by mixed-mode chemo-mechanical processes. Progressive deformation is intimately linked to hydration and introduction of potassium and silica-rich fluids, the latter being inferred from K-spar and quartz veinlets, the formation of muscovite-dominated deformation zones and striking rutile to titanite reaction textures. Though these observations this project aims to better understand the hydration and fluid migration occurring in these granit-

oids as well as the relative and exact timing of subsequent deformation in the area.

---

**Sharing Earth observation satellite technology and Indigenous mapping knowledge: recent Canadian Space Agency training activities in Atlantic Canada**

---

DIRK WERLE, GUY AUBÉ, JONATHAN LAJOIE,  
DALIA LAOUBI, AND JANIN HUARD

*Canadian Space Agency, Longueuil, Quebec J3Y 8Y9, Canada*  
<dirk.werle@asc-csa.gc.ca>

This presentation highlights scope, content and outcome of recent Earth Observation (EO) training and capacity building activities of the Canadian Space Agency (CSA) in Atlantic Canada. Planned for and with Indigenous Peoples, this work was based on guiding principles of Integrity, Reciprocity, Respect and Interaction, involving more than 50 participants on three occasions in 2024 and 2025. Combining presentations, discussions, field trips and ceremonies, we explored aspects of land use, environmental monitoring and issues of coastal and ocean governance through an interdisciplinary approach, with particular attention to Indigenous needs, perspectives, and knowledge systems. The value of satellite observations of coastal and marine environments, for example from Canada's RADARSAT Constellation Mission, was recognized as an integral part of the two-eyed-seeing and two-way learning approach. In practical terms, we introduced remote sensing concepts and examples of how optical and radar images can be a resource for participants interested in coastal areas, marine environments, impact of wildfires on land and air quality, mapping flood extent, and monitoring land use and land cover change. Hands-on activities and discussion enhanced the learning experience involving comparisons of detailed air photo series and synoptic satellite imagery. Overall training objectives and expected outcomes included the following: build awareness of the application potential of satellite-based EO; understanding the principles of remote sensing; accessing and applying satellite data; and engaging in respectful dialogue and two-way learning. The presentation includes Indigenous and space-technological perspectives on the-view-from-above with examples and comparisons of traditional geospatial knowledge and space technology. They explore the use of century-old traditional mental maps and recent satellite imagery for orientation, navigation and spatial awareness building. The presentation also highlights ongoing CSA collaborations and partnerships with indigenous Peoples and the international Committee on Earth Observation Satellites (CEOS) and its Working Group on Capacity Building and Data Democracy.

---

**Seeing inside basal Windsor Group microbial reefs in the Sydney Basin, Nova Scotia, Canada: serial-slab-based 3D reconstruction\***

---

KAYLA YOUNGCHIEF<sup>1,2</sup>, GRAYDON MACISAAC<sup>1</sup>,  
ZAILYNNE DURANT<sup>1,2</sup>, AND EDWARD J. MATHESON<sup>1</sup>

1. *Department of Math, Physics and Geology, Cape Breton University, Sydney, Nova Scotia B1M 1A2, Canada  
<kaylamaccormack11@gmail.com>*
2. *Department of Earth and Environmental Sciences, Acadia University, Wolfville, Nova Scotia B4P 2R6, Canada*

In the Glen Morrison area of the southern Sydney Basin, the basal Windsor Group Gays River Formation contains a prolific microbial reef complex up to 30 m thick that extends over >10 km<sup>2</sup>. Microbial reefs of similar scale are rare in the Phanerozoic rock record, and their internal architecture and growth mechanisms remain understudied. The exceptional preservation of the Glen Morrison reef complex offers a valuable opportunity to investigate microbial reef growth at the centimetre to decimetre scale, the scale at which reef architecture is constructed. This study presents the development of 3D mesoscale reconstructions using serial-slab-based digital reconstruction techniques. Selected specimens were sectioned with closely spaced cuts using a 24-inch slab saw. Each exposed slice was digitally imaged, processed to distinguish microbial fabrics, cavities, and cement, and imported into the SPIERS (Serial Palaeontological Image Editing and Rendering System) software package for alignment, segmentation, and 3D rendering. Through iterative methods development, the most detailed dataset was created with 26 slices at a spacing of 2.974 mm. While a primary challenge was achieving sufficiently small slice spacing for optimal 3D interpolation in SPIERS, image stacks, blocky 3D models, and downsized slice images enabled meaningful interpretation of microbial growth. The data reveal reefs built from irregular, interconnected microbial clots (thrombolites), with discrete growth events locally producing pseudo-layered or stromatolitic textures. Persistent clot growth generated vertically continuous elements with branching, dendritic-like geometries. Coupled with the absence of macroscopic biota, this suggests growth driven by localized sub-cm-scale microbial activity wherever conditions permitted. Persistence over broad areas and long timescales produced reef-scale structures, with early cementation contributing to framework integrity. This suggests a lack of self-sustaining ecological feedback characteristic of metazoan (e.g., coral) reefs, which engineer their environment to promote further growth. This indicates that the Glen Morrison reefs represent environmentally constrained, opportunistic accumulations.

\*Winner: AGS Rob Raeside Award for best undergraduate poster presentation

---

**Petrogenesis and geochemical evolution of Late Devonian peralkaline A-type granites in southwestern New Brunswick, Canada: insights from the Welsford, Parks Brook, and the Jake Lee alkali granites**

---

FAZILAT YOUSEFI<sup>1</sup>, DAVID R. LENTZ<sup>1</sup>,  
AND KATHLEEN G. THORNE<sup>2</sup>

1. *Department of Earth Sciences, University of New Brunswick, Fredericton, New Brunswick E3B 5A3, Canada  
<fazilat.yousefi@unb.ca>*
2. *New Brunswick Department of Natural Resources, Fredericton, New Brunswick E3C 2G6, Canada*

In southwestern New Brunswick, the Early Devonian 412±1 Ma Welsford (formerly 422 ± 1 Ma), 416 ± 5 Ma Parks Brook, and 418 ± 2.3 Ma Jake Lee (U–Pb zircon ages) peralkaline granites intrude along a NE-trending paleosuture zone. They exhibit an A-type affinity with variable degrees of mineralization, they are composed of quartz (15–20%), alkali feldspar, and plagioclase (40–75%), minor biotite, accessory zircon and apatite, and locally abundant riebeckite (10–20%). The McKeel Lake pegmatite-aplite dike swarm intruded the Welsford alkali granite and the dykes are notable for their Zr-REE-Y-Nb enrichment. The ratio of total alkalis (Na<sub>2</sub>O + K<sub>2</sub>O) to SiO<sub>2</sub>, together with major-element compositions (FeO<sup>T</sup> ≈ 1.5–9.4 wt.%, Al<sub>2</sub>O<sub>3</sub> ≈ 10.6–15.0 wt.%, and FeO<sup>T</sup>/MgO ratio ≈ 3–161) provides insights into parental magma sources, melting depth, tectonic setting, and mineralization potential. Geochemical data, including assessment files and whole-rock analyses, indicate anomalous high field strength elements, and REE enrichment. Rittmann serial (3.5 < σ < 8.8, and σ ≥ 8.8) and Apatitic indices (AI > 1) indicate that Welsford, Parks Brook, and Jake Lee granites exhibit alkaline features. The A-type within-plate tectonic setting of these intrusions is supported by Y + Nb > 60, Ta + Yb > 6, Zr + Nb + Ce + Y > 450, Zr > 688, Th/U > 5, Th > 20, and Eu/Eu\* = 0.26. Values of Ga/Al > 0.00025, Ce (> 100 ppm), Nb/Y (~0.2–0.5), Y (~105 ppm), and Nb (~33.5 ppm) indicate that the Welsford, Parks Brook, and Jake Lee intrusions formed in intraplate settings via low-degree partial melting of an enriched mantle, followed by fractional crystallization near the fayalite-magnetite-quartz buffer, with possible crustal interaction. The studied samples from these peralkaline granites range from A1-type, formed by highly fractionated mantle-derived melts, to A2-type granites, likely generated by partial melting of crustal material (under higher fO<sub>2</sub> conditions), as indicated by Nb/Y ratios ≤ 0.8.

---

**First intrusion-related Mo-Bi-Te-Au polymetallic system recognized in Chilean Patagonia**

---

MARCOS ZENTILLI<sup>1</sup>, JACOB HANLEY<sup>2</sup>,  
AND JUAN VALDÉS-EDWARDS<sup>3</sup>

1. *Department of Earth and Environmental Sciences,  
Dalhousie University 1355 Oxford Street,  
Halifax Nova Scotia B3H 4R2, Canada  
<marcos.zentilli@dal.ca>*

2. *Department of Earth Science, Saint Mary's University,  
Halifax, Nova Scotia B3H 3C3, Canada*

3. *Juan Valdés y Asociados, Santiago, Chile*

The Trapananda prospect in Rio Soler, in Aysén, Patagonia, latitude 47° S, Chile, contains quartz-Mo-Bi-Te-Au vein arrays within and near a granite apophysis of the North Patagonian Batholith that intruded pre-Jurassic carbonaceous phyllite and schist. Its mineralogy, comprised of molybdenite, native bismuth, and Bi-Te-S sulphosalts, is unknown elsewhere in Chile, and not compatible with the original description of the prospect as a porphyry-type Mo deposit. The 143.6 Ma (latest Jurassic) Re-Os date of molybdenite confirms a genetic association with the adjoining batholith. Two-phase fluid inclusions in quartz show no evidence of boiling (phase separation) but are coeval with molybdenite, Bi-Te-S sulphosalts minerals, and trails of Bi-Te-S semi-metal melt

inclusions. The fluid inclusions have an average salinity of 13.4 wt.% eq. NaCl and Th of 210°C and were trapped between ~270 and ~400°C at a minimum P of 0.7–2 kbar (min ~2.5–7 km depth). Together with the fluid inclusion systematics, the geochemistry of the igneous rocks, the vein textures, mineralogy and metal association (Mo-Bi-Te-Au) including semi-metal melts (which may have played a role in concentrating Au), and the stable isotope systematics (quartz  $\delta^{18}\text{O}$  +10 ‰; sulphide-sulphosalt  $\delta^{34}\text{S}$  -0.1 to -1.0 ‰; host schist  $\delta^{13}\text{C}$  = -22‰) are most compatible with metasediment-hosted, reduced-intrusion-related Au system (RIRGS). However, some features are not consistent with RIRGS, thus an intrusion-related polymetallic type of mineralization is favoured as a more general classification. The lack of native Au, which is contained in Bi-Te-S sulphosalt minerals as “invisible gold”, may explain why the deposit eluded discovery by traditional prospecting (panning) methods in this remote, glaciated region. In Atlantic Canada, despite a different tectonic setting, known intrusion-related Mo-Bi associations are recognized (e.g., Deep Cove and Long Lake, Nova Scotia) where detailed mineralogy and assaying have not been investigated with modern methods, and sulphosalts could hide valuable surprises.